
URINARY TRACT INFECTION IN FULL-TERM NEWBORN INFANTS: RISK FACTOR ANALYSIS

Mário Cícero Falcão, Cléa Rodrigues Leone, Renata A. P. D'Andrea, Roberta Berardi, Nilce A. Ono and Flávio Adolfo Costa Vaz

RHCFAP/2997

FALCÃO MC et al. - Urinary tract infection in full-term newborn infants: risk factor analysis. *Rev. Hosp. Clín. Fac. Med. S. Paulo* 55 (1):9-16, 2000.

SUMMARY: Objective: To analyze the correlation of risk factors to the occurrence of urinary tract infection in full-term newborn infants.

Patients and methods: Retrospective study (1997) including full-term infants having a positive urine culture by bag specimen. Urine collection was based on: fever, weight loss > 10% of birth weight, nonspecific symptoms (feeding intolerance, failure to thrive, hypoactivity, debilitate suction, irritability), or renal and urinary tract malformations. In these cases, another urine culture by suprapubic bladder aspiration was collected to confirm the diagnosis. To compare and validate the risk factors in each group, the selected cases were divided into two groups: Group I – positive urine culture by bag specimen collection and negative urine culture by suprapubic aspiration, and Group II – positive urine culture by bag specimen collection and positive urine culture by suprapubic aspiration .

Results: Sixty one infants were studied, Group I, n = 42 (68.9%) and Group II, n = 19 (31.1%). The selected risk factors (associated infectious diseases, use of broad-spectrum antibiotics, renal and urinary tract malformations, mechanical ventilation, parenteral nutrition and intravascular catheter) were more frequent in Group II (p<0.05). Through relative risk analysis, risk factors were, in decreasing importance: parenteral nutrition, intravascular catheter, associated infectious diseases, use of broad-spectrum antibiotics, mechanical ventilation, and renal and urinary tract malformations.

Conclusion: The results showed that parenteral nutrition, intravascular catheter, and associated infectious diseases contributed to increase the frequency of neonatal urinary tract infection, and in the presence of more than one risk factor, the occurrence of urinary tract infection rose up to 11 times.

DESCRIPTORS: Urinary tract infection. Risk factor. Newborn infant.

Usually, urinary tract infection in the neonatal period manifests after 72 hours of life. The incidence is between 0.1 to 10% in all infants. It is more frequent in male and preterm newborn infants, and may be as high as 10% in low birth weight infants^{17,20}. This incidence is directly proportional to the presence of several risk factors for this disease. Among the risk factors, beside those previously mentioned, are the presence of associated infectious diseases (sepsis, bronchopneumonia, septic arthritis, osteomyelitis, meningitis, etc.), the use of broad-spectrum antibiotics, mechanical ventilation, parenteral nutrition, intravascular catheter, and prolonged stay in the inten-

sive care unit^{17,20}. Renal and urinary tract malformations may be present in neonates with a urinary tract infection⁸.

Urinary tract infections are of interest to neonatologists for several reasons. First, they can be asymptomatic, so the diagnosis must be made by the examination and culture of a properly obtained specimen of urine^{7,11}. Second, they may indicate a serious underlying abnormality of the urinary tract, such as obstructive uropathy. Third, they

may have long-term consequences insofar as they may produce kidney damage, which may lead later in life to hypertension, recurrent infections, and kidney failure.

The knowledge of neonatal urinary tract infection risk factors or predisposing factors is important for identifying appropriate situations for performing serial urine culture to obtain an early diagnosis and adequate treatment¹³. The knowledge of the most important risk factors in a neonatal lower risk population is also important.

The purpose of this study was to analyze the contribution of the risk factors to the occurrence of urinary tract infection in full-term newborn infants.

From the Department of Pediatrics, Hospital das Clínicas, Faculty of Medicine, University of São Paulo.

PATIENTS AND METHOD

We conducted a retrospective study in the Child Institute “Prof. Pedro Alcântara” of the Hospital das Clínicas, University of São Paulo, Faculty of Medicine, including all full-term newborn infants (gestational age – 37 to 42 weeks) presenting positive urine culture by bag specimen collection.

Urine collection was based on clinical data as follows: fever ($T > 37.8^{\circ}\text{C}$)²⁴, weight loss > 10% of birth weight, or nonspecific symptoms (feeding intolerance, failure to thrive, hypoactivity, debilitate suction, irritability). In these cases, another urine culture was collected by suprapubic bladder aspiration to confirm diagnosis.

To compare and validate the risk factors in each group, the selected cases were divided into two groups: Group I – positive urine culture by bag specimen collection and negative urine culture by suprapubic aspiration, and Group II – positive urine culture by bag specimen collection and positive

urine culture by suprapubic aspiration. The selected risk factors were: associated infectious pathologies, use of broad-spectrum antibiotics, renal and urinary tract malformations, mechanical ventilation, parenteral nutrition, and intravascular catheter^{3,6,17}.

Figure 1 shows the design of the study.

The finding of 100 000 CFU/ml of the same microorganisms defined the positivity of the bag specimen collection²⁶. A suprapubic aspiration specimen was defined as positive at any CFU/ml rate²⁶.

The population of study was characterized by gestational age, birth weight, sex, perinatal asphyxia (Apgar score < 6 at 5 minute), and membrane rupture time (<24 hours and ≥ 24 hours). Symptoms were: fever ($T > 37.8^{\circ}\text{C}$), weight loss > 10% of birth weight or nonspecific symptoms (feeding intolerance, failure to thrive, hypoactivity, debilitate suction, or irritability).

The technique of bag specimen col-

lection was: antisepsy and sterile plastic bag fitted on the genitalia²⁵. If diuresis would not occur within 30 minutes, procedure would have been repeated until success²⁵.

The suprapubic aspiration specimen required a full bladder^{16,23}. The overlying skin was disinfected, the bladder was punctured above the symphysis pubis with a 25-gauge needle on a syringe, and about 2 ml of urine was aspirated^{2,21}.

All urine specimens were transported promptly to the laboratory⁹. Urine specimens were processed onto CLED, MacConkey, Thayer-Martin, thioglycolate, and blood agar plates. Culture plates were incubated at 35–37° C and read at 24–48 hours; if results were positive, the antibiogram was determined⁹. The absence of microorganisms within 48 hours characterized the culture as negative⁹.

Data were analyzed through chi-square, Fischer exact test, Student *t* test, and relative risk. Significance was defined as $p < 0.05$.




Figure 1 - Design of the study.
UC = urine culture

Table 1 - Population of the study – Group I and Group II.

	Group I n=42	Group II n=19	P	Test
Birth weight (g)	Average 3399.52 Variation 2560-4270 SD 418.36	3171.05 2420-4260 515.08	0.07	Student <i>t</i>
Sex	n (%)	n (%)		
Male	16 (38.1)	10 (52.6)	0.28	chi-square
Female	26 (61.9)	9 (47.4)	0.28	chi-square
Asphyxia Apgar 5' < 6	1 (2.4)	3 (15.8)	0.14	chi-square
Membrane Rupture time ≥ 24h	7(16.7)	5 (26.3)	0.47	chi-square

Table 2 - Clinical data – Group I and Group II.

	Group I N=42	Group II n=19	P	Test
	n (%)	n (%)		
Fever (T > 37.8°C)	38 (90.5)	15 (78.9)	0.31	chi-square
Weight loss > 10% of birth weight	20 (47.6)	3 (15.8)	0.01	chi-square
Nonspecific symptoms	4 (9.5)	10 (52.6)	0.0004	chi-square




Figure 2 - Clinical data (%) – Group I and Group II.

RESULTS

Sixty-one full-term newborn infants were included in this study, representing 5.1% of the total infants born alive in the study period. The diagnosis was confirmed (positive urine culture by suprapubic aspiration) on 19/61 (31.1%) of full-term infants included in the study (1.6% of the total infants born alive).

The infants were distributed in two groups according to the positivity of urine culture collected by suprapubic aspiration. Group I had 42 infants (68.9%) and Group II had 19 infants (31.1%).

The population of the study is shown in the Table 1 and clinical manifestations are shown in Table 2 and Fig. 2.

No complication occurred from specimen collection by the suprapubic method.

The microbial content of cultures of urine specimens was: *Escherichia coli* (42.9%), *Staphylococcus aureus* (10.5%) *Staphylococcus coagulase - negative* (26.3%), *Enterococcus faecalis* (15.8%) and *Klebsiella pneumoniae* (5.3%).


The studied risk factors (associated infectious pathologies, use of broad-spectrum antibiotics, renal and urinary tract malformations, mechanical ventilation, parenteral nutrition, and intra-vascular catheter) were more frequent in Group II (p<0.05). These results are described in Table 3 and Fig. 3.

Nine newborn infants (21.5%) presented infectious pathologies: omphalitis (33.3%), pyodermitis (33.3%), bronchopneumonia (22.2%), and sepsis (11.1%) in the Group I. In Group II this value was increased to 63.2% (meningitis - 8.3%, omphalitis - 25%, bronchopneumonia - 25%, and sepsis - 41.6%).

The urological malformations were as follows: hydronephrosis (27.3% in Group I and 9.1% in Group II), ureteropelvic junction obstruction (27.3%

Table 3 - Risk factors for urinary tract infection.

Risk factor	Group I n=42 n (%)	Group II n=19 n (%)	P	Test
Associated infectious diseases	9 (21.4)	12 (63.2)	0.001	chi-square
Use of broad-spectrum antibiotics	3 (7.1)	6 (31.6)	0.02	chi-square
Renal and urinary tract malformations	4 (9.5)	7 (36.8)	0.01	chi-square
Mechanical ventilation	1 (2.4)	4 (21.1)	0.04	chi-square
Parenteral nutrition	1 (2.4)	10 (52.6)	0.0006	chi-square
Intravascular catheter	1 (2.4)	5 (26.3)	0.01	chi-square

**Figure 3** - Risk factors for urinary tract infection.

in Group II), posterior uretero valves (27.3% in Group II, and polycystic kidney (9.1% in Group II).

Correlation between risk factors and urinary tract infection, in both studied groups, using relative risk analysis is shown in Table 4.

Table 5 shows the same analysis, drawing from more than one risk factor.

DISCUSSION

The first description of urinary tract

infection in the neonatal period dates from the beginning of the century, and the observations made are still valid. Due to the absence of specific symptoms, the confirmation of diagnosis is based on laboratory data⁴. The clinical presentation may vary from fever and other signs of sepsis to minimal changes, or the infant may be without signs⁴.

Until the decade of 1960, studies of urinary tract infection in the neonatal period were discrepant because they presented different criteria to define

this infection and used divergent methodologies¹⁷.

Currently, research indicates that urinary tract infection incidence is correlated to the presence of risk factors. The environment of the studied population, i.e., the intensive care unit or the nursery, should be taken into consideration.

The study population had a urinary tract infection frequency of 1.6%, a rate close to the low incidence limit of this disease in the neonatal period¹². This finding may be due to the low risk in this study population, because in the absence of risk factors the full-term infant rarely develops urinary tract infection.

The population of this study (61 full-term infants) presented nonspecific symptoms (Table 2 – Fig. 2) and positive urine culture collected by bag specimen, which was considered to be the presumptive urinary tract infection diagnosis. Diagnosis (positive urine culture by suprapubic aspiration) was confirmed in 19 full-term newborn infants, emphasizing the need for this method to make accurate etiologic diagnosis.

Another consideration concerning the incidence of the urinary tract infection is its relationship to associated infectious pathologies; when newborn infants develop sepsis, for instance, there is a higher chance of developing urinary tract infection^{5,6}. The results showed that in Group II, presence of infection was one of the most prevalent associated risk factors (63.2%), with significant statistical difference as compared to Group I (Table 3 – Fig. 3). Therefore, urinary tract infection should be considered in all infants who have signs of sepsis.

Prematurity and male gender are classically called risk factors⁸. There is also a proportional relationship with the presence of renal and urinary tract malformations¹⁷. Furthermore, a prolonged stay in the nursery may be con-

sidered as a risk factor, for instance, prematurity, perinatal asphyxia, mechanical ventilation, and inadequate nutrition. The incidence of neonatal urinary tract infection as a complication of intensive care has risen sharply in recent years.

Risk factors included in this research (associated infectious pathologies, use of broad-spectrum antibiotics, renal and urinary tract malformations,

mechanical ventilation, and intravascular catheter) were more frequent ($p < 0.05$) in Group II (Table 3 – Fig. 3).

Congenital renal and urinary tract defects, mainly the obstructive anomalies^{17,19}, exhibit a close relationship with urinary tract infection, and these were more frequent ($p < 0.05$) in Group II (Table 3 – Fig. 3).

The male presents a high incidence of renal and urinary tract malforma-

tions and sepsis, leading us to believe that these malformations increase the frequency of urinary tract infections. Circumcision seems to reduce the overall incidence of urinary tract infections, although a few studies have suggested that this procedure may be a predisposing factor for urinary tract infection, mainly due to the complications encountered¹⁰. In this study group, the absence of circumcision led us to exclude this procedure as a risk factor or protector factor of neonatal urinary tract infection. In our study, Group II presented no difference related to gender. The low sample size is one of the factors that could explain this finding.

Natural defenses in the urinary tract include antibacterial properties of urine and urinary tract mucosa, anti-adherence mechanisms, mechanical effects of urinary flow, presence of phagocytic cells, and immune mechanisms²². In the neonatal period, these mechanisms may fail, facilitating an urinary tract infection³. Furthermore, in the neonate, it is frequently difficult to know whether an urinary tract infection is the cause or the result of bacteremia¹⁴.

Another factor to be considered is the use of broad-spectrum antibiotics, since use of these antibiotics is necessary for treatment of undetermined infection. Use of broad spectrum antibiotics may change neonate natural flora, increasing infections by opportunistic microorganisms¹⁸. Fungal urinary tract infections, mainly *Candida albicans* well exemplify this fact¹⁵.

Table 3 and Fig. 3 reinforce that the use of broad-spectrum antibiotics (cefotaxime and vancomycin) can be considered as a factor risk, through analyzed data-frequency 31.6% ($p < 0.05$).

All invasive procedures, even with antiseptics, could lead to a bacterial contamination, especially in the neonatal period. Intravascular catheterization, often necessary to deliver prescription drugs or parenteral nutrition, can facili-

Table 4 - Relative risk (RR) analysis of the risk factors for urinary tract infection.

Risk factors	RR	CI:95%	P	Test
Associated infectious pathologies	3.27	1.51-7.04	0.001	chi-square
Use of broad-spectrum antibiotics	3.03	1.51-6.08	0.012	Fisher exact
Renal and urinary tract malformations	2.97	1.57-5.64	0.007	Fisher exact
Mechanical ventilation	2.99	1.61-5.53	0.029	Fisher exact
Parenteral nutrition	5.05	2.72-9.39	0.0009	Fisher exact
Intravascular catheter	3.27	1.84-5.83	0.009	Fisher exact

Table 5 - Relative risk (RR) analysis using more than one risk factor for urinary tract infection.

Risk factor association	PR	CI:95%	P	Test
Associated infectious diseases + Use of broad-spectrum antibiotics	2.95	1.50-2.78	0.003	chi-square Yates' corrected
Parenteral nutrition + Intravascular catheter	11.05	2.68-45.64	0.00004	Fisher exact
Parenteral nutrition + Intravascular catheter + Mechanical ventilation	8.11	2.55-25.75	0.00004	Fisher exact
Associated infectious diseases + Renal and urinary tract malformations	3.54	1.99-6.29	0.00003	chi-square Yates' corrected
Parenteral nutrition + Intravascular catheter + Renal and urinary tract malformations	3.98	1.54-10.28	0.006	Fisher exact

tate a bacteremia and consequently a urinary tract infection. Also, intubation, a necessary mechanical ventilation procedure, can provoke infection.

The frequency of mechanical ventilation, intravascular catheterization, and parenteral nutrition (Table 3 – Fig. 3) was higher in Group II.

Finally, all analyzed risk factors in this research (associated infectious pathologies, use of broad-spectrum antibiotics, renal and urinary tract malformations, mechanical ventilation, parenteral nutrition, and intravascular catheter) were present at high frequencies in Group II (Table 3 – Fig. 3), with statistical significance ($p < 0.05$). These

findings support the relationship between risk factors and urinary tract infection in the neonatal period.

The relative risk of urinary tract infection risk factors was, in decreasing importance: parenteral nutrition, intravascular catheter, associated infectious diseases, use of broad-spectrum antibiotics, mechanical ventilation, and renal and urinary tract malformations (Table 4).

Analysis using more than one urinary tract infection risk factor (Table 5) revealed an increased relative risk, rising from 2.95 to 11.05. The combination of intravascular catheter and

parenteral nutrition had the highest relationship with neonatal urinary tract infection frequency.

In conclusion, the results of this study showed that the presence of predisposing factors or risk factors (parenteral nutrition, intravascular catheter, associated infectious pathologies, use of broad-spectrum antibiotics, mechanical ventilation, and renal and urinary tract malformations) can contribute to increased urinary tract infection frequency in the neonatal period, and in the presence of more than one risk factor, the urinary tract infection occurrence rose up to 11 times.

RESUMO

RHCFAP/2997

FALCÃO MC e col. - Infecção urinária em recém-nascido de termo: análise de fatores de risco. **Rev. Hosp. Clín. Fac. Med. S. Paulo** 55 (1):9-16, 2000.

Objetivos: Analisar a contribuição dos fatores de risco para a ocorrência de infecção urinária em recém-nascidos de termo.

Casuística e metodologia: Estudo retrospectivo (1997), incluindo recém-nascidos de termo com urocultura positiva por saco coletor. A indicação desta coleta foi baseada em: hipertermia ($T > 37,8^{\circ}\text{C}$), perda de peso $> 10\%$ do peso de nascimento,

alterações do estado geral (recusa alimentar, ganho insuficiente de peso e hipoatividade) ou presença de malformações nefro-urológicas. Nesses recém-nascidos foi realizada punção suprapúbica para confirmação diagnóstica. Os recém-nascidos foram divididos em dois grupos, segundo o resultado das uroculturas: Grupo I (diagnóstico presuntivo de infecção urinária) e Grupo II (diagnóstico confirmado de infecção urinária), para avaliação dos fatores de risco pela análise do risco relativo.

Resultados: Foram estudadas 61 crianças (5,1% dos recém-nascidos de termo) - Grupo I $n=42$ (68,9%) e

Grupo II $n=19$ (31,1%). Os fatores de risco avaliados (patologias infecciosas associadas, uso prévio de antibióticos, malformações nefro-urológicas, ventilação mecânica, nutrição parenteral e o uso de cateteres) foram mais frequentes no Grupo II ($p < 0,05$). Analisando-se esses fatores, através de risco relativo, encontrou-se, em ordem decrescente de importância: nutrição parenteral, uso de cateteres, patologias infecciosas associadas, antibiótico prévio, ventilação mecânica e a presença de malformações nefro-urológicas.

Conclusões: Os resultados mostraram que a presença de nutrição

parenteral, uso de catéteres e patologias infecciosas associadas contribuíram para aumentar a frequência da infecção do

trato urinário no período analisado e, na presença de mais de um desses fatores, o risco elevou-se em até 11 vezes.

DESCRITORES: Infecção urinária. Fatores de risco. Recém-nascido.

REFERENCES

- BACHUR R & CAPUTO GL - Bacteremia and meningitis among infants with urinary tract infections. **Pediatr Emerg Care** 1995; **11**:280-284.
- BARKEMEYER BM - Suprapubic aspiration in very low birth weight infants. **Pediatrics** 1993; **89**: 457-458.
- BENADOR D, BENADOR N, SLOSMAN D et al. - Are younger children at highest risk of renal sequelae after pyelonephritis. **Lancet** 1997; **9**:349-317.
- BRION CP, SATLIN LM & EDELMANN CM - Renal disease. In: AVERY GB, FLETCHER MA & MACDONALD MG eds. **Neonatology: pathophysiology and management of the newborn**. 4th ed. Philadelphia, Lippincott, 199. p. 792-886.
- CHIU C, LIN T & BULLARD MJ - Identification of febrile neonates unlikely to have bacterial infections. **Pediatr Infect Dis J** 1997; **16**:59-63.
- CRAIN EF & GERSHEL JC - Urinary tract infection in febrile infants younger than 8 weeks of age. **Pediatrics** 1990; **86**:363-367.
- HANSSON S, BRANDSTROM P, JODAL U et al. - Low bacterial counts in infants with urinary tract infection. **J Pediatr** 1998; **2**:132:180.
- HELLERSTEIN S - Urinary tract infections: old and new aspects. **Pediatr Clin North Am** 1995; **42**:1433-1457.
- HENRY JB, LANZON RB & SCHUMANN GB - Basic examination of urine. In: HENRY JB ed. **Clinical Diagnosis and management by laboratory methods**. 19th. Philadelphia, Saunders, 1996. p. 411-56.
- HERZOG LW - Urinary tract infection and circumcision. A case-control study. **Am J Dis Child** 1989; **143**:348-350.
- HOBERMAN A, WALD ER, PENCHANSKY L et al. - Enhanced urinalysis as a screening test for urinary tract infection. **Pediatrics** 1993; **89**: 1196-1198.
- HOBERMAN A & WALD ER - Urinary infections in young children. **Pediatr Infect Dis J** 1997; **16**:11-17.
- HOBERMAN A - Is urine culture necessary to rule out urinary tract infection in young febrile children? **Pediatr Infect Dis J** 1996; **15**:304-309.
- JAMES-ELLISON MY, ROBERTS R, VERRIER-JONES K et al. - Mucosal immunity in the urinary tract: changes in sIgA, FSC and total IgA with age and in urinary tract infection. **Clin Nephrol** 1997; **48**:69-78.
- KARLOWICZ G & PHILLIPS JR - Prevalence of Candida species in hospital-acquired urinary tract infections in a neonatal intensive care unit. **Pediatr Infect Dis J** 1997; **16**:190-194.
- KIERMAN SC, PINCKERT TL & KESLER M - Ultrasound guidance of suprapubic bladder aspiration in neonates. **J Pediatr** 1993; **123**:789-791.
- KLEIN JO & LONG SS - Bacterial infections of the urinary tract. In: REMINGTON JS & KLEIN JO eds. - **Infectious diseases of the fetus and newborn**. 4th. Philadelphia, Saunders, 1995. p. 925-34.
- LOHR JA, DOWNS SM, DUDLEY S et al. - Hospital acquired urinary tract infections in pediatric patient: a prospective study. **Pediatr Infect Dis J** 1994; **13**:8-12.
- MAN P, CLAESON I, JOHANSON IM et al. - Bacterial attachment as a predictor of renal abnormalities in boys with urinary tract infection. **J Pediatr** 1989; **115**:915-922.
- MITCHELL CK, FRANCO SM & VOGEL RL - Incidence of urinary tract infection in a inner-city outpatient. **J Perinatol** 1995; **15**:131-134.
- MORELL RE, DURITZ G & OLTORF C - Suprapubic aspiration associated with hematoma. **Pediatrics** 1982; **69**:455.
- PISACANE A, GRAZIANI L, MAZZARELLA G et al. - Breast-feeding and urinary tract infection. **J Pediatr** 1992; **120**:87-89.
- POLLACK CV, POLLACK ES & ANDREW ME - Suprapubic bladder aspiration versus urethral catheterization in ill infants: success, efficiency, and complication rates. **Am Emerg Med** 1994; **23**:225-230.

24. VOORA S, SNIRIVASAN G, LILIE LD et al. - Fever in full-term newborns in the first four days of live. **Pediatrics** 1982; **69**:40-44.
25. WOODA GL - Specimen collection and handling for diagnosis of infectious diseases. In: HENRY JB ed. - **Clinical Diagnosis and management by laboratory methods**. 19th. Philadelphia, Saunders, 1999. p. 1311-1334.
26. WOODS GL, AYERS LW, WASHINGTON JA et al. - Medical bacteriology. In: HENRY JB ed. - **Clinical Diagnosis and management by laboratory methods**. 19th. Philadelphia, Saunders, 1996. p. 1132-1169.

Received for publication on the: 09/03/00