

DIFFERENT SIZES OF NET CAGES AND THE WEIGHT GAINS OF BULLFROG TADPOLES

JOSE MANDELLI JUNIOR
Pesquisador Científico
Instituto de Pesca

HENRIQUE ARRUDA SOARES
Pesquisador Científico
Instituto de Pesca

DORIVAL FONTANELLO
Pesquisador Científico
Instituto de Pesca

LUIZ ANTONIO PENTEADO
Pesquisador Científico
Instituto de Pesca

BENEDICTO DO ESPIRITO SANTO DE CAMPOS
Pesquisador Científico
Instituto de Zootecnia

MANDELLI JUNIOR, J.; SOARES, H.A.; FONTANELLO, D.; PENTEADO, L.A.; CAMPOS, B.E.S. Different sizes of net cages and the weight gains of Bullfrog tadpoles. *Rev.Fac.Med.Vet.Zootec. Univ. S.Paulo*, 24(2):209-212, 1987.

SUMMARY: The effect of different sizes of net cages upon the weight gains of Bullfrog tadpoles was studied in an outdoor facility, with a constant stock density of 0.5 individual per liter of water. Small baskets were found to be more effective for the Bullfrog tadpoles cultivation.

UNITERMS: Bullfrog; Weight of tadpoles; Housing, net-cages

INTRODUCTION

Much attention has always been dedicated by researchers in the field of livestock production to estimate both the proper size of the enclosure and the number of animals which should be kept in it.

As far as the bullfrog larvae are concerned, CULLEY et alii, 1 (1977) proposed a rearing system for bullfrog tadpoles in laboratory to diminish the problems arisen with the cultivation of large numbers of individuals within a small space. This effort aimed to contribute substantially to the standartization of the management techniques employed in many amphibian larvae and even in fish and aquatic invertebrate cultivations.

Basically those authors verified no differences in growth, consume and feed conversion when 8 to 13 tadpoles per liter densities were compared, employing rectangular baskets with capacity ranging from 25 to 30 liter of water. Yet some doubt remained whether increase in density could lead to lesser weight gain. SOARES, et alii, 6 (1983) verified in field conditions that the weight gain falls as density increases.

If something is already known as far as the density is concerned, it seems to be altogether the contrary in regard to the effect of the space availability upon the weight gain of Bullfrog tadpoles when the density is kept constant.

The refered bibliography scarcely informs on the volume of the tanks to be employed in the tadpole culture of this species both in laboratory and in field conditions. It is an important question to answer, as stated MARSCHALL, 4 (1978): "the question of the effect of the space disponibility in the same crowd conditions is still waiting for an answer".

The present work intends to contribute to the knowledge of the effect of the size of the tank upon the weight gain of Bullfrog tadpole aiming the development of the commercial frogculture in Brazil.

MATERIAL AND METHOD

The experiment was carried out at the Experimental Frogculture Facility of the Instituto de Pesca of the Secretaria

da Agricultura e Abastecimento do Estado de São Paulo, located in Parque Fernando Costa, São Paulo City, São Paulo State, Brazil.

The experiment ran from April, 4 th. through October, 1 st., 1985 and 90 Bullfrog tadpoles, belonging to the same spawn, at the stage 25 (GOSNER, 3, 1960) were utilized.

In a tank for fish culture, 20 rectangular parallelepiped screen-walled baskets were installed, being five of them 0.850 x 0.213 x 0.350 m (63.36 liters) (treatment A); five, 0.850 x 0.425 x 0.350 m (125.44 liters) (treatment B); five, 0.850 x 0.203 x 0.350m (84.19 liters) (treatment C); five, 0.850 x 0.850 x 0.350m (252.87 liters) (treatment D). All baskets were constructed with 6 meshes per centimeter nylon net. The water depth in all baskets was 0.200m. Also in all baskets the stocking rate was 0.5 tadpoles per liter.

The spring water flow was 50 liter per minute.

A commercial ration containing 36% of crude protein was offered ad libitum.

During experimental period, temperature was daily recorded at 10:00 a.m.

To estimate the weight gain during the period, the animals in each basket were weighed at the beginning and the end of the experiment with a scale of 0.1g sensibility.

The statistical design was the randomized blocks with four treatments

(A, B, C and D) and five replicates. Each of the four treatments was represented by a basket dimension as above mentioned. The smallest experimental units, i.e. the baskets, were randomly distributed within the blocks. Weight gains were submitted to analysis of variance, followed by F and Tukey tests at 1% probability level (SNEDECOR & COCHRAN, 5, 1980). To show schematically weight gains comparisons among treatments, the Duncan bar representation was employed (GOMES, 2, 1982).

Data were processed in a PC 1500 RP - SHARP.

In order to follow the progressive performance of tadpoles in each treatment during the experimental period; five extra weight measures were taken between the first and last ones. Thus, with the dates on the abscissa and the mean weights on the ordinate a bar diagram was constructed.

RESULTS AND DISCUSSION

Fig. 1 shows variations in the bullfrog tadpole mean weights in each treatment during experimental period.

Analysis of variance, Tukey test value and Duncan bars are represented on Tab. 1.

Tukey test minimum significant difference (M.S.D.) value was 2.0024, revealing that only contrast between A and D differed significantly.

TABLE 1 - Summary table of one-way analysis of variance on the weight-gains of Bullfrog tadpoles cultivation. Tukey test and Duncan bars representation.

SV	d.f.	S.Sqs	M.S.	F
Treatment	3	15.5518	5.1839	7.8212**
Block	4	3.7033	0.9255	1.3963
Error	12	7.9540	0.6638	
Total	19	27.2080		
M.S.D. Tukey 1% = 2.0024		C.V = 10.41%		
m A = 9.0155	m B = 8.0291	m C = 7.6957	m D = 6.5465	

(**) = P < 0.01

m A = 9.0156g

m B = 8.0291g

m C = 7.6957g

m D = 6.5465g

m = treat. mean weight gain

SV = Source of variation

d.f. = degree of freedom

S.Sqs. = Sum of Squares

M.S. = Mean Square

C.V. = Coefficient of variability

F = F test values




FIGURE 1 - Variations in mean weights in bullfrog tadpoles in each treatment during the experimental period.

Average water temperature during experimental period was $18.77\text{ }^{\circ}\text{C} \pm 2.09\text{ }^{\circ}\text{C}$.

From the results (Tab.1 and Fig.1) very trustful inferences can be drawn on the effect of the basket volume upon Bullfrog tadpole weight gains.

Statistical analysis of data showed that, as the volume of the basket grew bigger, the weight gain decreased. This phenomenon should not be attributable to possible differences in the quantities of rations which were offered to the animals along the experimental period, since all care was taken to offer excessive feed to the animals. Fig. 1 shows that in all weight measures from the second on the profile gotten kept constant and always favouring the smaller baskets.

At least for the time being, it would be advisable to use small, loosely crowded tanks in intensive cultures. Yet

much remain to be done until conclusive answers can be reached in this field.

CONCLUSION

The small parallelepiped baskets seem to enhance better weight gains than the larger ones in Bullfrog tadpole intensive culture.

ACKNOWLEDGMENTS

The authors are indebted to Albino Joaquim Rodrigues and José Marques dos Reis, scientific researchers, and to Mrs. Rodrigues as well, for their precious suggestions.

MANDELLI JUNIOR, J.; SOARES, H.A.; FONTANELLO, D.; PENTEADO, L.A.; CAMPOS, B.E.S. Diferentes tamanhos de tanque-rede e ganho de peso em girinos de rã-touro. (Rana Catesbeiana). Rev.Fac.Med.Vet. Zootec.Univ. S.Paulo, 24(2):209-212, 1987.

forma de paralelepipedo reto, sobre o ganho de peso em girinos de rã-touro, mantendo-se a densidade constante de 0,5 girinos por litro de água. Os resultados permitiram inferir que tanques pequenos são mais favoráveis à criação de girinos de rã-touro.

RESUMO: Estudou-se, a céu aberto, o efeito de diferentes tamanhos de tanque-rede, em

UNITERMOS: Rãs-touro; Peso, girinos; Instalações, tanques-rede

LITERATURE

- 1 - CULLEY JR., D.D.; MEYERS, S.P.; DOUCETTE JR., A.J. A high density rearing system for larval anurans. Lab.Anim., 6: 34-41, 1977.
- 2 - GOMES, P.P. Curso de estatística experimental. 2.ed. Piracicaba, Escola Superior de Agricultura "Luiz de Queiroz", 1982.
- 3 - GOSNER, K.L. A simplified table for staging anuran embryos and larvae with notes on identification. Herpetologica, 16:183-191, 1960.
- 4 - MARSCHALL, D.G. Development of testing procedures, feed formulation and protein requirements for Rana catesbeiana larvae. Baton Rouge, 1978. [Thesis of Master of Science, Louisiana State University].
- 5 - SNEDECOR, G.W. & COCHRAN, W.G. Metodos estadísticos. 7.ed. Mexico, Continental, 1980.
- 6 - SOARES, H.A.; FONTANELLO, D.; MANDELLI JR., J.; REIS, J.M.; PENTEADO, L.A.; CAMPOS, B.E.S. Efeito da densidade de população no ganho de peso de girinos de Rã-touro (Rana catesbeiana Shaw, 1802). Bol.Inst.Pesca, 10:47-51, 1983.

Recebido para publicação em 25/06/86
Aprovado para publicação em 25/02/87