

Effect of Gibberellic Acid (GA₃) Inflorescence Application on Content of Bioactive Compounds and Antioxidant Potential of Grape (*Vitis* L.) ‘Einset Seedless’ Berries

M. Kaplan^{1*}, A. Najda², K. Klimek³, A. Borowy¹

(1) Department of Pomology and Nurseries, University of Life Sciences, 58 Leszczyński Street, 20-068 Lublin, Poland

(2) Department of Vegetable Crops and Medicinal Plants, University of Life Sciences, 58 Leszczyński Street, 20-068 Lublin, Poland

(3) Department of Applied Mathematics and Informatics, University of Life Sciences, 28 Głęboka Street, 20-612 Lublin, Poland

Submitted for publication: June 2018

Accepted for publication: August 2018

Keywords: Seedless grape, hormonisation, DPPH, anthocyanins, correlation

Gibberellic acid (GA₃) is a plant growth regulator widely used in the cultivation of seedless grape varieties to increase their yield. Hormonisation treatment has beneficial effects on yield size and quality, yet its influence on the level of biologically active compounds and grape antioxidant activity has not yet been studied extensively yet. Clusters of 11-year-old ‘Einset Seedless’ grapevines trained according to the single Guyot pruning style were sprayed with GA₃ at 100, 200 or 300 mg/L dose once, twice or three times. Fruit harvested on 25 September were immediately examined for acidity, extract content, biologically active substances and antioxidant capacity using the DPPH test. In addition, correlations occurring between some parameters measured were calculated. Hormonisation had a negative effect on the content of extract, flavonoids and ascorbic acid, while it had no effect on the anthocyanin level. The antioxidant activity determined by the DPPH assay depended on dose and the number of treatments, and the analysed parameters were shown to decrease significantly with increasing application number. Gibberellic acid at 100 and 300 mg/L application rates had a significantly increased DPPH level compared to the control and 200 mg/L dose. The single GA₃ treatment, and when applied three times, and application rates at 100 and 200 mg/L were shown to have a significant influence on phenolic acid content. The level of tannins after a single GA₃ treatment and a 300 mg/L dose increased significantly.

INTRODUCTION

Grape consumption is strongly correlated with the reduced risk factor for developing chronic diseases, such as cardiovascular disorders and cancer (Arts & Hollman, 2005; Erdman *et al.*, 2007; Leifert & Abeywardena, 2008; Vislocky & Fernandez, 2010; Doshi *et al.*, 2015). This results from, among others, the presence of biologically active compounds like polyphenols, which display powerful antioxidant effects, along with anti-inflammatory, anti-carcinogenic and anti-platelet properties. Polyphenols also help dilate blood vessels, boost the immune system and play a neuroprotective role, attributed mainly to their ability to modulate and induce signalling pathways (Frankel, 1999; Stevenson & Hurst, 2007; Pezzuto, 2008; Dohadwala & Vita, 2009; Crozier *et al.*, 2010; Vislocky & Fernandez, 2010; Xia *et al.*, 2010; Doshi *et al.*, 2015). Polyphenols inactivate free radicals, chelate divalent metal ions, and thus lower their oxidant potential (Scalbert *et al.*, 2005).

The qualitative and quantitative composition, distribution and antioxidant activity of polyphenols in grapes are quite

variable and depend on species, cultivar, location in berry (skin, pulp, seeds, juice), climate-soil conditions (exposure to light, temperature, soil type), agrotechnical practices (irrigation, nutrient availability, application of plant growth regulators, harvest time, berry maturity, yield and berry size) and, finally, post-harvest conditions and storage-processing techniques (Kim *et al.*, 2003; Peña-Neira *et al.*, 2004; Jiang *et al.*, 2006; Montealegre *et al.*, 2006; Orak, 2007; Xia *et al.*, 2010; Liang *et al.*, 2011). Polyphenols are responsible for the major sensory attributes of products and beverages of plant origin, as they are the determinants of their appearance (colour) and taste, *i.e.* flavour, bitterness and astringency (Tomás-Barberán & Espin, 2001; Es-Safi *et al.*, 2007).

Seedless grape varieties have been on the rise in the world grape market because of their high quality and consumer preferences, and they enjoy increasing popularity not only as table grapes, but as raisins as well (Artés-Hernández *et al.*, 2006). One of the most promising commercial seedless cultivars that can be grown successively in cool-climate

*Corresponding author: E-mail address: magdalena.kaplan@up.lublin.pl [Tel.: 48 81 5247158]

Acknowledgments:

areas, such as Poland, is 'Einset Seedless', a pink grape with a unique strawberry-like flavour. This variety can be used for raisin production or for fresh consumption as table grapes. However, the natural berry size of the 'Einset Seedless' variety (± 2 g) is not large enough for table grape use and thus represents a problem for commercialisation. To overcome this problem, and to improve grape size and quality, plant growth regulators (most often gibberellic acid, GA₃) are applied globally (Harrell & Williams, 1987; Dimovska *et al.*, 2014; Nampila *et al.*, 2010; Kapłan *et al.*, 2017). Gibberellic acid promotes cell division, enhances earlier blooming and increases fruit size and yield. The effect of GA₃ application relies on the variety, dose and application time (Khan *et al.*, 2009; Nampila *et al.*, 2010; Dimovska *et al.*, 2014; Kapłan *et al.*, 2017). The earlier studies by the present authors showed a positive response of this cultivar to gibberellic acid treatment considering fruit set, and the size of clusters and berries (Kapłan, 2011; Kapłan *et al.*, 2017). These findings are of the utmost importance currently, because modern table grape production is expected to fully conform with the requirements of a market that is demanding improved grape quality, that is aiming for uniformly repeatable clusters, equal berry size, shape and uniform skin colour, as well as for increased resistance to transportation. An important attribute of grape berry quality has been proven to be the absence of seeds (Dimovska *et al.*, 2014).

The objective of the present study was to determine the parameters affecting the content of biologically active compounds and antioxidant activity in the 'Einset Seedless' grape variety subject to the dose and number of gibberellic acid applications.

The influence of gibberellic acid on the content of biologically active compounds and the antioxidant potential of grapes with regard to concentration and the number of treatments on grapevines is an innovative idea. The relevant available literature does not include any experiments exploring the effect of hormonisation on the antioxidant activity level in grapes.

MATERIALS AND METHODS

Plant materials

The field experiment assessed the effect of GA₃ dose and the number of treatments on the level of chosen secondary metabolites of grapes. The analysed fruit were obtained from the NOBILIS Vineyard (50°39'N; 21°34'E) located in the Sandomierska Upland in south-eastern Poland. The own-rooted vines of 'Einset Seedless' were planted in spring 2003 at a spacing of 2.0 x 1.0 m (5 000 units/ha) on lessive soil developed from loess deposits. Throughout the experiment, regular soil-protective measures against diseases, pests and weeds were carried out in compliance with the current grapevine protection programme. The grapevines were not watered. An average number of clusters per vine was 17 to 18, while grape yield averaged 3.5 kg per vine. Grapes were harvested on 2014-09-25. The grapevines were pruned in the single Guyot pruning style. The inflorescences of 11-year-old 'Einset Seedless' grapevines were treated with GA₃ spray at three dose levels, *viz.* 100, 200 or 300 mg/L, once (seven days after full bloom, when 70% of berries in the cluster were 1 mm in diameter), twice (seven and 14 days after full

bloom, when 70% berries in the cluster were 1 and 3 mm in diameter respectively) or three times (seven, 14 and 21 days after full bloom, when 70% of berries in the cluster were 1, 3 and 6 mm in diameter respectively). The solution contained 99% gibberellic acid and an adhesive and wetting SILWET Gold agent at 0.015% dose, *i.e.* 150 μ L. The solution was prepared immediately before the treatment. The clusters were treated with a hand pump sprayer, covering the pedicels and berries thoroughly. On average, 50 mL of solution was enough to thoroughly cover all the grape clusters on a vine. The untreated grapes constituted the control.

The field experiment was set up in a randomised block design, including 10 combinations with five replications comprising plots of three grapevines each. The harvested fruit underwent laboratory tests at the Laboratory for Vegetable and Herbal Material Quality in the Department of Vegetable Crops and Medicinal Plants of the University of Life Sciences in Lublin. The research material comprised the 'Einset Seedless' grape variety ('Fredonia' \times 'Canner', Reisch *et al.*, 1986) subjected to hormonisation with gibberellic acid (GA₃).

Chemicals

All reagents and solvents were analytical grade chemicals from Merck (Darmstadt, Germany), Sigma Chemical Co. (St. Louis, MO, USA) or POCh (Gliwice, Poland). GA₃ was obtained from Acros Organics™ (Thermo Fisher Scientific, Geel, Belgium) and SILWET Gold from Chemtura Europe Limited (Warsaw, Poland).

Physicochemical analyses

Fruit extract content was measured on harvest day using a refractometer, Abbe WAY 2W (EnviSense, Poland) while squeezing the juice from 100 representative berries collected from each combination. In order to determine biologically active compounds and antioxidant activity, grapes were transported to the laboratory on harvest day, stored in a cooler at 8°C for 16 hr, and finally underwent the chemical analyses. Titratable acidity (TA) was determined in accordance with Polish Norm PN-90/A/75101/02.

Determination of L-ascorbic acid

The fresh and comminuted grape fruits (5 g) were extracted twice for 30 min with 2.5 ml 4.0% (m/V) L-cysteine and 10.0 ml water by sonification. All aqueous extracts were combined and diluted with water to 25 mL. The samples were analysed using high performance liquid chromatography. Analyses were done with a LaChrom-Merck HPLC system with a photodiode array detector (DAD L-7450), and all separations were done in a Lichrospher 100 RP18 column (250.0 \times 4.0 mm, 5.0 μ m; Merck). The mobile phase consisted of 0.0272 g/L KH₂PO₄ adjusted to pH 2.40 with H₃PO₄, applied in isocratic elution for 30 min. The flow rate was adjusted to 1.0 mL/min. The detection wavelength was set to DAD at $\lambda = 254.0$ nm. Samples of 20.0 μ L were injected. All separations were performed at 24.0°C. The peaks were assigned by spiking the samples with standard compounds and comparing the UV spectra and retention times (ascorbic acid, 5.66 min) (Najda, 2017). Calibration curves were obtained from five doses of each external

standard (0.01 to 1.40 mg/mL). The regression coefficient (R_2) of the calibration curve for ascorbic acid was equal to $Y = 85.231, X = 18.787$. The RSD value for the repeatability ($n = 4$) of standard solution was 0.40% (0.01 mg/mL ascorbic acid). The limits of quantitation (LOQ) and detection (LOD) of ascorbic acid were 0.16 and 0.04 mg/L respectively. All solvents used were HPLC grade (Merck). Reference standards were obtained from Sigma-Aldrich.

Total phenolic acid estimation was carried out according to the Arnov method (Polish Pharmacopoeia, 2002). One millilitre of sample was mixed with 5 ml of distilled water, 1 mL 0.5 M HCl, 1 mL of Arnov reagent and 1 mL 1M NaOH, and subsequently adjusted to 10 ml with distilled water. The absorbance was measured at 490 nm. The total phenolic acid content was expressed as caffeic acid equivalent (CAE).

Estimation of anthocyanins by means of colorimetry

Samples of raw material (1.0 g) were extracted with 50 ml HCl (1 mol/dm) and heated in a water bath for 1 hr. The obtained extract was hydrolysed with 20 ml n-butanol, and then two 10 ml n-butanol portions were added as a solution. Anthocyanin extracts were rinsed with n-butanol in a 50 mL flask. The absorbance was measured immediately at 533 nm (Miłkowska & Strzelecka, 1995).

The percentage of anthocyanins, expressed as delphinidin chloride, was calculated from the expression:

$$P = \frac{A \times V \times F}{M}$$

where: P = total anthocyanins (mg/100 g), A = absorbance at 533 nm, V = value of butanol phase (50 mL), F = coefficient expressed as delphinidin chloride (2.6), and m = mass of sample to be examined (mg).

Determination of antiradical activity (AA)

A 0.1 mL aliquot of the methanol extract prepared above was mixed with 3.9 mL of an 80% ethanolic 0.6 mM DPPH solution. The tubes were vortexed for 15 s and allowed to stand for 180 min, as described by Cai *et al.* (2003). After this, the absorbance of the mixture was measured at $\lambda = 517$ nm wavelength using the HITACHI UV-Vis spectrophotometer (UV-Vis model U-2900. Shimadzu, Kyoto, Japan). Most tested compounds reacted completely within 180 min under these conditions. The reaction time for vitamin C was less than 1 min due to its fast oxidation. Ethanol (80%) was used as a blank solution, and DPPH solution without test samples (3.9 ml of DPPH + 0.1 ml of 80% ethanol) served as the control. All tests were performed in triplicate. The antiradical activity of the test samples was expressed as the median effective dose for radical scavenging activity (EC_{50}): TP (mg) of antioxidant (test sample) required for a 50% decrease in absorbance of DPPH radicals and inhibition (%) of DPPH absorbance = $(A_{\text{control}} - A_{\text{test}}) \times 100 / A_{\text{control}}$. A plot of absorbance of DPPH vs dose of antioxidant was made to establish the standard curves (dose-response curves) and to calculate that $EC_{50} \cdot A_{\text{control}}$ is the absorbance of the control (DPPH solution without the test sample), and A_{test} is the absorbance of the test sample (DPPH solution plus 0.1 mL of 5 μ M test compound). Ascorbic acid served as a standard.

The results of the assay were expressed relative to an ascorbic acid equivalent.

Estimation of total flavonoids

The studied material was investigated for total content of flavonoids using a modified Christ and Müller method, calculated for quercetin, QE (Polish Pharmacopoeia, 2014). Absorbance was measured at 425 nm on a HITACHI U-2900 spectrophotometer.

The content of flavonoids was calculated from the equation:

$$X = \frac{8.75 \times A}{m}$$

where m (g) was the amount of fresh mass

Tannin estimation

The tannins were determined by the Folin-Ciocalteu method. About 0.1 mL of the sample extract was added to a volumetric flask (10 mL) containing 7.5 mL of distilled water, 0.5 mL of Folin-Ciocalteu phenol reagent and 1 mL of 35% Na_2CO_3 solution, and diluted to 10 mL with distilled water. The mixture was shaken well and kept at room temperature for 30 min. A set of reference standard solutions of gallic acid (20, 40, 60, 80 and 100 μ g/mL) was prepared in the same manner as described earlier. The absorbance for the test and standard solutions was measured against the blank at 725 nm with a UV/Visible spectrophotometer. The tannin content was expressed in terms of % of GAE of extract (Polish Pharmacopoeia, 2011) according to the European Pharmacopoeia (2008).

Statistical analysis

The results obtained in this study were analysed statistically using one-way analysis of variance and Tukey's confidence intervals. The inference was based on a significance level at $p < 0.05$. The estimation of correlations occurring between the qualitative parameters of the grapes was done by counting Pearson's correlation coefficients. Multidimensional analysis techniques were employed to show the similarities in the groups in such a way that the homogeneous objects could be found in the same cluster. The results of the cluster analysis are graphically shown in a dendrogram. All the statistical analyses were done using SAS Enterprise Guide 5.1. software.

RESULTS AND DISCUSSION

Table 1 presents the average monthly air temperatures and monthly precipitation totals in the 2014 growing season as compared to the multi-year average of 1998 to 2008. It can be observed that the weather conditions in the study year favoured grapevine cultivation. The average air temperature during the vegetation period was higher than the average multi-year one. Notably, an average air temperature lower than the average multi-year one was recorded only in August. The average precipitation total in the 2014 growing season was found to be higher compared to the average multi-year total. The rainfall sum in May, July and August was higher than that determined in the multi-year observations.

Sugars are one of the major components that determine

fruit quality and are responsible for their sweetness. The sugar/organic acid ratio in fruit plays the most important role in the final flavour of grapes (Topalovic & Mikulic-Petkovsek, 2010). The statistical analysis indicated a significant influence of GA₃ dose and application number on total extract content, vitamin C level and total acidity of the 'Einset Seedless' grape variety (Table 2). It was found that, irrespective of GA₃ dose and number of treatments, the content of extract and vitamin C in the hormonised fruit was significantly lower compared to the control, except for the grapevines subjected to this treatment once. The significantly lowest level of both estimated parameters was recorded in the

fruit treated with GA₃ three times and with 200 mg/L GA₃. The study of Al-Atrushy (2016) showed that an increase in the number of applications and dose of gibberellic acid significantly increased the total sugar level. Dimovska *et al.* (2014) applied GA₃ twice and three times at a dose of 5, 10 and 20 mg/L and did not observe any significant effect on extract content in 'Flame Seedless' grapes.

The vitamin C content in 'Umran' grapes after 10 mg/L GA₃ application was lower than in the control, whereas at higher doses, *viz.* 30 and 50 mg/L GA₃, the hormonisation had the most beneficial impact on the parameter under study (Rachna & Singh, 2013). The vitamin C content after two

TABLE 1

Average monthly air temperatures and monthly precipitation totals in the 2014 growing season.

Month	Mean air temperature (°C)		Amount of precipitation (mm)	
	2014	Mean 1988-2008	2014	Mean 1988-2008
April	10.77	8.8	42.6	45.7
May	14.51	14.2	112.2	57.0
June	17.2	16.9	54.2	68.7
July	20.9	19.1	97.0	82.4
August	18.26	18.4	96.8	58.7
September	14.81	13.4	32.4	57.0
October	9.16	8.6	36.6	37.9
Mean (°C)	15.09	14.2	-	-
Amount of precipitation (mm)	-	-	471.80	407.4

* according to the Sandomierz weather station

TABLE 2

Effect of hormonisation on extract content, vitamin C level and total acidity in 'Einset Seedless' grapevines.

Combination	Solid content (°Bx)	Ascorbic acid (mg/100 g)	Total acidity (%)	
Dose (mg/L GA ₃)	Control	19.63 ± 0.12 D	72 ± 1 D	0.2 ± 0.1 A
	100	19.46 ± 0.38 C	47 ± 22 C	0.3 ± 0.0 B
	200	18.78 ± 1.35 A	31 ± 4 A	0.2 ± 0.1 A
	300	19.00 ± 1.02 B	41 ± 13 B	0.3 ± 0.1 B
	Mean	19.08 ± 0.92	39 ± 13	0.3 ± 0.1
	p-value	< 0.0001	< 0.0001	0.0001
Number of applications	Control	19.63 ± 0.12 C	72 ± 1 D	0.2 ± 0.1 A
	1	19.40 ± 0.35 C	41 ± 12 B	0.4 ± 0.1 B
	2	19.08 ± 1.07 B	51 ± 19 C	0.3 ± 0.1 B
	3	18.76 ± 1.33 A	27 ± 2 A	0.2 ± 0.1 A
	Mean	19.08 ± 0.92	39 ± 11	0.3 ± 0.1
	p-value	< 0.0001	< 0.0001	0.0003
Treatment × Number of applications	p-value	< 0.0001	< 0.0001	0.0021

* Mean values marked with the same letters do not differ significantly at $P < 0.05$; NS: not significant.

applications of GA₃ in seven seedless grape varieties was found to increase by 10% to 27% depending on variety, yet this was not confirmed in the present study (Gougoulis & Masheva, 2010). Awad and Al-Qurashi (2012) used gibberellic acid in the cultivation of date palms of the 'Barhee' variety and showed a positive influence of hormonisation on the vitamin C level in the fruit. The application of 100 and 150 mg/L GA₃ significantly increased the vitamin C content compared to the control and the application of 50 mg/L GA₃. It was found that the vitamin C level depended on the sugar content of the fruit of the grapevine variety under study. Moreover, the effect of gibberellic acid dose on vitamin C concentration was exactly the same as on sugar level.

Laszlo and Saayman (1990) and Topalovic and Mikulic-Petkovsek (2010) found that the acidity of grapes correlated with their taste, which results from the presence of tartaric and malic acid, whose contents reach as much as 90% in grapes. Total acidity in the fruits varied between 0.2 and 0.4%, and it differed significantly between the combinations under study. It was shown that grapes treated with 100 and 300 mg/L GA₃ and with gibberellic acid solution applied once and twice displayed significantly higher total acidity than the control. Wholly different relationships were reported by Al-Atrushy (2016), who studied grapes subjected to hormonisation (irrespective of dose and number of treatments) and found significantly lower acidity than in the control. Similarly, Dimovska *et al.* (2014) applied GA₃ at a dose of 20 mg/L and found that, irrespective of number of treatments, the application significantly decreased the acidity level. Rachna and Singh (2013) assessed the influence of gibberellic acid on the content of chosen biologically active compounds in fruits of *Zizyphus mauritiana* Lamk. cv. 'Umran' and observed an unfavourable effect of hormonisation at 50 mg/L GA₃ application rate on total acidity at harvest. On the other hand, the study of Kok (2017) did not show any significant influence of GA₃ applied in combination with a biostimulant on the acidity of 'Cardinal' grapes.

The antioxidant capacity of grape material is associated with the presence of secondary metabolites, *i.e.* phenolic acids, anthocyanins, flavonoids and tannins. The level of phenolic acids in the fruits studied depended significantly on the GA₃ dose and the number of applications (Table 3). It was found that GA₃ applied at 100 and 200 mg/L doses, as well as in single and three sprays, significantly increased the phenolic acid content in 'Einset Seedless' grapes.

Anthocyanins are a group of the most important phenolic components of dark grape cultivars, as they directly affect the intensity of berry skin coloration, which is the key quality attribute determining the market value and consumer acceptance of the grapes (Kok, 2017). In addition, anthocyanins, with their strong antioxidant properties, are involved in protection against fungal and bacterial infections. Notably, anthocyanin synthesis mostly occurs only in the grape berry skin (Doshi *et al.*, 2015). The hormonisation treatment did not significantly affect the anthocyanin level in the fruit of the grapevine variety analysed here (Table 3). This research finding is confirmed by Dimovska *et al.* (2014), whose study also did not show any significant effect of hormonisation on the content of these compounds in 'Flame Seedless' grape berries. Different results were reported by

Gougoulis and Masheva (2010) after GA₃ was applied twice, giving rise to a 30% increase in anthocyanin content in 'Kishmish Tjurkmenski' fruit. A vast body of studies indicates that the level of anthocyanins and tannins relies largely on cultivar, species, degree of maturity of the fruit, climate and site of fruit production (Mazza, 1995; Mattivi *et al.*, 2002; Muñoz-Espada *et al.*, 2004; Yang *et al.*, 2009).

The antioxidant potential of extracts of the fruit analysed was determined by the DPPH method and ranged from 56.272 to 83.652 µM TE/g. This depended significantly on the number of treatments and the dose (Table 3). The treatments applied had a significantly positive effect on the parameter under study in most combinations. It was found that the control grapes and those to which 100 and 300 mg/L GA₃ had been applied displayed significantly higher antioxidant activity compared to grapes treated with 200 mg/L GA₃. The number of hormonisation treatments also had a significant effect on the parameter analysed compared to the control combination, since increasing the number of applications significantly decreased the antioxidant potential. Tian *et al.* (2011) demonstrated that 100 mg/L GA₃ applied twice had an unfavourable effect on the level of antioxidant capacity measured by the DPPH assay in 'Muscat Hamburg' fruit. The authors highlighted the opposite relationship when studying the anatomical parts of the plant, that is leaf, stem and tendril. Gougoulis and Masheva (2010) noted a beneficial influence of hormonisation that increased the antioxidant potential in seedless grape cultivars by 16% to 42%.

The flavonoid content in the grapes under investigation varied between 0.083 and 0.103 mg of cyanidin 3-glucoside equivalents per 100 grapes, and differed significantly between the combinations assessed. The grapes subjected to hormonisation had significantly less flavonoids than the control (Table 3). It was observed that an increasing dose of gibberellic acid promoted a significant increase in flavonoid level. After three hormonisation treatments, the fruit displayed a significantly lower flavonoid content than the other hormonisation treatments and the control grapes. These results are consistent with those reported by Tian *et al.* (2011), who noted that GA₃ application decreased total flavonoid content substantially in grape pulp and skin. However, contrary results were obtained by Gougoulis and Masheva (2010), who assessed 'Trakijskaperla' fruit with amber yellow grape berries and violet-red 'Kishmish Tjurkmenski' fruit after GA₃ treatment applied twice. The authors observed an increase in flavonoid content by 10% and 12% respectively against the control.

Tannins occur in grape skin, seeds and pedicels. Their amount in fruit juice (must) and wine is related to grapevine viticultural practices, vine load and environmental conditions, maceration procedures and fermentation conditions (Matthew & Nuzzo, 2007). Tannins possess several vital properties that affect the colour and colour stability of the fruit, the depth of mouthfeel and astringency (Weston, 2005). The analysis showed that hormonisation with 300 mg/L GA₃ and a single application had a significant effect on tannin content. Similar relationships were reported by Awad and Al-Qurashi (2012), who treated the date palm 'Barhee' cv. with 100 and 150 mg/L of GA₃. However, the available literature does not provide any data on the impact

TABLE 3
Hormonisation and antioxidant activity of 'Einset Seedless' grapes.

Combination	Phenolic acid (mg 100/g FM)	Total anthocyanin (mg 100/g FM)	DPPH (μ M TE/g FM)	Total flavonoids (mg 100/g FM)	Tannins (%)
Control	0.051 \pm 0.000 B	9.930 \pm 0.501 A	66.386 \pm 0.081 B	0.103 \pm 0.001 D	0.0643 \pm 0.007 A
100	0.055 \pm 0.006 C	10.220 \pm 0.507 A	82.063 \pm 1.524 C	0.083 \pm 0.005 A	0.0680 \pm 0.009 A
200	0.056 \pm 0.006 C	10.434 \pm 0.439 A	56.272 \pm 9.681 A	0.090 \pm 0.004 B	0.0745 \pm 0.017 A
300	0.048 \pm 0.004 A	10.163 \pm 0.549 A	83.652 \pm 1.403 D	0.098 \pm 0.005 C	0.1514 \pm 0.182 B
Mean	0.053 \pm 0.005	10.272 \pm 0.498	73.996 \pm 4.203	0.090 \pm 0.005	0.098 \pm 0.069
p-value	< 0.0001	0.4944	< 0.0001	< 0.0001	0.0242
Control	0.051 \pm 0.000 A	9.930 \pm 0.501 A	66.386 \pm 0.081 A	0.103 \pm 0.001 C	0.0643 \pm 0.007 A
1	0.053 \pm 0.010 B	10.353 \pm 0.509 A	76.390 \pm 8.524 D	0.091 \pm 0.004 B	0.1629 \pm 0.176 B
2	0.051 \pm 0.002 A	10.384 \pm 0.480 A	75.424 \pm 11.610 C	0.092 \pm 0.008 B	0.0673 \pm 0.014 A
3	0.055 \pm 0.006 C	10.080 \pm 0.493 A	70.172 \pm 19.922 B	0.088 \pm 0.009 A	0.0637 \pm 0.013 A
Mean	0.053 \pm 0.006	10.272 \pm 0.494	73.996 \pm 13.375	0.090 \pm 0.007	0.098 \pm 0.068
p-value	< 0.0001	0.3844	< 0.0001	< 0.0001	0.0689
Treatment \times Number of applications	p-value < 0.0001	0.4370	< 0.0001	< 0.0001	0.0298

* Mean values marked with the same letters do not differ significantly at $P < 0.05$; NS: not significant.

of dose and the number of GA₃ sprays on the tannin level of grape berries. It was shown that the application of 300 mg/L GA₃ and a single GA₃ application significantly increase the level of the parameter studied, and similar relationships were found when assessing the antioxidant potential using the DPPH assay. Notably, a high tannin level was observed to be modified by a high DPPH level.

An interaction between gibberellic acid dose and number of treatments was found to significantly affect the chosen secondary metabolites in grapes of the 'Einset Seedless' variety, except for anthocyanins.

The Pearson's coefficient indicates a strong correlation between total extract content and a dose of 100 mg/L and a single application of GA₃, vitamin C level and treatment applied once (Table 4). A strong negative correlation was noted between total extract content and two applications of 200 mg/L GA₃; vitamin C and 300 mg/L GA₃ dose; and two and three times applied sprays as well as between total acidity and dose 200 mg/L GA₃. A negative correlation was observed between vitamin C content and application at 200 mg/L GA₃, and between total acidity and treatment at 300 mg/L GA₃ dose.

The Pearson's coefficient for the parameters determining the antioxidant activity of fruit showed a strong correlation between total phenolic acids and a dose of 100 mg/L GA₃ and a treatment rate of 300 mg/L GA₃, flavonoid level and number of applications, as well as between tannin content and a single GA₃ treatment (Table 5). A strong negative correlation existed between total phenolic acids and a dose of 200 mg/L GA₃ and the treatment applied three times, and

between the DPPH parameter and 200 mg/L GA₃ application, and between flavonoid content and single GA₃ treatment and applied two times. The correlations were established between the anthocyanin level and a single GA₃ application, the DPPH parameter and 300 mg/L GA₃ application rate, the flavonoid content and 300 mg/L GA₃ dose, and the tannin level and a single GA₃ treatment. A negative correlation was found between total phenolic acids and a single application, anthocyanin content and a 300 mg/L GA₃ application rate, and between tannin content and treatment with 200 and 300 mg/L GA₃ dose.

The dendrogram shows two separate clusters, in that one object is a clear outlier (Fig. 1). This is the control, which differs distinctively from the other combinations. The latter display high similarity at the group level. These are: 100 and 300 mg/L GA₃ dose (group 1) and 200 mg/L GA₃ dose (group 2). Both clusters are quite similar to each other.

The dendrogram (Fig. 2) made it possible to define the similarity in the effect of the number of gibberellic acid applications on a level of antioxidant activity in the fruit. The results serve to define two separate clusters exhibiting some similarities. It was shown that a single gibberellic acid treatment and GA₃ applied twice affect antioxidant activity very similarly, whereas the similarity with the control combination is noted at three times the treatment.

The PC sum (PC1 and PC2 components) of the total variation in traits for the GA₃ doses reached 81.1% – 2.47% for PC1 and 28.63% for PC2 (Fig. 3a). PC1 contained the level of flavonoids, vitamin C, anthocyanins and phenolic acids, whereas PC2 contained the potential of sugars,

TABLE 4
Correlation coefficient for chosen quality parameters defining fruit taste.

Combination	Solid content	Ascorbic acid	Total acidity
Dose (mg/L GA ₃)			
100	0.7952	-0.1309	0.1598
200	-0.8047	-0.4736	-0.9607
300	-0.0566	-0.9977	-0.6495
Number of applications			
1	0.9389	0.7633	0.2738
2	-0.8401	-0.7832	0.2165
3	-0.0217	-0.9221	-0.2582

TABLE 5
Correlation coefficient of chosen secondary metabolites.

Combination	Phenolic acid	Total anthocyanin	DPPH	Total flavonoids	Tannins
Dose (mg/L GA ₃)					
100	0.7043	0.0803	0.1858	-0.8238	-0.2604
200	-0.9707	-0.1361	-0.9577	-0.8816	-0.6893
300	0.9743	-0.6100	0.6494	0.5386	-0.6304
Number of applications					
1	-0.5000	0.4886	0.1306	0.7267	0.6362
2	-0.1218	-0.4064	-0.0680	0.8895	-0.1986
3	-0.7777	-0.2571	0.0873	0.9834	-0.0333




FIGURE 1

Branching-tree diagram for antioxidant activity in 'Einset Seedless' grapevines subject to gibberellic acid dose.




FIGURE 2

Branching-tree diagram for antioxidant activity of 'Einset Seedless' grapevines subject to number of applications.




FIGURE 3

PCA map visualising relationship between the content of a chosen secondary metabolite and dose (a), and number of GA₃ applications (b).

tannins, acidity and DPPH. The control displayed a high content of extract and flavonoids, along with a low level of anthocyanins and phenolic acids. The PCA analysis showed differences between the application rates, *i.e.* fruit subjected to 100 mg/L GA₃ treatment had a high vitamin C content. The application rates of 200 and 300 mg/L GA₃ contributed to high tannin levels and, what was particularly noteworthy, 300 mg also increased acidity.

The PC sum of total variation for the analysed number of GA₃ applications was 84% (60.73% and 23.7% respectively). PC1 indicates the phenolic acid level, while PC2 shows the other secondary metabolites, as well as the level of vitamin C and extract. The control fruit were characterised by a high content of flavonoids, vitamin C and extract, while the remaining components were present at low levels. The hormonisation spray, irrespective of the number of applications, had a beneficial effect on the level of phenolic acids, anthocyanins, DPPH and fruit acidity.

CONCLUSIONS

This study assessed the chosen biologically active compounds of 'Einset Seedless' grapes subjected to a gibberellic acid dose and a varied number of applications. The hormonisation treatment had an adverse effect on the content of extract, vitamin C and flavonoids in grapes. However, this treatment did not have significant influence on the anthocyanin level in the fruit of the grapevine variety under study. Antioxidant activity, as determined by the DPPH assay, depended on the dose and the number of treatments, and the analysed parameter was shown to decrease significantly with an increasing number of applications. Gibberellic acid at 100 and 300 mg/L application rates significantly increased the DPPH level compared to the control and the 200 mg/L dose. The single GA₃ treatment and the treatment applied three times, and application rates of 100 and 200 mg/L, were shown to have a significant influence on phenolic acid content. The level of tannins after a single GA₃ treatment and a 300 mg/L dose increased significantly.

LITERATURE CITED

Al-Atrushy, S.M.M., 2016. Effect of GA₃ dose and frequency on yield and quality of 'Zark' grape. *Jordan J. Agric. Sci.* 12(4), 1183-1191.

Artés-Hernández, F., Tomás-Barberán F.A. & Artés, F., 2006. Modified atmosphere packaging preserves quality of SO₂-free 'Superior seedless' table grapes. *Postharvest Biol. Technol.* 39, 146-154.

Arts, I. & Hollman, P., 2005. Polyphenols and disease risk in epidemiologic studies. *Am. J. Clin. Nutr.* 81, 317S-325S.

Awad, M.A. & Al-Qurashi, A.D., 2012. Gibberellic acid spray and bunch bagging increase bunch weight and improve fruit quality of 'Barhee' date palm cultivar under hot arid conditions. *Sci. Hort.* 138, 96-100.

Cai, Y.Z., Sun, M. & Corke, H., 2003. Antioxidant activity of betalains from plants of the amaranthaceae. *J. Agric. Food Chem.* 51(8), 2288-2294.

Crozier, A., Del Rio, D. & Clifford, M.N., 2010. Bioavailability of dietary flavonoids and phenolic compounds. *Mol. Aspects Med.* 31, 446-467.

Dimovska, V., Petropulos, V.I., Salamovska, A. & Ilieva, F., 2014. Flame Seedless grape variety (*Vitis vinifera* L.) and different concentration of gibberellic acid (GA₃). *Bulg. J. Agric. Sci.* 20, 137-142.

Dohadwala, M. & Vita, J.A., 2009. Grapes and cardiovascular disease. *J. Nutr.* 139, 1788S-1793S.

Doshi, P., Adsule, P., Banerjee, K. & Oulkar, D., 2015. Phenolic compounds, antioxidant activity and insulin tropic effect of extracts prepared from grape (*Vitis vinifera* L.) by products. *J. Food Sci. Technol.* 52, 181-190.

Erdman, J., Balentine, D., Arab, L., Beecher, G., Dwyer, J. T., Folts, J., Harnly, J., Hollman, P., Keen, C.L., Mazza, G., Messina, M., Scalbert, A., Vita, J., Williamson, G. & Burrowes J., 2007. Flavonoids and heart health. Proceedings of the ILSI North America Flavonoids Workshop. *J. Nutr.* 137, 718S-737S.

Es-Safi, N., Ghidouche, S. & Ducrot, P.H., 2007. Flavonoids: Hemisynthesis, reactivity, characterization and free radical scavenging activity. *Molecules* 12, 2228-2258.

European Pharmacopoeia, 2008 (6th ed). Council of Europe, Strasbourg.

Frankel, E.N., 1999. Natural phenolic antioxidants and their impact on health. In: Packer, L. (ed). *Antioxidant food supplements in human health*. Academic Press, London. pp. 385 – 392.

Gougoulis, N. & Masheva, L., 2010. Effect of gibberellic acid (GA₃) on polyphenols content and antioxidative activity of some table grape varieties of *Vitis vinifera* L. *Oxid. Commun.* 33(3), 652-660.

Harrell, D.C. & Williams, L.E., 1987. Net CO₂ assimilation rate of grapevine leaves in response to trunk girdling and gibberellic acid application. *Plant Physiol.* 83, 457-459.

Jiang, H., Ji, B.P., Liang, J.F., Zhou, F., Yang, Z.W. & Zhang, G.Z., 2006. Changes of contents and antioxidant activities of polyphenols during fruit development of four apple cultivars. *Eur. Food Res. Technol.* 223, 743-748.

Kaplan, M., 2011. Effect of growth regulator application technique on quality of grapevine 'Einset Seedless' variety (In Polish). *Acta Agrobot.* 64(4), 189-196.

Kaplan, M., Najda, A., Baryła, P. & Klimek, K., 2017. Effect of gibberellic acid dose and number of treatments on yield components of "Einset Seedless" grapevine cultivar. *Hort. Sci.* 44(4), 195-200.

Khan, M., Hafeez-ur-Rahman, A., Ahmed, M., Abbas, G. & Ahmed, N., 2009. Effect of gibberellic acid on growth and fruit yield of grape cultivar 'flame seedless'. *Int. J. Biol. Biotech.* 6(4), 265-268.

Kim, D.O., Jeong, S.W. & Lee, C.Y., 2003. Antioxidant capacity of phenolic phytochemicals from various cultivars of plums. *Food Chem.* 81, 321-326.

Kok, D., 2017. Grape growth, anthocyanin and phenolic compounds content of early ripening Cv. Cardinal table grape (*V. vinifera* L.) as affected by various doses of foliar biostimulant applications with gibberellic acid. *ErwerbObstbau* 58, 1-7.

Laszlo, J.C. & Saayman, D., 1990. Optimum harvesting stage for Sultanina as table grape. *Decid. Fruit Grow.* 40(3), 101-105.

Leifert, W.R. & Abeywardena, M.Y., 2008. Cardio protective actions of grape polyphenols. *Nutr. Res.* 28(11), 729-737.

Liang, Z., Sang, M., Fan, P., Wu, B., Wang, L., Duan, W. & Li, S., 2011. Changes of polyphenols, sugars, and organic acid in 5 *Vitis* genotypes during berry ripening. *J. Food Sci.* 76(9), 1231-1238.

Matthew, M.A. & Nuzzo, V., 2007. Berry size and yield paradigms on grapes and wines quality. *Acta Hort.* 754, 423-436.

Mattivi, F., Zulian, C., Nicolini, G. & Valenti L., 2002. Wine, biodiversity, technology, and antioxidants. *Ann. N.Y. Acad. Sci.* 957, 37-56.

Mazza, G., 1995. Anthocyanins in grapes and grape products. *Crit. Rev. Food Sci. Nutr.* 35(4), 341-71.

- Milkowska, K. & Strzelecka, H., 1995. Flos Hibisci – metody identyfikacji i ocena surowca. *Herba Polonica* 41(1), 11-16.
- Montealegre, R.R., Peces, R.R., Vozmediano, J.L.C., Gascueña, J.M. & Romero, E.G., 2006. Phenolic compounds in skins and seeds of ten grape *Vitis vinifera* varieties grown in a warm climate. *J. Food Compos. Anal.* 19, 687-693.
- Muñoz-Espada, A.C., Wood, K.V., Bordelon, B. & Watkins, B.A., 2004. Anthocyanin quantification and radical scavenging capacity of Concord, Norton, and Marechal Foch grapes and wines. *J. Agric. Food Chem.* 52(22), 6779-6786.
- Najda, A., 2017. Zmienność ontogenetyczna mięty (*Mentha species*) czynnikiem warunkującym zawartość składników bioaktywnych w surowcu. In Polish. University of Life Sciences, Lublin, 1, 178.
- Nampila, R., Bing-Shiun, Ch., Ching-Cheng, Ch. & YauShiang, Y., 2010. Effect of GA₃ and CPPU on berry size of seedless grapes. *Hort. NCHU* 35(3), 53-64.
- Orak, H.H., 2007. Total antioxidant activities, phenolics, anthocyanins, polyphenoloxidase activities of selected red grape varieties and their correlation. *Sci. Hort.* 111, 235-241.
- Peña-Neira, A., Dueñas, M., Duarte, A., Hernández, T., Estrella, I. & Loyola, E., 2004. Effects of ripening stages and of plant vegetative rigor on the phenolic composition of grapes (*Vitis vinifera* L.) cv. Cabernet Sauvignon in the Maipo Valley (Chile). *Vitis* 43(2), 51-57.
- Pezzuto, J., 2008. Grapes and human health: A perspective. *J. Agric. Food Chem.* 56(16): 6777-6784.
- Polish Norm PS, PN-90 A-75101/04 - Fruit and vegetable preserves. Sample preparation and physicochemical methods of examination. Determination of total acidity (in Polish).
- Polish Pharmacopoeia, 2002. VI, Wyd. PTFarm, Warszawa.
- Polish Pharmacopoeia, 2011. IX, Wyd. PTFarm, Warszawa.
- Polish Pharmacopoeia, 2014. X, Wyd. PTFarm, Warszawa.
- Rachna & Singh, S., 2013. Effect of gibberellic acid on periodical changes in bio-chemical composition of ber cv. Umran. *HortFlora Res. Spectrum* 2(1): 25-29.
- Reisch, B.I., Remaily, G.W., Pool, R.M. & Watson, J.P., 1986. 'Einset Seedless' grape. *HortScience* 21, 155-156.
- Scalbert, A., Manach, C., Morand, C., Remesy, C. & Jimenez, L., 2005. Dietary polyphenols and the prevention of diseases. *Crit. Rev. Food Sci. Nutr.* 45, 287– 306.
- Stevenson, D.E. & Hurst, R.D., 2007. Polyphenolic phytochemicals: Just antioxidants or much more? *Cell. Mol. Life Sci.* 64, 2900-2916.
- Tian, S., Wang, Y., Du, G. & Li, Y., 2011. Changes in contents and antioxidant activity of phenolic compounds during gibberellin-induced development in *Vitis vinifera* L. 'Muscat'. *Acta Physiol. Plant.* 33, 2467-2475.
- Tomás-Barberán, F.A. & Espin, J., 2001. Phenolic compounds and related enzymes as determinants of quality in fruits and vegetables. *J. Sci. Food Agric.* 81, 853-876.
- Topalovic, A. & Mikulic-Petkovsek, M., 2010. Changes in sugars, organic acids and phenolics of grape berries of cultivar Cardinal during ripening. *J. Food Agric. Environ.* 8(3), 223-227.
- Vislocky, L.M. & Fernandez, M.L., 2010. Biomedical effects of grape products. *Nutr. Rev.* 68(11), 656-670.
- Weston, L.A., 2005. Grape and wine tannins and phenolics, their roles in flavor, quality and human health. *Proc. 29th Annual New York Wine Industry Workshop, Month? 2005, New York, USA.* pp. 6 – 15.
- Yang, J., Martinson, T.E. & Liu, R.H., 2009. Phytochemical profiles and antioxidant activities of wine grape. *Food Chem.* 116, 332-339.
- Xia, E., Deng, G.F., Guo, Y.J. & Li, H.B., 2010. Biological activities of polyphenols from grapes. *Int. J. Mol. Sci.* 11, 622-646.