

# On the nucleolar and cytoplasmic RNA density during "cell dedifferentiation" represented by blastic transformation of human mature T lymphocytes – a cytochemical study

Karel Smetana, Petra Otevřelová, Ilona Jirásková, Ivan Kalousek

Institute of Hematology and Blood Transfusion, Prague, Czech Republic

**Abstract:** The present study was undertaken to provide information on the nucleolar and cytoplasmic density in specimens stained for RNA during "cell dedifferentiation" represented by blastic transformation of mature T lymphocytes. Nucleolar and cytoplasmic RNA's were visualized using a simple cytochemical method followed by computer assisted densitometry and size measurements of digitised images. An increased nucleolar and cytoplasmic RNA density accompanying the blastic transformation was significant after 48 hours of cultivation with phytohemagglutinin (PHA) when stimulated cells were characterized the largest nucleolar size reflecting S or G2 phase of the cell cycle. On the other hand, significantly larger ratio of the nucleolar to cytoplasmic density was noted only after a shorter cultivation when stimulated cells were presumably in the G1 phase. Thus the increased nucleolar and cytoplasmic RNA density together represented an accompanying phenomenon of the cell proliferation and cycling state. From the methodical point of view, the nucleolar and cytoplasmic RNA densitometry appeared as a simple as well as useful additional method to study "dedifferentiation" or various cell states at the single cell level. In addition, it was also interesting that the increase of the nucleolar diameter in stimulated cells was much larger than that of the nucleolar density. Such difference suggested that the RNA content in nucleoli was related mainly to their size.

**Key words:** nucleolar and cytoplasmic density, lymphocytes, blastic transformation

## Introduction

It is generally known that human mature resting T lymphocytes stimulated with phytohemagglutinin (PHA) "dedifferentiate" and transform back to proliferating blastic stage [1,3]. Such transformation of small mature resting to large cycling blastic cells is accompanied by nucleolar enlargement, which is related to the increased nucleolar biosynthetic activities of stimulated cells [1,3-5,24]. In addition, ring shaped nucleoli transform to large nucleoli with a relatively uniform distribution of RNA [1,15,18]. On the other hand, the information on the nucleolar and cytoplasmic RNA density in the course of blastic transformation is missing although it seems to be likely that it might be different.

The present study was undertaken to provide more information on the nucleolar and cytoplasmic density in specimens stained for RNA during "cell dedifferentiation" represented by blastic transformation of mature T lymphocytes. Nucleolar and cytoplasmic RNA were visualized using a simple cytochemical method followed by computer assisted densitometry and nucleolar diameter measurements of digitised images. The results indicated that an increasing density accompanying the blastic transformation was significant after 48 hours of cultivation with PHA when the cells were in S/G2 phase of the cell cycle. On the other hand, significantly larger ratio of the nucleolar to cytoplasmic RNA density was noted only after a shorter cultivation when stimulated cells were presumably in the late G1 phase.

## Materials and methods

Human lymphocytes were isolated from the peripheral blood of healthy volunteers using a discontinuous density gradient [2] of

**Correspondence:** K. Smetana, Institute of Hematology and Blood Transfusion, U nemocnice 1, Prague 2, Czech Republic 128 20; fax.: (+4202) 71977249, e-mail: karel.smetana@uhkt.cz

**Table 1.** Nucleolar and cytoplasmic density in specimens stained for RNA.

Time of cultivation (hrs)	Nucleoli (arbitrary density units)	Cytoplasm (arbitrary density units)	No/Cy	No D ( $\mu\text{m}$ )##
0	77.80 $\pm$ 22.43 <sup>a</sup>	75.12 $\pm$ 22.90	1.07 $\pm$ 0.29	1.0 $\pm$ 0.2
24	82.15 $\pm$ 12.96	66.90 $\pm$ 19.56 <sup>§</sup>	1.25 $\pm$ 0.31 <sup>*</sup>	2.1 $\pm$ 0.4 <sup>*</sup>
48	107.70 $\pm$ 19.96 <sup>*Ⓢ</sup>	95.5 $\pm$ 24.79 <sup>*Ⓢ</sup>	1.15 $\pm$ 0.24	2.8 $\pm$ 0.8 <sup>*</sup>

Legend: # – based on more than 40 measurements for each group; ## – see ref. [22]; \* – statistically different from 0 hrs using t-test ( $p < 0.009$ ); Ⓢ – statistically different from 24 hours using t-test ( $p < 0.001$ ); § – statistically different from nucleoli using t-test ( $p < 0.001$ ); a – mean and standard deviation; No/Cy – ratio of nucleolar to cytoplasmic RNA density; NoD – mean nucleolar diameter.

Histopaque (Sigma, St. Louis, MO USA). Cells,  $1 \times 10^6/\text{ml}$ , were suspended in medium RPMI 1640 supplemented with 10% heat-inactivated FBS and stimulated with 10  $\mu\text{g}/\text{ml}$  PHA-P from Sigma at 37°C in a 5% CO<sub>2</sub> humidified atmosphere for time frames up to 48 hrs [8,12]. Then, harvested cells were prepared for the light microscopic observations using a Shandon Cytospin 2 cytocentrifuge (Shandon Southern Products, UK) – 800 rpm for 10 min. and unfixed cytopsins were stained for RNA with acidified methylene blue at pH=5.3 [13,21].

Micrographs were captured with a Camedia digital photo camera C.4040 ZOOM (Olympus, Japan) placed on Jenalumar microscope (Zeiss, Germany) equipped with the double adapter to provide a larger magnification of resulting images. Then the images were processed and nucleolar size was measured with Quick Photogram (Olympus, Japan). Mean nucleolar diameters were determined on the screen at magnification  $\times 4300$  and were based on two measurements of the long and minor axis for each nucleolus [10,14,17]. It should be mentioned that the nucleolar size and distribution of RNA facilitated to estimate the cell cycle phase of studied cells [6,9,11,20,23,24].

The nucleolar and cytoplasmic density in specimens stained for RNA was measured after image conversion to grey scale using the NIH Image Program – Scion for Windows (Scion Corp., USA). The density was expressed in relative arbitrary units, which were calculated by subtracting measured density of nucleoli or cytoplasm from the density surrounding the measured cell. Such calculation and standardisation of arbitrary density units facilitated the comparison of results in various portions of cytopsins, which occasionally exhibited various artificial densities due to both preparation and staining techniques. This approach decreased artificial measurements and thus provided better results than the background adjusted to zero, which depended on the investigator. In contrast to the nucleolar size measurements, the nucleolar and cytoplasmic densities were measured at lower magnification using dry 50x objective to decrease the resolving power of the microscope. Such approach facilitated to cover a larger and more uniform density measurements of nucleolar and cytoplasmic regions stained for RNA.

## Results


### Nucleolar size and types

Quantitative data are in the Table 1. The nucleolar size during blastic transformation gradually increased. Maximal values of mean nucleolar diameter in stimulated lymphocytes were noted after 48 hrs of cultivation and apparently reflected S and G2 phase [6,9,11,16,20,23,24]. In resting mature lymphocytes representing cells in G0 or /early G1 phase, nucleoli were


small and mostly ring shaped (Fig. 1) [see also 3,6,18,23,24]. After 24 hours of cultivation in the presence of PHA, stimulated cells were characterised by enlarged nucleoli with more or less distinct nucleolonemas which were described in late G1 and especially S or G2 phase of the cell cycle (Fig. 2) [see also 3,6,15,23,24].

### Nucleolar and cytoplasmic density in specimens stained for RNA

For quantitative data see the Table 1. In control resting lymphocytes the nucleolar and cytoplasmic density was almost the same (Fig. 3). The nucleolar to cytoplasmic density ratio was about one. In transforming lymphocytes to blastic state (Figs. 4,5), the nucleolar density increased but such increase was significant only after 48 hours of cultivation with PHA. On the other hand, the cytoplasmic density was apparently smaller at first but then significantly increased. Therefore, the increased nucleolar to cytoplasmic density ration in stimulated cells significantly increased only after 24 hours of cultivation in the presence of PHA. It was also interesting that the nucleolar density in stimulated cells increased less dramatically than the nucleolar diameter (Fig. 6).



**Fig. 1.** Resting lymphocyte with a ring shaped nucleolus (arrow and insert), (magnification approx.  $\times 3500$ ). **Fig. 2.** Stimulated transformed and "dedifferentiated" blastic cell with a large nucleolus with a more uniform distribution of RNA (arrow), which shows nucleolonemas at larger magnification (insert), (magnification approx.  $\times 2100$ ).



**Fig. 3.** Nucleolar (No – thin black line) and cytoplasmic (Cy – thick black line) RNA density in a small resting lymphocyte. Nucleolus – arrow. The lower part of the Figure contains nucleolar (No) and cytoplasmic (Cy) densitograms. Numbers in densitograms indicate density values calculated by subtracting measured values from the background surrounding the investigated cell. The density scale is below densitogram (magnification approx.  $\times 2200$ ). **Fig. 4.** Nucleolar and cytoplasmic RNA density in a stimulated and transformed blastic cell after 24 hours of cultivation in the presence of PHA. For other legend see Fig. 3 (magnification approx.  $\times 1700$ ). **Fig. 5.** Nucleolar and cytoplasmic RNA density in a stimulated and transformed blastic cell after 48 hours of cultivation in the presence of PHA. For other legend see Fig. 3 (magnification approx.  $\times 1300$ ).


**Discussion**

The relatively increased nucleolar RNA density during blastic transformation (present study) is not surprising and represents a further phenomenon, which is related to the generally known increased nucleolar RNA transcription [3-5,7,15,16,23]. Moreover, the known transformation of ring shaped nucleoli in resting lymphocytes to enlarged nucleoli with less or more distinct nucleolonemas in transformed blastic cells is also in harmony with the increased nucleolar biosynthetic activity and return to the cell cycle [3,4,6,7,15,20,23, 24]. It should be added that the increase of the nucleolar diameter in stimulated cells was much larger than that of the nucleolar density. Such difference suggests that the RNA content in nucleoli depends mainly on their size. A similar RNA density of small and large nucleoli was also noted previously in leukaemia granulocytic progenitors [19].

The delayed increase of the RNA cytoplasmic density in stimulated cells seems to be interesting and resulted in an increased nucleolar to cytoplasmic RNA density ratio. Such delay might be easily explained by the preceding increased nucleolar RNA transcription and assembly of ribosomes before their transport to cytoplasm [3,5,7,23]. On the other hand, when biosynthetic activities of nucleoli in resting cells are repressed [4,6,7,15,24], the nucleolar and cytoplasmic RNA densities are similar but smaller (present results). At this occasion it should be mentioned that in stimulated and cycling blastic cells the nucleolar and cytoplasmic RNA densities were also similar. However,

they were significantly higher than in resting lymphocytes. In that case, both the nucleolar and cytoplasmic biosynthetic activities are known to substantially increase in comparison with resting cells [3-5,7,9,15].

From the methodical point of view the present study, regardless of the interpretation, clearly demonstrated that the cytochemical RNA densitometry was an useful procedure that might be complementary to



**Fig. 6.** Nucleolar diameter (NoD) and density (NoDn) in stimulated lymphocytes. Mean values of the nucleolar diameter are multiplied by 10 and mean values of the nucleolar density are divided by 10. Standard deviations are in the Table 1. Note that the increase of the nucleolar diameter was larger than that of the nucleolar density.

other methodical approaches facilitating to distinguish resting and stimulated – "dedifferentiated" cycling cells at the single cell level. In addition, such approach appeared to be useful in smear or cytopins specimens, in which the number of investigated cells might be very limited.

**Acknowledgement:** The present study was supported in part by the Research Development Project VZ 0002373601 of the Ministry of Health of the Czech Republic.

## References

- [ 1 ] Astaldi G, Lisiewicz J. *Lymphocytes*. Naples: Idelson; 1971.
- [ 2 ] Böyum A. Separation of leukocytes from blood and bone marrow. *Scan J Clin Lab Invest. Suppl.* 1968;21:9-109.
- [ 3 ] Busch H, Smetana K. *The nucleolus*. New York: Academic Press; 1970.
- [ 4 ] Carson DA. *Biochemistry and function of lymphocytes and plasma cells*. In: Williams WJ, Beutler E, Erslev AJ, Lichtman MA, eds. *Hematology*. New York: McGraw-Hill; 1983: 903-908.
- [ 5 ] Cline MJ. *The white cell*. Cambridge: Harvard University Press; 1975.
- [ 6 ] Gani R. The nucleoli of cultured human lymphocytes. *Exp Cell Res.* 1976;97:249-258.
- [ 7 ] Henry PH, Levin MJ, Levin AS. *Composition and biochemistry of lymphocytes and plasma cells*. In: Williams WJ, Beutler E, Erslev AJ, Rundles RW, eds. *Hematology*. New York: McGraw-Hill; 1972:782-790.
- [ 8 ] Kalousek I, Krizkova P. Lymphocyte mitogenic transformation is accompanied by phosphorylation of the nucleolar transcription factor UBF. *Cell Mol Biol. (Noisy-le-grand)* 2000; 46:163-1171.
- [ 9 ] Kwiatkowska M, Maszewski J. Functional and structural heterogeneity of nucleoli: the dependence of the activity of transport of newly synthesized rRNA on nucleolar size and phase of the cell cycle. *Folia Histochem Cytochem.* 1985;23:117-126.
- [ 10 ] Monge JM, Val-Bernal JF, Buelta L, Garcia-Castrillo L, Asensio L. Selective nuclear morphometry as prognostic factor of survival of renal carcinoma. *Histol Histopathol.* 1999;14:119-123.
- [ 11 ] Nagl W. *Zellkern und Zellzyklen*. Stuttgart: Verl. Eugen Ulmer; 1976
- [ 12 ] Nowell PC Differentiation of human leukaemic leukocytes in tissue culture. *Exp Cell Res.* 1960;19:267-277.
- [ 13 ] Ochs RL. Methods used to study structure and function of the nucleolus. *Methods Cell Biol.* 1998;53:303-301.
- [ 14 ] Politi EN, Lazaris AC, Kavantzias A, Kountselini H. Comparison between morphometry and immunostaining of malignant cells in no-small cell lung cancer. *Anal Quant Cytol Histol.* 2003;25:169-176.
- [ 15 ] Raška I, Koberna K, Malinský J, Fidlerová H, Mašata M. The nucleolus and transcription of ribosomal genes. *Biol Cell.* 2004;96:579-594.
- [ 16 ] Schnedl W, Schnedl M. Nukleolus Zahl und Grösse während des Zellzyklus. *Z. Zellforsch.* 1972;126:374-382.
- [ 17 ] Setala L, Lipponen P, Kosma VM, Marin S, Eskelinen M, Syrjanen K, Alhava E. Nuclear morphometry as predictor of disease outcome in gastric cancer. *J Pathol.* 1977;181:46-50.
- [ 18 ] Smetana K. Nucleoli in maturing blood cells. *Top Rev Haematol.* 1980;1:115-137.
- [ 19 ] Smetana K, Klamová H, Mikulenkova D, Pluskalovi M, Hrkal Z. The nucleolar size and density in human early granulocytic progenitors – myeloblasts. *Europ J Histochem.* 2006; 50:119-124.
- [ 20 ] Smetana K, Kuzelova K, Zapotocky M, Starkova J, Hrkal Z, Trka J. Mean diameter of nucleolar bodies in cultured human leukemic myeloblasts is related to the S and G2 phase of the cell cycle. *Europ. J Histochem.* 2007;51:269-274.
- [ 21 ] Smetana K, Lejnar J, Potměšil M. A further contribution to the demonstration of RNA and nucleoli of blood cells in smear preparations. *Folia Haematol.* 1969;91:381-384.
- [ 22 ] Smetana K, Otevřelová P, Kalousek I. Heterochromatin density in the course of cell "dedifferentiation" represented by blastic transformation of human mature T lymphocytes. *Appl Biomed.* 2007;5:189-193.
- [ 23 ] Wachtler F, Schwarzacher HG, Ellinger A. The influence of the cell cycle ion structure and number of nucleoli in cultures human lymphocytes. *Cell Tissue Res.* 1982;225:155-163.
- [ 24 ] Wachtler F, Stahl HG. The nucleolus. *Micron.* 1983;24:473-505.

Submitted: 29 November, 2007

Accepted after reviews: 2 June, 2008