

Increased apoptosis in human knee osteoarthritis cartilage related to the expression of protein kinase B and protein kinase $C\alpha$ in chondrocytes

Qingzhi Chen^{1,2} Bing Zhang³, Tingting Yi³, Chun Xia¹

¹Zhongshan Hospital, Xiamen University, Fujian, China

²The Third Hospital of Xiamen, Fujian, China

³Medical School, Xiamen University, Fujian, China

Abstract: Protein kinase B (Akt) and protein kinase $C\alpha$ (PKC α) play important roles in the regulation of cell apoptosis. The aim of this study was to investigate the expression of Akt and PKC α in chondrocytes of human knee osteoarthritic (OA) cartilage, further evaluating their role in chondrocyte apoptosis during OA progression. Human knee OA cartilages were obtained from 38 patients undergoing knee arthroplasty, which is the medium-late stage of OA. Healthy knee cartilages were obtained from 11 amputees. The samples taken from the condyle of femur were collected routinely for morphological, immunohistochemical and Western blot detection, respectively. Light microscopy and laser-scanning confocal microscopy were used for morphological observation. The optical density with computer image analysis evaluated the intensity of immunohistochemical reaction of Akt and PKC α in OA cartilage. Western blot detected the protein expression levels. The results indicated that Akt and PKC α were involved in OA progression, along with the increase of cell apoptosis. In OA cartilage, Akt decreased ($p < 0.05$) and PKC α increased ($p < 0.05$). There was a negative correlation and interaction between Akt and PKC α ($r = -0.8$). These results demonstrated that both Akt and PKC α are related to increased chondrocyte apoptosis in human OA cartilage. The correlation between human OA progression, the role of Akt and PKC α , and chondrocyte apoptosis allows for new therapeutic strategies to be considered. (*Folia Histochemica et Cytobiologica* 2012, Vol. 50, No. 1, 137–143)

Key words: Akt, PKC α , correlation, chondrocytes, human OA cartilage

Introduction

Osteoarthritis (OA) is characterized by a slow focal destruction of articular cartilage, causing a roughening and thinning of the weight-bearing regions of the articular surface, resulting in progressive immobility and pain. Chondrocytes, as only a single cell type in articular cartilage, is responsible for the homeostasis of the tissue by synthesizing extracellular matrix that

surrounds the cells and provides the important biophysical characteristics of the tissue [1]. Recently, several studies have provided evidence that chondrocyte apoptosis and dedifferentiation play an important role in OA [1, 2]. In addition, apoptosis has been positively correlated with the severity of cartilage destruction and with matrix depletion in human specimens [3]. However, the intrinsic signaling pathways regulating cell apoptosis are not fully understood.

Protein kinase C (PKC) is a subfamily of serine/threonine kinases involved in the regulation of various cell responses, such as cell proliferation, differentiation, and apoptosis. PKC α , one of 12 types of PKC isoforms, mediates a number of cellular fates in some mammal cell lines, including normal articular chondrocytes [4]. In addition, Akt (also known as protein kinase B), another subfamily of serine/threonine

Correspondence addresses:

B. Zhang, Medical School, Xiamen University, Fujian, China;
tel.: + 592 2188675, fax: + 86 592 2188680;
e-mail: cristal66@xmu.edu.cn

C. Xia, Zhongshan Hospital, Xiamen University, Fujian, China;
tel.: + 592 2993080, fax: + 86 592 5921461;
e-mail: chunxia@xmu.edu.cn

kinases, has emerged as a crucial regulator of diverse cellular processes, such as apoptosis, survival, proliferation, differentiation and metabolism [5–7]. For instance, activated by a variety of stimuli through a phosphorylation mechanism, Akt is known to be a potent inhibitory signal for apoptosis in several kinds of cells [5–8]. Especially, Akt1 in osteoblasts and osteoclasts controls bone remodeling [8] and the actin cytoskeleton mediates NO-induced regulatory effects in chondrocytes by modulating the down-regulation of PI3K-Akt [9]. Taken together, both PKC α and Akt are involved in the regulation of cellular fates. However, it is not clear whether PKC α and Akt also regulate the cell responses of chondrocytes in OA cartilage such as cell apoptosis, which is one of the crucial elements in OA progression.

The present study was undertaken to investigate whether PKC α and Akt signal pathways play roles in OA progression, with chondrocyte apoptosis occurring in OA cartilage, and, if so, further to investigate the correlation between them.

Our results indicated differing roles for Akt and PKC α , along with an increase of chondrocyte apoptosis in OA cartilage, and there was a negative correlation and interaction between Akt and PKC α . This study might provide a molecular basis for studying the regulatory mechanism of chondrocyte apoptosis in human OA cartilage, and a novel target for the treatment of OA.

Materials and methods

Human articular cartilage. After approval by the Ethical Committee of Zhongshan hospital, Xiamen University, human OA cartilage was obtained from 38 patients (55–67 years, average age 63.2 years) undergoing knee arthroplasty, which is the medium-late stage of OA (Mankin score 8–12, average value 10.3) [10]. Healthy knee cartilage was obtained from 11 amputees (31–48 years, average age 43.1 years), of whom four had been amputated because of bone tumor, and seven had been amputated due to trauma. All samples were taken from the condyle of femur. Each sample was divided into two parts: one was fixed in paraformaldehyde and embedded for observation using a light microscope and a laser-scanning confocal microscope, and the other was preserved in the freezer (at -80°C) for protein detection.

Histomorphology. The sample was fixed in 4% paraformaldehyde for 48 hr, decalcified in 15% EDTA (pH 7.0, 37°C) for two weeks, and then paraffin-embedded for further routine histological preparation. Serial frontal $4\ \mu\text{m}$ sections were observed under a light microscope in order to evaluate matrix abundance and cellularity after hematoxylin and eosin staining. In normal articular cartilage, the zones are

easy to delineate, and their cells are clearly distinguishable, including four zones: superficial zone, transitional zone, radical zone, and calcification. Since the superficial cartilage zone had been substantially destroyed in the OA group, only the chondrocytes in the transitional zone and the radical zone were observed in this study.

Immunohistochemistry. As described in the manufacturer's instructions (MAIXIN.BIO China), the sections were incubated overnight at 4°C with primary antibody: Akt (1:100 dilutions, Cell Signaling Technology, Danvers, MA, USA) or PKC α (1:150 dilutions, Cell Signaling Technology, Danvers, MA, USA) and subsequently, with secondary antibody (1:400) for 60 min, respectively. Diaminobenzidine (DAB) was used to visualize the immunohistochemical reaction, followed by counterstaining with hematoxylin. Finally, dark brown cells were considered to be positive under a light microscope and were counted throughout the microscopically magnified fields ($\times 400$) of each articular cartilage section. According to previous studies [11–13], the average optical density (OD) of cells is the symbol of protein expression, so it was detected by ImagePro Plus6.0 system in each cartilage zone, followed by statistical analysis.

TUNEL. For *in situ* visualization of apoptotic cells, the TUNEL method was performed according to the manufacturer's instructions with in Situ Cell Death Detection Kit, POD (Roche Diagnostics). In TUNEL staining, dark brown cells were considered to be positive, and were counted throughout microscopically magnified fields ($\times 400$) of each articular cartilage section. The percentage of positive cells was analyzed by SPSS v.15.0 for Windows.

Laser-scanning confocal microscopy. Articular cartilage was fixed in 4% paraformaldehyde. For staining endogenous Akt and PKC α proteins, sections were incubated with anti-Akt or anti-PKC α antibody (1:100 or 1:150 dilutions, Cell Signaling Technology, Danvers, MA, USA) followed by FITC or Texas Red conjugated secondary antibodies (1:200 dilutions, Santa Cruz Biotechnology, Inc. CA, USA). Cells were stained by 4',6-diamidino-2-phenylindole (DAPI, 50 mg/ml, Sigma) to visualize the nuclei simultaneously [14]. The stained cells were finally visualized under laser-scanning confocal microscopy (Leica Tcs Sp2 SE).

Western blot and immunoprecipitation. Articular cartilage was lysed in RIPA buffer (10 mmol/L Tris pH7.4, 150 mmol/L NaCl, 1% TrionX-100, 1% deoxycholic acid, 0.1% SDS, 5 mmol/L EDTA pH 8.8, 1 mmol/L PMSE, 10% Cocktail (Roche), 1% dithiothreitol) for 30 min at 4°C [14]. Lysates were centrifuged at $12,000 \times g$ for 30 min at 4°C . Protein concentration was determined using the Bio-Rad protein assay system according to the manufacturer's instructions (Bio-Rad, Hercules, CA, USA). Protein extracts were electrophoresed on 8% or 10% denaturing gel and electroblot-

ted onto a nitrocellulose membrane. The membranes were incubated with different antibodies as required at 4°C overnight, followed by the addition of a corresponding secondary antibody at room temperature for 1 hr. An ECL kit (Pierce, Thermo Scientific, Rockford, IL, USA) was used to detect the antibody reactivity.

For immunoprecipitation [14], protein extracts were incubated with the appropriate antibody for 1 hr, and were subsequently incubated with protein A-sepharose beads (Sigma) for 1 hr. The protein-antibody complexes that were recovered on beads were subjected to Western blot analysis as described above.

Statistical analysis. Data was expressed as mean \pm standard deviation. Statistical analyses were done using SPSS v.15.0 for Windows. Independent two-sample *t*-test and paired two-sample *t*-test were used for the comparison of protein expression between OA group and normal group. P values of less than or equal to 0.05 were considered significant. Pearson correlation test was used for the correlation between Akt and PKC α .

Results

Chondrocyte apoptosis increases in human OA cartilage

The histological differences between the human OA group and the normal group were initially observed, and it was found that the zones in the articular cartilage of the OA group were irregular compared to the normal group because of the cellular changes that took place (Figure 1A). These changes included the articular cartilage becoming thinner, and the superficial zone being virtually destroyed. Moreover, the arrangement of chondrocytes throughout transitional and radical zones were irregular, following the decrease of the chondrocyte number and the rupture of the tidemark, which is at the junction of the articular hyaline cartilage and adjacent subchondral bone (Figure 1A).

The decrease of chondrocyte number is crucial to the loss of articular cartilage in OA. To confirm whether chondrocyte apoptosis was present in human OA cartilage, we performed the TUNEL method in the OA and the normal group. TUNEL is a common method for detecting DNA fragmentation that results from apoptotic signaling cascades. The assay relies on the presence of nicks in the DNA which can be identified by terminal deoxynucleotidyl transferase, an enzyme that will catalyze the addition of dUTPs that are secondarily labeled with a marker [15]. Although TUNEL staining cells were present in all zones in the normal and in the OA group, there was a significant increase in the number of TUNEL staining cells in the OA group compared to the normal group

(Figure 1B, Table 1, $p < 0.05$). Moreover, the number of TUNEL staining cells in the transitional zone was significantly higher than that in the radical zone (Figure 1B, Table 1, $p < 0.01$).

Akt and PKC α are involved in human OA progression

The result of immunohistochemistry showed that the average OD of the heavy staining cells for Akt in the OA group was significantly lower than that of the normal group (Figure 2A, Table 1, $p < 0.05$). Like the TUNEL staining result, the average OD of the cells for Akt in the transitional zone was significantly higher than that in the radical zone, whether in the OA or the normal group (Figure 2A, Table 1, $p < 0.01$).

Furthermore, Western blot results showed that the expression level of total Akt in the OA group decreased compared to that in the normal group, while the P-Akt increased in the OA group (Figure 3). These findings confirmed that Akt and its phosphorylation are involved in OA progression. Nevertheless, the result of immunohistochemistry showed that the average OD for PKC α in the OA group is significantly higher than in the normal group (Figure 2B, Table 1, $p < 0.05$). In addition, the expression of PKC α is more prominent in the transitional zone than the radical zone, whether in the OA or the normal group (Figure 2B, Table 1, $p < 0.01$). The result of the Western blot analysis also showed that the expression level of PKC α increased in the OA group compared to that in the normal group (Figure 3).

Correlation between Akt and PKC α in human OA progression

The distribution of Akt and PKC α in chondrocytes was initially observed under laser-scanning confocal microscopy. Both Akt and PKC α mainly localized in the cytoplasm of chondrocyte in both the OA group and the normal group (Figures 4A, C for Akt, Figures 4B, D, for PKC α). The distribution of Akt and PKC α did not alter in OA progression. Then, as shown in Figure 5, Akt bound to PKC α , but the degree of their interaction in the OA group was similar to that in the normal group (Figure 5). Furthermore, there exists a negative correlation between Akt and PKC α by analyzing the average OD of Akt and PKC α (Table 1, $r = -0.8$).

Discussion

Chondrocyte apoptosis has been implicated as one of the etiological features in articular cartilage degradation during OA development [1–3]. In this study, Akt and PKC α participate in human OA progression,




Figure 1. Comparison of histological and chondrocyte apoptosis in normal and OA group. (A) Histological changes in normal and OA group. Left: normal group, Right: OA group (magnification $\times 50$). (B) Apoptotic chondrocytes in normal and OA group. Left: TUNEL staining chondrocytes (brown) (magnification $\times 200$, $\times 400$). Right: percentage of apoptotic chondrocytes in normal and OA group ($*p < 0.01$)




Figure 2. Expression of Akt and PKC α by immunohistochemical staining in normal and OA group. (A) Expression of Akt in normal and OA group. Left: Akt positive chondrocytes (magnification $\times 400$). Right: Histogram of the average OD of chondrocytes ($*p < 0.01$). (B) Expression of PKC α in normal and OA group. Left: PKC α positive chondrocytes (magnification $\times 400$). Right: Histogram of the average OD of chondrocytes ($*p < 0.01$)

Table 1. Expression of Akt, PKC α and chondrocyte apoptosis in normal and OA group

	Akt		PKC α		TUNEL	
	Normal	OA	Normal	OA	Normal	OA
Transitional	*1176.74 \pm 135.39	*356.5 \pm 65.87	*155.97 \pm 17.56	*621.47 \pm 103.19	*13.4 \pm 3.2%	*34.5 \pm 3.7%
Radical	775.33 \pm 122.58	18620 \pm 40.93	88.26 \pm 26.29	175.71 \pm 114.26	6.2 \pm 1.8%	21.8 \pm 2.8%
Total	#1076.03 \pm 140.54	271.18 \pm 102.22	126.16 \pm 41.46	#398.59 \pm 39.57	10.7 \pm 2.6%	#28.4 \pm 3.6%

* $p < 0.01$, # $p < 0.05$, $R = -0.802$




Figure 3. Expression level of Akt, p-Akt and PKC α in normal and OA group. Protein extracts from articular cartilages were prepared as described in the 'Materials and methods' section, and then analyzed by Western blot analysis with an antibody against Akt or P-Akt or PKC α . The blots were normalized to an endogenous protein (GAPDH). The data is representative of ten independent experiments, each yielding similar results

along with the increase of chondrocyte apoptosis. There exists a negative correlation and interaction between Akt and PKC α .

Firstly, apoptosis is significantly higher in OA cartilage than in normal cartilage (Figure 1B, Table 1, $p < 0.05$), consistent with the belief that chondrocyte apoptosis is an indication of OA cartilage [1–3, 16]. Furthermore, we found that the increase of chondrocyte apoptosis in the transitional zone was higher than that in the radical zone in the OA cartilage, something also found in other authors' reports [17].

This increase of chondrocyte apoptosis would thus explain the failure of maintenance of the cartilage matrix and the loss of articular cartilage in OA. In 2004, Roach et al. [18] suggested the term 'chondroptosis' to define the type of cell death present in articular cartilage, which included the presence of both apoptotic (late stage) and autophagic (early stage) processes [19]. Here, the chondrocytes obtained are most likely undergoing chondrocyte apoptosis of 'chondroptosis', due to the fact that they were obtained from patients in the medium-late stage for undergoing knee arthroscopy (Mankin score 8–12, average value 10.3).




Figure 4. Subcellular distribution of Akt and PKC α in the chondrocytes of normal and OA group. As described in the 'Materials and methods' section, the cells were immunostained with Akt or PKC α -antibody, followed by Texas-conjugated secondary antibody (A & C for Akt) and FITC-conjugated secondary antibody (B & D, for PKC α), respectively. The nuclei were simultaneously indicated by DAPI staining. The fluorescent images were observed under a laser-scanning confocal microscope. The data is representative of 15 independent experiments, each yielding similar results

The activation of the chondrocyte by various extracellular factors resulted in an imbalance between the synthesis and degradation of extracellular matrix in OA progression. All these factors act through specific receptors that transmit the signals to the nucleus to activate the transcription of matrix metalloproteinases and inflammatory genes. Targeting these intrinsic signaling pathways in osteoarthritis is a novel approach to modulate this imbalance. In 2008, Keisuke et al. [20] reported that the activation of Akt in embryonic chondrogenesis enhanced chondrocyte proliferation. Other articles also have demonstrated enhanced apoptosis in osteoblasts, testis cells, thymocytes, and embryo fibroblasts in Akt1 $^{-/-}$ mice [8].




Figure 5. Interaction between Akt and PKC α in normal and OA group. As described in the ‘Materials and methods’ section, tissue lysates were immunoprecipitated with Akt or PKC α antibody, and then subjected to Western blotting with PKC α or Akt antibody. IgG expression served as a control for indicating similar proteins in each lane. The same lysates were applied to ascertain the position and expression of PKC α and Akt by Western blotting (Input). The data is representative of five independent experiments, each yielding similar results

In this study, Akt not only declined in OA cartilage, but also its decrease was higher in the transitional zone, consistent with the increase of chondrocyte apoptosis in the transitional zone (Figure 2A, Table 1). In addition, Western blot analysis also indicated the decrease of Akt level in the OA cartilage (Figure 3).

Taken together, we suggest that Akt decline accompanied by chondrocyte apoptosis hastens human OA progression. However, the high expression level of phosphorylation of Akt (P-Akt) in the OA group (Figure 3) might act as a compensatory role to protect chondrocytes from the apoptosis in the OA cartilage. The latest paper by García S et al. [21] also reported that the endogenous phosphorylation of Akt protects rheumatoid arthritis (RA) fibroblast such as synovial cells against the apoptosis induced by Fas. Elevated production of TRB3, an inhibitor of Akt activation, in osteoarthritic chondrocytes, would increase chondrocyte death in OA cartilage [22]. Therefore, the decrease of Akt level makes for the increase of chondrocyte apoptosis in the OA group, and Akt may be a novel target for new therapeutic strategies in OA.

Recently, growing evidence has indicated that PKC α not only plays an essential role in chondrocyte apoptosis [5, 9], but is also involved in the early stage of OA progression [23, 24]. Ca²⁺influx-dependent

PKC α -MAPK signaling cascades have been implied to be activated in the experimental dog OA model system, indicating that the PKC α pathway may mediate development and progression of OA [23]. PKC α appears in chondrocytes in the early stage of OA progression [24]. Here, the increase of chondrocyte apoptosis in the late stage of OA (Figures 2B, 3, Table 1) and the high expression level of PKC α , showed that the increase of PKC α might promote the chondrocyte apoptosis in OA cartilage, suggesting its involvement in the total progression of OA.

Interaction between Akt and PKC α has been reported [9, 11, 14]. Previous studies, both in our lab and others, have suggested that Akt could be regulated by PKC α negatively in cancer cells [14, 25]. On the other hand, the overexpression of PKC also could stimulate Akt activity and suppress cytokine-dependent apoptosis [26]. Thus, PKC α might be a potential regulator, positively or negatively, for Akt. In the current case, the co-localization of Akt and PKC α in the cytoplasm of chondrocyte (Figure 4) provided a clue to the next step: a study on the interaction between them. Unfortunately, the alternation of the bind between them in OA progression could not be observed in this study (Figure 5). Combined with the analysis in Table 1, we suggest that Akt might be associated with PKC α in OA chondrocyte, and PKC α may regulate Akt negatively. It is necessary to study their phosphorylation in future in order to elucidate their interaction in OA progression. It also implied that it is crucial to pay attention to the antagonism between them in therapy of OA and interesting to delineate further the correlation between Akt with PKC α in OA cartilage.

However, there exists some limitation in this study because of the difficulty in obtaining human OA cartilage. It is regrettable not to be able to pay more attention to studying the relationship between the range age differences and the expression of Akt or PKC α in the normal and the OA group.

In summary, our study demonstrated that both Akt and PKC α related to the increased chondrocyte apoptosis in human OA cartilage. This is a very significant finding, indicating that Akt and PKC α play differing roles in human OA progression. It may provide a molecular basis for molecular diagnosis and therapeutic strategies. For instance, the change of the ratio of Akt/PKC α might provide reference points for doctors in diagnosing the degree of OA or determining therapeutic strategies.

Thus, the correlation between human OA progression, the role of Akt and PKC α , and chondrocyte apoptosis allows for the possibility of new therapeutic strategies.

Acknowledgements

The Natural Science Foundation of Fujian, China (No.C0910644), the National Natural Science Foundation of China (No.81072015) and the Natural Science Foundation of Xiamen, China (No. 3502Z20084015) supported this study.

References

1. Kuhn K, D'Lima DD, Hashimoto S et al. Cell death in cartilage. *Osteoarthritis Cartilage*. 2004;12:1–16.
2. Kim HA, Blanco FJ. Cell death and apoptosis in osteoarthritic cartilage. *Curr Drug Targets*. 2007;8:333–345.
3. Kim HA, Suh D-1, Song Y-W. Relationship between chondrocyte apoptosis and matrix depletion in human articular cartilage. *J Rheumatol*. 2001;28:2038–2045.
4. Kim SJ, Kim HG, Oh CD et al. p38 kinase-dependent and -independent inhibition of protein kinase C zeta and -alpha regulates nitric oxide-induced apoptosis and dedifferentiation of articular chondrocytes. *J Biol Chem*. 2002;277:30375–30381.
5. Brazil DP, Hemmings BA. Ten years of protein kinase B signaling: a hard Akt to follow. *Trends Biochem Sci*. 2001;26:657–664.
6. Franke TF, Hornik CP, Segev L et al. PI3K/Akt and apoptosis: size matters. *Oncogene*. 2003;8:22:8983–8998.
7. Chen WS, Xu PZ, Gottlob K et al. Growth retardation and increased apoptosis in mice with homozygous disruption of the Akt1 gene. *Genes Dev*. 2001;15:2203–2208.
8. Kawamura N, Kugimiya F, Oshima Y et al. Akt1 in osteoblasts and osteoclasts controls bone remodeling. *PLoS ONE*. 2007;2:e1058.
9. Kim SJ, Hwang SG, Kim IC et al. Actin cytoskeletal architecture regulates nitric oxide-induced apoptosis, dedifferentiation, and cyclooxygenase-2 expression in articular chondrocytes via mitogen-activated protein kinase and protein kinase C pathways. *J Biol Chem*. 2003;278:42448–42456.
10. Mankin HJ, Dorfman H, Lippiello L et al. Biochemical and metabolic abnormalities in articular cartilage from osteoarthritic human hips. II. Correlation of morphology with biochemical and metabolic data. *J Bone Joint Surg Am*. 1971;53:523–537.
11. Oh CD, Chun JS. Signaling mechanisms leading to the regulation of differentiation and apoptosis of articular chondrocytes by insulin-like growth factor-1. *J Biol Chem*. 2003;19:278:36563–36571.
12. Słuczana-Głąbowska S, Laszczyńska M, Wylot M et al. Morphological and immunohistochemical comparison of three rat prostate lobes (lateral, dorsal and ventral) in experimental hyperprolactinemia. *Folia Histochem Cytobiol*. 2010;48:447–454.
13. Wang WZ, Liu HO, Wu YH et al. Estrogen receptor α (ER α) mediates 17 β -estradiol (E2)-activated expression of HBO1. *J Exp Clin Cancer Res*. 2010;29:140–146.
14. Zhang B, Xia C. 12-O-tetradecanoylphorbol-1, 3-acetate induces the negative regulation of protein kinase B by protein kinase Calpha during gastric cancer cell apoptosis. *Cell Mol Biol Lett*. 2010;15:377–394.
15. Negoescu A, Guillermet C, Lorimier P et al. Importance of DNA fragmentation in apoptosis with regard to TUNEL specificity. *Biomed Pharmacother*. 1998;52:252–258.
16. Sharif M, Whitehouse A, Sharman P et al. Increased apoptosis in human osteoarthritic cartilage corresponds to reduced cell density and expression of caspase-3. *Arthritis Rheum*. 2004;50:507–515.
17. D'Lima D, Hermida J, Hashimoto S et al. Caspase inhibitors reduce severity of cartilage lesions in experimental osteoarthritis. *Arthritis Rheum*. 2006;54:1814–1821.
18. Roach HI, Aigner T, Kouri JB. Chondroptosis: a variant of apoptotic cell death in chondrocytes? *Apoptosis*. 2004;9:265–277.
19. Maiuri MC, Zalckvar E, Kimchi A et al. Self-eating and self-killing: crosstalk between autophagy and apoptosis. *Nat Rev Mol Cell Biol*. 2007;8:741–752.
20. Keisuke Kita, Tohru Kimura, Norimasa Nakamura et al. PI3K/Akt signaling as a key regulatory pathway for chondrocyte terminal differentiation. *Genes to Cells*. 2008;13:839–850.
21. García S, Liz M, Gómez-Reino JJ et al. Akt activity protects rheumatoid synovial fibroblasts from Fas-induced apoptosis by inhibition of Bid cleavage. *Arthritis Research & Therapy*. 2010;12:R33.
22. Cravero JD, Carlson CS, Im HJ et al. Increased expression of the Akt/PKB inhibitor TRB3 in osteoarthritic chondrocytes inhibits insulin-like growth factor 1-mediated cell survival and proteoglycan synthesis. *Arthritis Rheum*. 2009;60:492–500.
23. Boileau C, Martel-Pelletier J, Brunet J et al. PD-0200347, an alpha2delta ligand of the voltage gated calcium channel, inhibits in vivo activation of the Erk1/2 pathway in osteoarthritic chondrocytes: a PKCalpha dependent effect. *Ann Rheum Dis*. 2006;65:573–580.
24. Yeh CC, Chang HI, Chiang JK et al. Regulation of plasminogen activator inhibitor 1 expression in human osteoarthritic chondrocytes by fluid shear stress: role of protein kinase Calpha. *Arthritis Rheum*. 2009;60:2350–2361.
25. Doornbos RP, Theelen M, Van Der Hoeven PC et al. Protein kinase Czeta is a negative regulator of protein kinase B activity. *J Biol Chem*. 1999;274:8589–8596.
26. Li W, Zhang J, Flechner L et al. Protein kinase C-alpha overexpression stimulates Akt activity and suppresses apoptosis induced by interleukin 3 withdrawal. *Oncogene*. 1999;6564–6572.

Submitted: 11 April, 2011

Accepted after reviews: 14 November, 2011