

# Early trabeculation and closure of the interventricular foramen in staged human embryos

M. Rauhut-Klaban, M. Bruska, W. Woźniak

Department of Anatomy, Medical University, Poznań, Poland

[Received 3 December 2007; Accepted 15 January 2008]

*Internal differentiation of the ventricles was studied in staged serially sectioned human embryos of developmental stages 13–19 (postovulatory days 32–46). At stage 13 the trabeculation of both ventricles was advanced and the muscular part of the interventricular septum well marked. Dorsal and ventral endocardial cushions were fused and the atrioventricular canal was divided into two parts. In embryos at stage 18 the membranous interventricular septum was developing and the interventricular foramen was obliterated. At stage 19 the membranous part of the interventricular septum was becoming more cellular in structure. (Folia Morphol 2008; 67: 13–18)*

**Key words:** human embryology, embryonic period, heart development, differentiation of ventricles

## INTRODUCTION

During intrauterine development the heart is the first organ to begin mechanical function and this is initiated before structural organogenesis is complete. This raises the possibility that its early mechanical function affects its own morphogenesis [17]. Although morphogenesis of the heart varies between species the general steps during its development are common. These steps involve the following: determination and formation of cardiomyocytes, formation of the heart tube, looping, growth of the chambers, formation of the endocardial cushions, valvulogenesis, and septation.

Valvular and septal defects are among the most common and most deleterious of all cardiac malformations [15]. The endocardial cushions play a central role in cardiac septation and valve formation [13, 24]. In the development of the myocardium and endocardial cushions an important role is played in turn by the neural crest cells, which migrate from

the cardiac neural crest region. Ablation of this region results in myocardial dysfunction and septal malformation [10, 20, 32].

Trabeculation and septation are interlinked events. Trabeculation has been implicated in enhancing contractility, ventricular septation and intraventricular conduction and in helping to direct blood flow before septation [7, 8, 11, 16–18].

Although there has been considerable research into human embryonic [2, 3, 5, 9, 16, 21–23, 25] and foetal hearts [1], as well as those of animals [12, 14, 29], there are still controversies as to the development of the interventricular septum and the exact timing of the differentiation of the ventricles, and differing conclusions have been reached regarding the time of closure of the interventricular foramen [5, 9, 16, 23].

Understanding of the mechanics of the development itself will provide the information required for the proper interpretation of postnatal morphology [2–4]. The aim

of the present study is to establish the exact time of closure of the interventricular septum and the formation of the trabeculae in staged human embryos.

### MATERIAL AND METHODS

Twelve human embryos ranging in age from 32 to 46 days were studied using a light microscope. The embryos belonged to the collection of the Department of Anatomy of the Medical University of Poznań. The embryos studied were from stage 13 to stage 19 (Table 1). They were sectioned serially in the sagittal, frontal, and horizontal planes and stained according to routine histological methods and with silver. From each embryo histological sections relevant to this study were selected and photographed.

### RESULTS

In the embryos at stage 13 the dorsal and ventral endocardial cushions were fused and the atrioventricular canal was divided into right and left atrioventricular canals. The right and left atrioventricular sulci were evident and the interventricular sulcus was deep. The muscular part of the interventricular septum was well marked (Fig. 1). Each ventricle consisted of the atrioventricular inlet, the trabeculated part and the outlet, which was directed to the bulbus cordis, in which the bulbar ridges are evident (Fig. 2).


The dorsal atrioventricular cushion descended downward to the ventricular wall (Fig. 3), supply-

**Table 1.** Developmental stage, age in postovulatory days, and plane of section of the embryos examined

Catalogue No.	Stage	Age [days]	Section
B171	13	32	Frontal
B218	13	32	Sagittal
B174	13	32	Horizontal
B272	14	33	Sagittal
B195	14	33	Sagittal
B115	15	36	Frontal
PJK21	15	36	Sagittal
PJK8	16	39	Horizontal
PJK2	17	41	Horizontal
BŁ4	18	44	Sagittal
B66	19	46	Horizontal
B173	19	46	Horizontal

ing evidence that the dorsal endocardial cushion contributes to the formation of the interventricular septum. In the peripheral part of the ventricular wall the compact myocardium was formed (Fig. 2).


The interventricular flange partially overlapped the interventricular septum and the atrioventricular canal was located to the left. The outflow tract, like the bulbus cordis, originated from the right ventricle. The bulbus continued as a truncus arteriosus. The




**Figure 1.** Frontal section of an embryo at stage 13. Cresyl violet,  $\times 50$ ; a — endocardial cushion, b — left ventricle, c — interventricular sulcus, d — interventricular septum (muscular part), e — right ventricle.



**Figure 2.** Horizontal section through the ventricles in an embryo at stage 13. Haematoxylin and eosin,  $\times 100$ ; a — bulbus cordis with ridges, b — left ventricle, c — interventricular septum, d — right ventricle.



**Figure 3.** Horizontal section of an embryo at stage 13. Toluidine blue,  $\times 50$ ; a — dorsal endocardial cushion, b — ventral endocardial cushion, c — bulbus cordis.




**Figure 4.** Oblique horizontal section of the heart in an embryo at stage 14. Cresyl violet,  $\times 50$ ; a — endocardial cushion, b — interventricular flange, c — muscular interventricular septum; d — trabeculae in the right ventricle.

conotruncal ridges were fused and the outflow tract divided, forming two blood streams (Fig. 4, 5).

In addition, during stages 14 and 15 the interventricular septum was elevated and the expansion of the left ventricle and the growth of the left atrium

were observed. The primary interventricular foramen was shifted to the left in the direction of the aortic outflow.

In embryos at stages 16 and 17 the interventricular flange was oriented to the left of the interven-



**Figure 5.** Sagittal section of an embryo at stage 14. Haematoxylin and eosin,  $\times 50$ ; a — fused conotruncal ridges, b — right atrioventricular canal.

tricular septum (Fig. 6) and the atrioventricular cushions were to the right of this flange. The auricles were enlarged and the secondary interventricular foramen was between the elevated muscular interventricular septum and the endocardial cushions.

There was also a counter-clockwise shift of the right and left ventricular outflow tracts from the horizontal to the sagittal axis.

In embryos at stage 18 the membranous interventricular septum developed from the cushion material and, together with the muscular septum, completed the separation of the right and left ventricles (Fig. 7).

In embryos at stage 19 the site of fusion of the membranous and muscular interventricular septum was an area of more cellular structure. Additionally, the aortic and pulmonary blood streams were completely divided (Fig. 8).

## DISCUSSION

Trabeculation is evident in the early phase of the development of the heart. It first becomes evident along the inner myocardial layers near the maximum or greater curvature of the looped primitive ventricle [7, 8, 19]. The pattern of primitive trabecular ridges runs dorsoventrally (circumferential to the heart tube) and appears similar in different species [28]. Forces produced by the peristaltic contractile pattern stimulate patterned trabecular morphogenesis [6]. The primary mechanical consequences of ventricular trabeculation are more uniform transmural stress distribution and increased intramyocardial blood flow. Trabeculations are also important in co-ordinating intraventricular conduction. Patterns of trabeculation specific for the morphologically left, as opposed to the right, ventricles become apparent at the beginning of ventricular septation, and the



**Figure 6.** Oblique frontal section of an embryo at stage 16. Haematoxylin and eosin,  $\times 50$ ; a — interventricular flange, b — endocardial cushion, c — left ventricle, d — right atrium.



**Figure 7.** Sagittal section of an embryo at stage 18. Haematoxylin and eosin,  $\times 50$ ; a — membranous interventricular septum, b — right atrioventricular canal.



**Figure 8.** Oblique horizontal section of an embryo at stage 19. Haematoxylin and eosin,  $\times 50$ ; a — complete interventricular septum, b — left ventricle.

differences are more pronounced in birds than in mammals. The trabeculations in the left ventricle are generally thicker than those in the right ventricle [28, 33, 34].

In the present study the ventricular trabeculation was found in all embryos at stage 13 (the beginning of the 5<sup>th</sup> week). This is in accordance with descriptions of O’Rahilly and Müller [26]. It has to be pointed out that the trabeculations in the left ventricle are thicker and the centripetal growth of the trabeculations is seen.

The muscular part of the interventricular septum in human embryos begins to develop early in the 5<sup>th</sup> week (stages 11 and 12) [26]. In our study this part of the interventricular septum was well marked in embryos at stage 13. Genes *Hand1* and *Hand2* play an essential role in the formation of the interventricular septum and trabeculation [30].

Two opposed hypotheses have been proposed for the development of the interventricular septum [14]. Several authors [26, 27, 31] suggest that the muscular part of the interventricular septum is formed passively by apposition of the expanding ventricular walls. More recent studies have shown, however, that there is a multi-step model of interventricular septum development [14]. An initial passive growth phase delineates the apical/medial aspect of the interventricular septum, including a contribution by cardiomyocytes derived from the inner curvature of the heart. The second step is active in-growth of the left and right derived cardiomyocytes. The third wave of septal growth involves the dorsoventral proliferation of cells within the septum.

The membranous part of the interventricular septum results from fusion of the subendocardial tissue from the right and left bulbar ridges and the dorsal endocardial cushion. The contribution of all

sources to the membranous septum was demonstrated in the present study. It was also shown that the interventricular foramen is obliterated at the end of the 6<sup>th</sup> week (stage 18).

## REFERENCES

- Allwork SP, Anderson RH (1979) Developmental anatomy of the membranous part of the ventricular septum in the human heart. *Br Heart J*, 41: 275–280.
- Anderson RH, Wilkinson JL, Arnold R, Lubkiewicz K (1974) Morphogenesis of bulboventricular malformations: (1) consideration of embryogenesis in the normal heart. *Br Heart J*, 36: 242–255.
- Anderson RH, Webb S, Brown NA, Lamers W, Moorman A (2003) Development of the heart: (2) septation of the atriums and ventricles. *Heart*, 89: 949–958.
- Anderson RH, Webb S, Brown NA, Lamers W, Moorman A (2003) Development of the heart: (3) formation of the ventricular outflow tracts, arterial valves, and intrapericardial arterial trunks. *Heart*, 89: 1110–1118.
- Bartelings MM, Wenink ACG, Gittenberger-de Groot AC, Oppenheimer-Dekker A (1986) Contribution of the aortopulmonary septum to the muscular outlet septum in the human heart. *Acta Morphol Neerl-Scand*, 24: 181–192.
- Bartman T, Hove J (2005) Mechanics and function in heart morphogenesis. *Dev Dyn*, 233: 373–381.
- Ben-Shachar OF, Arcilla RA, Lucas RV, Manasek FJ (1985) Ventricular trabeculations in the chick embryo heart and their contribution to ventricular and muscular septal development. *Circ Res*, 57: 759–766.
- Chalice CE, Viragh S (1973) The architectural development of the early mammalian heart. *Tissue Cell*, 6: 447–462.
- Conte G, Grieco M (1984) Closure of the interventricular foramen and morphogenesis of the membranous septum and ventricular septal defects in the human heart. *Anat Anz*, 155: 39–55.
- Conway SJ, Godt RE, Hatcher CJ, Leatherbury L, Zolotouchnikov W, Brotto MA, Copp AJ, Kirby ML, Creazzo TL (1997) Neural crest is involved in development of abnormal myocardial function. *J Mol Cell Cardiol*, 29: 2675–2685.
- De Jong F, Ophthof T, Wilde AA, Janse MJ, Charles R, Lamers WH, Moorman AF (1992) Persisting zones of slow impulse conduction in developing chicken hearts. *Circ Res*, 71: 240–250.
- De Le Cruz M, Castillo MW, Villavicencio LG, Valencia A, Moreno-Rodriguez RA (1997) Primitive interventricular septum, its primordium, and its contribution in the definitive interventricular septum: in vivo labelling study in the chick embryo heart. *Anat Rec*, 247: 512–20.
- Eisenberg LM, Markwald RR (1995) Molecular regulation of atrioventricular valvuloseptal morphogenesis. *Circ Res*, 77: 1–6.
- Franco D, Meilhac SM, Christoffels VM, Kispert A, Buckingham M, Kelly RG (2006) Left and right ventricular contributions to the formation of the interventricular septum in the mouse heart. *Dev Biol*, 294: 366–375.
- Goodwin RL, Nesbitt T, Price RL, Wells JC, Yost MJ, Potts JD (2005) Three-dimensional model system of valvulogenesis. *Dev Dyn*, 233: 122–129.
- Goore DA, Jesse EE, Lillehei CW (1970) The development of the interventricular septum of the human heart. *Correlative Morphogenetic Study*, Chest, 58: 453–467.
- Hartman T, Hove J (2005) Mechanics and function in heart morphogenesis. *Dev Dyn*, 233: 373–381.
- Hogers B, De Ruyter MC, Baasten AM, Gittenberger-de Groot AC, Poelmann RE (1995) Intracardiac blood flow patterns related to the yolk sac circulation of the chick embryo. *Circ Res*, 76: 871–877.
- Icardo JM (1996) Developmental biology of the vertebrate heart. *J Exp Zool*, 275: 144–161.
- Li Y-X, Zdanowicz M, Young L, Kumiski D, Leatherbury L, Kirby L (2003) Cardiac neural crest in zebrafish embryos contributes to myocardial cell lineage and early heart function. *Dev Dyn*, 226: 540–550.
- Lodewyk HS, Mierop SV, Kutsche LM (1985) Development of the ventricular septum of the heart. *Heart Vess*, 1: 114–119.
- Mcbride RE, Moore GW, Hutchins GM (1981) Development of the outflow tract and closure of the interventricular septum in the normal human heart. *Am J Anat*, 160: 309–331.
- Meredith MA, Hutchins GM, Moore GW (1979) Role of the left interventricular sulcus in formation of the interventricular septum and crista supraventricularis in normal human cardiogenesis. *Anat Rec*, 194: 417–28.
- Molin DGM, Bartram U, Vander Heiden K, Van Iperen L, Speer CP, Hierck BP, Poelmann RE, Gittenberger-de Groot AC (2003) Expression patterns of Tgf $\beta$ 1–3 associate with myocardialisation of the outflow tract and the development of the epicardium and the fibrous heart skeleton. *Dev Dyn*, 227: 431–444.
- Odgers PNB (1938) The development of the pars membranacea septi in the human heart. *J Anat*, 72: 247–259.
- O’Rahilly R, Müller F (2001) *Human Embryology and Teratology*. 3<sup>rd</sup> ed. Wiley-Liss, New York, Chichester, Weinheim, Brisbane, Singapore, Toronto.
- Rychter Z, Rychterova V, Lemez I (1979) Formation of the heart loop and proliferation structure of its wall as a base for ventricular septation. *Herz*, 4: 86–90.
- Sedmera D, Prexieder T, Vuillemin M, Thompson RP, Anderson RH (2000) Developmental patterning of the myocardium. *Anat Rec*, 258: 319–337.
- Szostakiewicz-Sawicka H, Panek-Mikuła J, Prejzner-Morawska A (1978) The membranous part of the interventricular septum of the heart in primates. *Folia Morphol*, 37: 225–235.
- Togi K, Yoshida Y, Matsumae H, Nakashima Y, Kita T, Tanaka M (2006) Essential role of Hand2 in interventricular septum formation and trabeculation during cardiac development. *Bioch Bioph Res Comm*, 343: 144–151.
- Van Microp LHS, Kutsche LM (1985) Development of the interventricular septum of the heart. *Heart Vess*, 1: 114–119.
- Waldo KL, Miyagawa-Tomita S, Kumiski D, Kirby ML (1998) Cardiac neural crest cells provide new insight into septation of the outflow tract: aortic sac to ventricular septal closure. *Dev Biol*, 196: 129–144.
- Wenink ACG (1981) Embryology of the ventricular septum. Separate origin of its components. *Virchows Arch*, 390: 71–79.
- Wenink ACG (1992) Quantitative morphology of the embryonic heart: an approach to development of the atrioventricular valves. *Anat Rec*, 234: 129–135.