

# Orbital size measurement based on computed tomography imaging for surgical safety

A. Skorek<sup>1</sup>, E. Szurowska<sup>2</sup>, D. Raczynska<sup>3</sup>, T. Szmuda<sup>4</sup>, D. Stodulski<sup>1</sup>

<sup>1</sup>Department of Otolaryngology, Medical University of Gdansk, Poland

<sup>2</sup>Department of Radiology, Medical University of Gdansk, Poland

<sup>3</sup>Department of Ophthalmology, Medical University of Gdansk, Poland

<sup>4</sup>Department of Neurosurgery, Medical University of Gdansk, Poland

[Received 14 August 2013; Accepted 10 March 2014]

*Determining the orbital size makes it possible to manoeuvre safely within the orbit during a surgical procedure. Based on the measurements performed on a multi-layer head computed tomography images, the length was determined of the medial, superior, inferior and lateral orbital walls. Also angles were determined between the superior and inferior walls, between the medial and lateral walls, between the inferior wall and Frankfurt plane and between the anterior and posterior segments of the orbital wall. With these measurements it was possible to establish that the safe space for surgical exploration of the orbit (that is the space between the orbital margin and optic canal) is approximately 40 mm. Moreover, it was determined that the medial wall is parallel to the vertical axis of the body and that the angle between the inferior wall and the Frankfurt plane is 19.7°. The angle between the posterior segment of the inferior wall (posterior to the inferior orbital fissure) and the anterior segment is 130.8°. These data will significantly increase the safety of orbital surgeries. (Folia Morphol 2014; 73, 3: 314–320)*

**Key words:** orbit, surgical exploration, computed tomography imaging

## INTRODUCTION

The orbit is a complex, conical structure formed by bones, with the base opening towards the front and the apex towards the back of the head. The orbit contains, above all, the eyeball, the optic nerve and the eyeball muscles. All these structures are tightly bound by ligaments, supporting elements made of connective tissue, and by adipose tissue. The orbit may also be the site of primary or secondary (usually originating from the adjacent paranasal sinuses) inflammatory or neoplastic processes. Another possible issue is evaluating the extent to which the bone elements have been preserved after a trauma and the presence foreign bodies in the orbit. When choosing the appropriate method of surgical treatment, both

the location of the pathological process and the important anatomical structures should be taken into consideration. Such treatment is often performed by teams, such as the following: ophthalmologist, otolaryngologist, and neurosurgeon. It also requires precise knowledge of the orbital anatomy and its possible variability. The key term in the orbital surgery is the “safe space”, namely the distance from the outer bone margin to the optic canal.

The purpose of this paper is to determine the orbital dimensions, angles between the orbital walls and distances between the outer margin of the orbit and its apex (the opening of the optic canal) basing on the images obtained in computed tomography (CT) scans performed in three planes: horizontal, sagittal and frontal.

## MATERIALS AND METHODS


Evaluated were CT images of the head (in the orbital area) of patients with no previous trauma in this area, chronic diseases of the bones and sinuses or surgeries of the neurocranium and splanchnocranium. The study subjects were patients who appeared at the Clinical Emergency Department of the Medical University of Gdansk (head: Prof. A. Basiński, MD PhD) for neurological reasons (headaches, dizziness, status post an epileptic seizure, suspected circulatory disorders within the central nervous system). The group included 100 patients (50 women and 50 men) aged between 18 and 93 years (60.3 years in average). All patients agreed to participate in the study. The CT scans were performed in 3 planes: horizontal, sagittal and frontal, in 0.6 mm layers. All the obtained values were provided in millimetres and entered into the Microsoft Office Excel 2007 spread sheet. Each examination started with adjusting the head position, so that the distance between the nasal ridge (the nasion) and the outermost point of the orbit was equal on both sides (with a margin of 0.5 mm, namely less than 0.6% difference between the sides). If the difference was more than 0.5 mm (due to difficulties in adjusting the position of the head), the results were discarded.

The distance was measured between the lateral margin of the orbit (on the level of the lateral rectus muscle attachment to the orbital globe) to the lateral portion of the optic canal border, and from the verge of the posterior lacrimal crest to the medial portion of the optic canal border (Fig. 1). If the difference in distances between the sides (in 1 patient) was 0.5 mm or more (which was 1.13% and 1.15% of the results obtained on both sides, respectively), the results were discarded and no further measurements were made.


CT images were positioned in the sagittal plane to measure the orbital depth, namely the distance from the superior margin to the apex of the orbit (upper border of the optic canal) and from the inferior margin to the orbital apex (lower border of the optic canal) (Fig. 2).

Basing on the CT images also the angles were evaluated between the lateral and medial walls and superior and inferior walls of the orbit. The evaluation was made on the horizontal and sagittal planes respectively, with the middle of the optic canal as the vertex of the angle and the rays of the angle resting on the bony margins of the orbit (Fig. 3).

Moreover, the angle in the inferior wall between the line anterior to the inferior orbital fissure and posterior to it was evaluated in the sagittal plane.



**Figure 1.** Computed tomography image in the horizontal plane. Measuring the lateral and medial distances.




**Figure 2.** Computed tomography image in the sagittal plane. Measuring the superior and inferior distances.


The vertex of this angle was the bone margins of the inferior orbital fissure. The angle was evaluated between the Frankfurt plane (the plane passing through the superior margin of the external auditory meatus — porion and the inferior margin of the orbit — orbitale) and the inferior orbital wall (Fig. 4).

### Statistical analysis

Descriptive statistics, including mean value, standard deviation (SD), and range were included. The statistical testing was based on the distribution of continuous variables that was verified by the Kolmogorov-Smirnov test. Student's t-test or non-parametric



**Figure 3.** Method of measuring the angles between the medial and lateral walls (A) as well as inferior and superior (B) walls.



**Figure 4.** The angle in the inferior wall (A) and the angle between the Frankfurt plane (red line) and the inferior margin of the orbit (B).

tric Mann-Whitney U test was applied for relevant comparison. Software Statistica v.10.0 (StatSoft, Inc.) was used for statistical analysis, with the significance level less than 0.05 in calculations.

## RESULTS

### Measurements of the orbital size

The average length between the lateral margin and the optic canal was 44.43 mm — men and 43.23 mm — women on the right and respectively 44.10 mm and 43.38 mm on the left side (SD 3.28–3.47 mm). No significant side to side and gender differences were found. The average length was 43.78 mm.

The length between the medial margin and the optic canal was 43.71 mm — men and 42.13 mm — women on the right and 43.56 mm — men

and 42.68 mm — women on the left side (SD 3.34–3.43 mm). No statistically different values were observed between sides and gender. The average length was 43.02 mm.

The average length between the superior margin and the optic canal was 45.75 mm — men and 44.22 mm — women on the right and 45.79 mm — men and 44.67 — women on the left side (SD 3.56–4.33 mm). No significant side to side and gender differences were found. The average length was 45.15 mm.

The length between the inferior margin and the optic canal was 43.73 mm — men and 41.94 mm — women on the right and 43.95 mm — men and 42.19 mm — women on the left side (SD 2.85–3.13 mm). No statistically significant side to side and gender differences were observed. The average length was 42.96 mm. The results are shown in the Table 1.



**Table 1.** Orbital sizes [mm]

Site of measurement	Right side		Left side		Average value
	Men	Women	Men	Women	
Lateral margin — optic canal	44.43	43.23	44.10	43.38	43.78
		*p = 0.52, **p = 0.94			
Medial margin — optic canal	43.71	42.13	43.56	42.68	43.02
		*p = 0.82, **p = 0.35			
Superior margin — optic canal	45.75	44.22	45.97	44.67	45.15
		*p = 0.69, **p = 0.83			
Inferior margin — optic canal	43.75	41.94	43.95	42.19	42.96
		*p = 0.71, **p = 0.34			

\*p — right side vs. left side; \*\*p — men vs. women

**Table 2.** Values of the orbital angles [°]

Site of the angle	Value of the angle				Average value
	Right side		Left side		
	Men	Woman	Men	Woman	
The angle between the medial and lateral walls	50.2	49.6	50.4	49.7	50.0
		*p = 0.65, **p < 0.01			
The angle between the inferior and superior walls	44.7	43.6	44.0	43.2	43.9
		*p = 0.87, **p = 0.51			
The angle in the inferior wall	130.5	130.0	132.9	129.6	130.8
		*p = 0.73, **p = 0.40			
The angle between the Frankfurt plane and the inferior margin	19.9	19.1	20.3	19.3	19.7
		*p = 0.74, **p = 0.11			

\*p — right side vs. left side; \*\*p — men vs. women

### Orbital angles

The average angle between the medial and lateral walls was 50.2° — men and 49.6° — women on the right and 50.4° — men and 49.7° — women on the left side. No statistically significant side to side differences were found in measurement results. The difference between the angle size in men and women was statistically significant. The average angle was 50.0° (SD 3.82–3.95°).

The average angle between the inferior and superior walls was 44.7° — men and 43.6° — women on the right and 44.0° — men and 43.2° — women on the left side. No statistically significant side to side and gender differences were found. The average angle was 43.9° (SD 5.45–5.23°) (Fig. 3).


The average angle in the inferior wall was 130.5° — men and 130.0° — women on the right and 132.9° — men and 129.6° — women on the left side (SD

8.4–8.9°). The average angle was 130.8°. The average angle between the Frankfurt plane and the inferior margin of the orbit was 19.9° — men and 19.1° — women on the right and 20.3° — men and 19.3° — women on the left side (SD 4.6–5.6°). The average angle was 19.7°. No statistically significant side to side and gender differences were found in all measurement results (Fig. 4, Table 2).

As a portrayal of our analysis, a patient with an orbital tumour was presented herein. A meticulous preoperative knowledge of orbital measurements and angles is necessary in the contemporary microsurgery of these lesions (Fig. 5).

### DISCUSSION

In order to determine orbital size basing on the CT imaging, fixed, repetitive and easily recognisable points need to be established, on which the measu-



**Figure 5.** The validation of the orbit's measurements for clinical practice. Medial, superior and inferior walls of the orbit were marked consecutively by blue arrows (A. Axial; B, C. Sagittal preoperative T1 magnetic resonance imaging plane). Patient's head and small schwannoma of the optic nerve were reconstructed for neurosurgical planning purpose (D–F). The tumour was located in the optic canal (A–F red arrows). Intraoperative screenshots (G–I) present the advanced visualisation methods. The 'injected pictures' in the operating microscope vision depict the outlines of the appropriate osteotomy (G, white arrows), optic nerve and chiasm (H, yellow arrows) as well as the lateral wall of the orbit (blue arrows). The navigation (G–I right upper corner) enables the constant monitoring of the removed anatomical structures of the orbit. Screenshots from a navigation station (J–O) show current intraoperative visualisation capabilities with respect to the lateral wall of the orbit (K, M, green dashed line and blue arrows) or outlined optic nerve (L, M, O, yellow arrows).

rements will depend. Such points need not be identical with the landmarks, which are used for better intraoperative and topodiagnostic orientation. The points used for measurements, however, are used to represent the size of the orbit (distance from the base to the apex, determining the angles between the walls or particular elements of the walls, etc.) [24].

The starting point for measurements in the medial wall was, in this study, the verge of the posterior lacrimal crest, and the final point was the medial optic canal border. Similar final points, to which the distance from the starting point was measured, were provided by Ji et al. [12], who used the frontomaxillary suture as the starting point (easy to find on a dry skull and intraoperatively, but difficult to establish in CT imaging). Other authors provided as a measurement point (e.g. to determine the distance to the ethmoidal foramina) the anterior or posterior lacrimal crest [5, 6, 11, 14]. The average result of the present measurements was 43.02 mm (no statistically significant side to side and gender differences), which is similar to the results obtained by other authors (43.77 mm and 42.0 mm) [1, 23].

Within the lateral wall, the distance was measured between the bony point on the level of the lateral rectus muscle attachment to the orbital globe and the lateral portion of the optic canal border. Many authors recommend the frontozygomatic suture as the starting point of measurement. This

point is a surgical landmark, but is difficult to locate in CT images [6, 10, 14, 17, 23]. The starting point in this study is analogous to the point proposed by Ji et al. [12] — it is an external bony point in the intersection between the horizontal line that divides the orbit in two equal parts (upper and lower) and the lateral bony margin. This point is equivalent to the Whitnall's tubercle — approximately 1 cm inferior to the frontozygomatic suture and approximately 4–5 mm behind the lateral orbital rim, where attached are the lateral canthal tendon, the lateral rectus check ligament, lateral horn of the levator aponeurosis, suspensory ligament of the lower lid (Lockwood's ligament), the lacrimal gland fascia and the orbital septum [5, 16]. Lee et al. [15] used for their measurements such horizontal CT images, in which the length of the lateral rectus muscle was largest. In the present study this length was 43.78 mm in average (no statistically significant side to side and gender differences). This result was within the range obtained in the measurements of the distance from the frontozygomatic suture to the optic canal: 43.0 mm ÷ 47.1 mm [17, 22].

The starting point in the measurements within the superior wall was usually the supraorbital foramen [12, 14, 17, 23]. The distances to this foramen from the optic canal were within the range from 40 mm to 48.65 mm. In the measurements performed in the present study, the starting point was the superior

orbital margin in the sagittal plane and the final point was the upper border of the optic canal. The result was 45.15 mm (no statistically significant side to side and gender differences). In other studies, the starting point for measurements in the lower wall was the infraorbital foramen or the bony margin of the orbit above the infraorbital foramen. The results were  $40 \div 54$  mm, 50.3 mm in average, and 45.5 mm, respectively [11, 14, 17, 21, 24]. In the present study, the bony margin of the orbit was used as the starting point and the distance to the optic canal was 42.96 mm in average (no statistically significant side to side and gender differences). Both these distances are within the standard error, and the differences between the results may be caused by differences in the applied measurement methods and the subjects who underwent the measurement.

For surgical exploration of the orbit, the distances between the bony margin to the unsafe zone (superior and inferior orbital fissure, ethmoidal vessels) and the critical zone (optic canal) need to be determined [3, 13, 23]. Based on the analysis of the obtained results it was established that the safe space (distance between the bony margin of the orbit and the optic canal) in each of the orbital wall is approximately 40 mm. Other authors obtained similar results [20, 22].

The size of the orbit was evaluated based on the angles between the lateral and medial walls and superior and inferior walls of the orbit. For the evaluation of the inferior wall, the angle was used between this wall and the Frankfurt plane — this angle in the study group was  $19.7^\circ$ , and between the anterior and posterior segment of the orbital floor —  $130.8^\circ$  (no statistically significant side to side and gender differences). According to Nagasao et al. [19], the angle between the orbital floor and the Frankfurt plane is larger and amounts to  $30^\circ$ . That study was performed on Japanese subjects [19]. The angle between the medial and lateral wall of the orbit in the present study was  $50.0^\circ$  (no statistically significant side to side differences were observed, but the angle size was statistically significant greater in male gender), while other authors obtained a smaller angle of  $45^\circ$  [22]. However, Dortzbach [7] emphasised that there is a large variety of angles between the orbital walls, depending on gender and ethnicity. The angles between the orbital walls (orbital geometry) determine the orbital volume [22]. Al-Sukhun et al. [4] stated that the larger the orbital volume, the less resistant the orbit is to a trauma — a 1 mm difference in the wall spacing

reduces the resistance by approximately 50%. That study was performed on a statistical computer model.

Laryngological surgical approaches to the orbit lead through the medial wall (transethmoidal approach) or inferior wall (transantral approach) [3, 9, 18]. The angles between these walls require different preparation methods. Analysing the obtained study results it should be noted that the medial wall is practically parallel to the vertical axis of the body. On the other hand, in the analysis of the angle between the inferior wall and the Frankfurt plane, the inferior wall was determined to be sloping “upwards”. In the examined study group this angle was  $19.7^\circ$  in average. The angle between the posterior segment of the inferior wall (posterior to the inferior orbital fissure) and the anterior segment is  $130.8^\circ$ . This needs to be taken into account during surgical procedures in this area. This position of the inferior wall also affects the resistance to trauma of the inferior margin [2, 8].

## CONCLUSIONS

The measurements performed on head CT images made it possible to determine the distance from the outer margin of the orbit to the optic canal, which amounts to 42.96 mm for the inferior wall, 45.15 mm for the superior wall, 43.02 mm for the medial wall and 43.78 mm for the lateral wall. Therefore, the “safe distance” for all orbital walls is approximately 40 mm. Angle between the medial and lateral walls was  $50^\circ$  and between superior and inferior walls was  $43.9^\circ$ . Average angle in inferior wall was  $130.8^\circ$  and between Frankfurt plane and inferior margin was  $19.7^\circ$ .

## REFERENCES

1. Abed SF, Sham P, Shen S, Adds PJ, Uddin JM (2012) A cadaveric study of ethmoidal foramina variation and its surgical significance in Caucasians. *Br J Ophthalmol*, 96: 118–121.
2. Ahmad F, Kirkpatrick WNA, Lyne J, Urdang M, Garey LJ, Waterhouse N (2003) Strain gauge biomechanical evaluation of forces in orbital floor fractures. *Br J Plast Surg*, 56: 3–9.
3. Akdemir G, Tekdemir I, Altm L (2004) Transethmoidal approach to the optic canal: surgical and radiological microanatomy. *Surg Neurol*, 62: 268–174.
4. Al-Sukhun J, Penttilä H, Ashammakhi N (2012) Orbital stress analysis. Part V: systematic approach to validate a finite element model of human orbit. *J Craniofac Surg*, 23: 669–674.
5. Burkat CN, Lemke BN (2005) Anatomy of the orbit and its related structures. *Otolaryngol Clin N Am*, 38: 825–856.
6. Chastain JB, Sindwani R (2006): Anatomy of the orbit, lacrimal apparatus and lateral nasal all. *Otolaryngol Clin N Am*, 39: 855–864.

7. Dortzbach RK ed. (1994) *Ophthalmic plastic surgery: prevention and management of complications*. Raven Press, New York.
8. Fujino T (1974) Experimental "blow-out" fracture of the orbit. *Plast Reconstr Surg*, 54: 81–82.
9. Goldberg RA, Kim AJ, Kerivan KM (1998) The lacrimal keyhole, orbital door jamb and basin of the interior orbital fissure. Three areas of deep bone in the lateral orbit. *Arch Ophthalmol*, 116: 1618–1624.
10. Gönül E, Erdogan E, Düz B, Timurkaynak E (2003) Transmaxillary approach to the orbit: an anatomic study. *Neurosurgery*, 53: 935–941.
11. Hwang K, Baik SH (1999) Surgical anatomy of the orbit of Korean adults. *J Craniofac Surg*, 10: 129–133.
12. Ji Y, Qian Z, Dong Y, Zhou H, Fan X (2010) Quantitative morphometry of the orbit in Chinese adults based on a three-dimensional reconstruction method. *J Anat*, 217: 501–508.
13. Jones DEP, Evans JNG (1967) "Blow-out" fractures of the orbit: an investigation into their anatomical basis. *J Laryngol Otol*, 81: 1109–1120.
14. Karakas P, Bozkir MG, Oguz Ö (2002) Morphometric measurements from various reference points in the orbit of male Caucasians. *Surg Radiol Anat*, 24: 358–362.
15. Lee H, Lee Y, Ha S, Park M, Baek S (2011) Measurement of width and distance of the posterior border of the deep lateral orbital wall using computer tomography. *J Craniomaxillofac Surg*, 39: 606–609.
16. Liesegang TJ, Skuta GL, Cantor LB eds. (2005) *Basic and clinical science course. Część 7. [Oczodół, powieki i układ łzowy]*. Wydawnictwo Medyczne Urban & Partner, Wrocław.
17. McQueen CT, Di Ruggiero DC, Campbell JP, Shockley WW (1995) Orbital osteology: a study of the surgical landmarks. *Laryngoscope*, 105: 783–788.
18. Michel O (2000) Transnasale Chirurgie der Orbita. *HNO*, 48: 4–17.
19. Nagasao T, Miyamoto J, Nagasao M, Ogata H, Kaneko T, Tamaki T, Nikajima T (2006) The effect of striking angle on the buckling mechanism in blowout fracture. *Plast Reconstr Surg*, 117: 2373–2380.
20. Nitek S (2005) *Morfologia i morfometria struktur kostnych oczodołu w aspekcie bezpieczeństwa wybranych dośń operacyjnych*. Rozprawa doktorska. Akademia Medyczna w Warszawie.
21. Regensburg NI, Wiersinga WM, van Velthoven ME, Berendschot TT, Zonneveld FW, Baldeschi L, Saeed P, Mourits MP (2011) Age and gender-specific reference values of orbital fat and muscle volumes in Caucasians. *Br J Ophthalmol*, 95: 1660–1663.
22. Rene C (2006) Update on orbital anatomy. *Eye*, 20: 1119–1129.
23. Rontal E, Rontal M, Guilford FT (1979) Surgical anatomy of the orbit. *Ann Otol Rhinol Laryngol*, 88: 382–386.
24. Wanyura H (1996) Anatomia chirurgiczna oczodołu. *Czas Stomat*, 49: 567–578.