


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/pjnns>

Review article

Treatment of lumbar disc herniation with radiculopathy. Clinical practice guidelines endorsed by The Polish Society of Spinal Surgery

Dariusz Latka^{a,*}, Grzegorz Miekisiak^b, Pawel Jarmuzek^c,
Marcin Lachowski^d, Jacek Kaczmarczyk^e

^aDepartment of Neurosurgery, Regional Medical Center, Opole, Poland

^bDepartment of Neurosurgery, Specialist Medical Center, Polanica-Zdroj, Poland

^cDepartment of Neurosurgery, Regional Neurosurgery and Neurotrauma Center, Zielona Gora, Poland

^dMedical University of Warsaw, Warsaw, Poland

^eDepartment of Orthopedics and Traumatology, Poznan University of Medical Sciences, Poznan, Poland

ARTICLE INFO

Article history:

Received 19 November 2015

Accepted 2 December 2015

Available online 17 December 2015

Keywords:

Spinal surgery

Herniated lumbar disc

Radiculopathy

Guidelines

Degenerative disc disease

ABSTRACT

Introduction: Herniated lumbar disc (HLD) is arguably the most common spinal disorder requiring surgical intervention. Although the term is fairly straightforward, the exact pathology and thus the clinical picture and natural history may vary. Therefore, it is immensely difficult to formulate universal guidelines for surgical treatment.

Aim: The aim of this paper is to organize the terminology and clear the inconsistencies in phraseology, review treatment options and gather available published evidence to address the clinical questions to create a set of clinical guidelines in relevant to the topic.

Methods and results: Twelve queries, addressing optimal surgical treatment of the HLD have been formulated. The results, based on the literature review are described in the present work. The final product of the analysis was a set of guidelines for the surgical treatment of symptomatic HLD. Categorized into four tiers based on the level of evidence (I–III and X), they have been designed to assist in the selection of optimal, effective treatment leading to the successful outcome.

Conclusions: The evidence based medicine (EBM) is becoming ever more popular among spinal surgeons. Unfortunately this is not always feasible. Lack of uniform guidelines and numerous conflicts of interest introduce flaws in the decision making process. The key role of experts and professional societies is to provide high value recommendation based on the most current literature. Present work contains a set of guidelines for the surgical treatment of HLD officially endorsed by the Polish Spine Surgery Society.

© 2015 Polish Neurological Society. Published by Elsevier Sp. z o.o. All rights reserved.

* Corresponding author at: Department of Neurosurgery, Regional Medical Center, al. W. Witosa 26, 45-418 Opole, Poland.

Tel.: +48 601853513.

E-mail address: dlatka@mp.pl (D. Latka).

<http://dx.doi.org/10.1016/j.pjnns.2015.12.001>

0028-3843/© 2015 Polish Neurological Society. Published by Elsevier Sp. z o.o. All rights reserved.

Introduction

Herniated lumbar disc (HLD) is the most common spine disorder requiring surgical intervention. It is a problem in daily practice of spine specialists. Although HLD is quite consistent in clinical presentation, it is due to the variable pathology and natural history that indications for surgical intervention are not precise. The necessity of choosing the right surgical technique makes the situation even more complex. New methods and technologies are constantly being adapted in treatment of spine disorders. Currently, not only do we have means of intraoperative visualization (naked eye, microscope, endoscope), but also various methods of tissue dissection (classic and tubular retractors, percutaneous techniques) and a range of systems for supplementation and internal instrumentation. Variety of therapeutic options imposes making decisions based on trustworthy and up-to-date articles (EBM). Unfortunately, there are still no clear therapeutic guidelines. Spine surgeons are often subjects of external, nonobjective influence, also from the medical industry. In these circumstances, the main role of expert groups and scientific societies is to create recommendations based on up-to-date articles to be used in medical practice. Such recommendations provide help in choosing the optimally effective procedure for a functional cure.

Definition and classification of HLD

In order to create a systematic review of literature on lumbar discopathy patients with radiculopathy the authors used the definition of HLD based on the consensus of North American Spine Society published in 2001 [1]. According to the consensus:

HLD is a localized displacement of the disc material outside the anatomic borders of intervertebral space that causes pain, weakness or numbness in myotomal or dermatomal distribution.

HLD is not a homogenous disorder. Equally important in indications for surgical treatment and choosing the right method and surgical approach is the degree of lumbar disc herniation, being the basis for pathologic classification [2] and the spatial relationship on which the topographic classification of the disease is based [3].

Methods

Twelve queries connected with optimal surgical treatment of HLD were designed. The results, based on the review of literature are presented in this article. The guidelines for a surgical treatment of symptomatic disc herniation with radiculopathy in adult population were created, classified into four grades using the four levels of quality specified by GRADE [4], according to the Cochrane Back Review Groups [5] (Table 1).

Table 1 – Assessment criteria for scientific evidence and grading of recommendations.

Initial quality levels of scientific evidence		
Type of study	Quality level of a scientific study	Grade
RCT	High	4
–	Medium	3
Observational study	Low	2
Different studies	Very low	1
Modification of grade/points		
Score reducing factors:		
Serious (–1) or very serious (–2) limitation to study quality		
Important inconsistency (–1)		
Some (–1) or major (–2) uncertainty about directness		
Data not precise or lacking (–1)		
High probability of selective reporting (–1)		
Score increasing factors:		
Strong evidence of association – significant relative risk >2 ($p < 0.5$) based on consistent results of two or more observational studies, with no plausible confounders (+1)		
Very strong evidence of association – significant relative risk >5 ($p < 0.2$) based on direct evidence, without concerns about credibility with no major threats to validity (+2)		
Evidence of a dose response gradient (+1)		
All plausible confounders have reduced the effect (+1)		
Final recommendation classes in relations to the strength of scientific evidence		
Score (pts)	Recommendation class	
4	I	
3	II	
2	III	
No evidence	X	
Based on Atkins et al. [4] with modifications.		

Results

Natural history of the disease

HLD with radiculopathy is characterized by a favorable natural history. Many articles on correlation of radiological improvement with clinical improvement were identified. In a vast majority of patients there is an improvement after few weeks. No evident radiological correlation with prognosis of spontaneous remission was found [6–21].

Surgical or conservative treatment? Indications for a surgery

Six controlled, randomized or quasi-randomized clinical trials comparing results of surgical treatment (discectomy) with results of conservative treatment were identified [15,22–26], one of which was classified as Class I evidence [15], the others as Class II.

Class II recommendation: in particular patients, short-term result of surgical treatment is better when compared with conservative treatment. Patients, who improved after

surgical treatment, fit the pattern: pain radiating consistently with the level of HLD-nerve conflict, positive stretch test (straight leg raising test positive at 10-45°), convincing diagnostic imaging (level, side) [26]. Reoperated patients were not included in any of cited studies.

Class X recommendation: based on literature analysis, there is no clear evidence favoring surgical treatment over conservative treatment in longer observation (longer than 1 year).

What is the optimal time from the onset of symptoms to the surgical treatment?

The query returned four prospective controlled trials/studies (Class II) [27-30] and 9 retrospective studies of Class III [31,32,33,34,35,36,37,38]. Analysis of the literature allows to formulate the recommendations:

Class II recommendation: longer delay from the onset of symptoms to surgical treatment is associated with worse outcomes.

Class III recommendation: the threshold period after which the treatment outcome is significantly worse is 6 months.

Which patients have the best outcomes when treated surgically?

Authors of three articles of sufficient quality attempted to identify factors predicting favorable outcome of surgical treatment. Two articles concerned the endoscopic techniques [39,40] and the remainder pertained to the open surgery [41].

Class II recommendation: better outcome can be expected in patients younger than 40, with severe pain radiating to the leg, and when the symptoms are present for less than 3 months.

What is the effectiveness of intradiscal techniques?

Percutaneous discectomy

Among the Randomized Controlled Trials (RCT) analyzed on the effectiveness of percutaneous discectomy [42-45], none met the minimal quality criteria, mainly because of the short follow-up outcomes only. A few prospective studies without control and/or without randomization [46-49] were found.

Class III recommendation: percutaneous discectomy is an effective method of treatment.

No advantage of the method over the open surgery was found (Class X recommendation).

Intradiscal electrotherapy (IDET)

Class X recommendation: there is no clinical evidence [50] proving IDET's effectiveness or lack of thereof in radicular pain treatment.

What is the effectiveness of various surgical techniques (microdiscectomy, open discectomy, tubular techniques) used in the treatment of HLD?

Sixteen articles describing RCTs addressing the effectiveness and comparing treatment outcomes of various techniques were identified [51-66].

Class I recommendation: either microdiscectomy, open discectomy or discectomy with the use of tubular technology are effective treatment methods of the HLD with radiculopathy.

Class X recommendation: because of the lack of sufficient scientific evidence of proper quality, no technique can be called superior.

Sequestrectomy or discectomy?

The query returned one RCT [58,67,68] comparing results of the microdiscectomy with sequestrectomy, and 5 observational, mainly retrospective studies [69-74].

Class X recommendation: because of the lack of scientific evidence of sufficient quality, the superiority of sequestrectomy over discectomy cannot be proven.

Should the means of prevention of epidural adhesions be used?

Ten RCTs on safety and effectiveness of techniques and materials limiting the development of epidural adhesions were identified and analyzed [75-83]. Two studies have shown the advantage of using anti-adhesion gel in comparison with control group [77,84], although little power of these studies does not allow to form firm recommendation. One study has shown no relationship between the outcomes and the presence of adhesions on MRI scans [83].

Class X recommendation: there is no scientific evidence supporting the use of barrier methods in prevention of epidural adhesions.

Is the antibiotic prophylaxis effective?

Among the large number of articles pertaining to this topic, only two describe the results of RCTs on the effectiveness of antibiotic prophylaxis in spinal surgery [85,86]. All analyzed studies are of low value because of many confounders [87-92].

Class II recommendation: the antibiotic prophylaxis applied immediately before the spine surgery decreases the risk of infectious complications.

Class X recommendation: there is not enough evidence, that the prophylactic administration of antibiotics after the surgery decreases the risk of infectious complications.

Can lumbar fusion be an appropriate primary method of treatment of the HLD with radiculopathy?

No RCTs were identified, only a few observational studies [93-100], mainly retrospective of low value. Many confounding

factors and the lack of standardization prevent from performing meta-analyses.

Class X recommendation: available clinical evidence does not allow to create recommendations or contraindications for spondylodesis as a primary method of treatment of the HLD with radiculopathy.

Can lumbar fusion be used as the treatment of recurrent HLD?

The result of the query was one article describing results of a RCT with high risk of bias [101] and a few observational studies. In the literature there is no data on observations longer than 3 years [102-104]. An important confounding factor affecting meta-analyses is heterogeneity of clinical manifestation among various groups.

Class III recommendation: Lumbar fusion is an effective method of treatment for recurrent HLD, especially when the clinical presentation is dominated by axial pain and/or signs of instability.

What are the medium- (1-4 years) and long-term outcomes of the surgical treatment of HLD with radiculopathy in comparison with the conservative treatment?

Five RCTs comparing short- and medium-term (up to 2 years) outcomes from treatment were identified [15,22,26,105,106]. The available long-term results come only from an RCT of poor quality [23] or observational studies, either prospective [7] or retrospective [107]. Although numerous articles pointed to advantage of surgical treatment in any time period, none of these studies was of sufficient quality allowing for the formulation of recommendations.

Class X recommendation: based on the available literature, superiority of surgical treatment over conservative treatment in HLD with radiculopathy cannot be proven, neither in medium- nor long-term assessment.

Discussion and conclusions

Our analysis confirms a long and well-known belief among spine surgeons that there still is not enough high quality medical evidence allowing standardization of diagnostic and therapeutic decisions, even in regards to a very common disorder such as HLD. Currently, the main sources of data used to create of guidelines are the results of RCTs [108]. In spine surgery, conduction of such studies poses a great challenge, not only because of the amount of workload and associated great costs. Many confounders must be considered, some of the very difficult to measure. For example the most frequent indication for surgery is pain and poor quality of life, and these factors are notoriously difficult to assess and further studies on significance of psychosocial factors are required [109]. Very often well designed and well performed studies may lead to completely contradicting conclusions making the formulation of clinical recommendations virtually impossible [110]. There

is a certain hope in recent advances in spine surgery registries, such as SpineTango. First analyses comparing results of RCTs with data from registries are promising [111], with similar results but at largely reduced costs.

At the same time, pressure of taxpayers imposes standardization of treatment. There are many attempts to organize the general of treatment of spinal disorder, ranging from establishing the formal nomenclature [1] to detailed indications for performing particular surgical procedures. Most often, they are created by expert groups working under auspices of professional societies, such as North American Spine Society recommendations [112]. Such formal guidelines are required not only for making optimal clinical decisions but also play an important role in socio-economic perspective, shaping payers' policies (NASS Coverage Recommendations) [113]. Formulation of such guidelines should take into account various regional factors, in particular the socioeconomic status.

This publication, influenced to some degree by aforementioned NASS publication [112], is an attempt to create recommendations for the most common surgical spine pathologies encountered in a typical practice. We hope it will undergo further refinement by panels of expert as new evidences become available.

The analysis of literature presented in this paper allowed for the formulation of guidelines shown below. This publication has been endorsed by the Polish Society of Spinal Surgery.

- (1) The herniation of nucleus pulposus causing neurological symptoms and signs of irritation or deficits may be treated surgically.
- (2) Patients treated surgically should be symptomatic, and there should be a clear correlation between radiological imaging and clinical presentation.
- (3) Given the natural history of the disease, surgical treatment is superior over conservative treatment, mainly because of the shorter recovery of symptoms, i.e. the benefit is in short-term.
- (4) The indication for immediate surgical treatment is neurological deficit: cauda equina syndrome and/or reduced muscle strength.
- (5) In cases with painful radiculopathy, surgical treatment should be offered after 6 weeks of ineffective conservative treatment, but not later than 6 months after the onset of symptoms. Patients in remission should not be treated surgically.
- (6) The mainstay of surgical treatment is discectomy without fusion. Effectiveness of discectomy is similar regardless of the method of viewing aids used (endoscopy, microscope, open) or the method of tissue retraction (percutaneous, tubular, classic retractors), therefore the selection of technique can be based on surgeon's preference. In case of recurrent disc herniation, lumbar fusion can be considered, especially in patients with severe axial pain and/or signs of instability.
- (7) The choice between full discectomy and sequestrectomy can also be based on surgeon's preference – there is no evidence for superiority of either method.
- (8) There is no evidence supporting the use of barrier methods in prevention of epidural adhesions when performing the transcanal discectomy.

- (9) The antibiotic prophylaxis applied immediately before the spine surgery decreases the risk of infectious complications.
- (10) There is no evidence for effectiveness of IDET and hence it should not be recommended.
- (11) The percutaneous discectomy is an effective method of treatment. There is no evidence for its superiority over open discectomy.

Conflict of interest

The authors declare no conflict of interest.

Funding

No financial support was received for this study.

Ethics

The work described in this article has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans; Uniform Requirements for manuscripts submitted to Biomedical journals.

REFERENCES

- [1] Fardon DF, Milette PC. Nomenclature and classification of lumbar disc pathology, recommendations of the Combined task Forces of the North American Spine Society, American Society of Spine Radiology, and American Society of Neuroradiology. *Spine (Phila PA 1976)* 2001;26:E93-113.
- [2] Caplan LR. Evaluation of patients with suspected herniated lumbar discs with radiculopathy. *Eur Neurol* 1994;34:53-60.
- [3] Wiltse LL, Berger PE, McCulloch JA. A system for reporting the size and location of lesions in the spine. *Spine (Phila PA 1976)* 1997;22:1534-7.
- [4] Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, et al. Grading quality of evidence and strength of recommendations. *Br Med J* 2004;328:1490. <http://dx.doi.org/10.1136/bmj.328.7454.1490>
- [5] Furlan AD, Pennick V, Bombardier C, van Tulder M. 2009 updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine (Phila PA 1976)* 2009;34:1929-41.
- [6] Atlas SJ, Deyo RA, Keller RB, Chapin AM, Patrick DL, Long JM, et al. The Maine Lumbar Spine study, Part II: 1-year outcomes of surgical and nonsurgical management of sciatica. *Spine (Phila PA 1976)* 1996;21:1777-86.
- [7] Atlas SJ, Keller RB, Wu YA, Deyo RA, Singer DE. Long-term outcomes of surgical and nonsurgical management of sciatica secondary to a lumbar disc herniation: 10 year results from the Maine Lumbar Spine study. *Spine (Phila PA 1976)* 2005;30:927-35.
- [8] Baldwin NG. Lumbar disc disease: the natural history. *Neurosurg Focus* 2002;13:1-4.
- [9] Benoist M. The natural history of lumbar disc herniation and radiculopathy. *Jt Bone Spine* 2002;69:155-60.
- [10] Bush K, Cowan N, Katz DE, Gishen P. The natural history of sciatica associated with disc pathology: a prospective study with clinical and independent radiologic follow-up. *Spine (Phila PA 1976)* 1992;17:1205-12.
- [11] Cowan NC, Bush K, Katz DE, Gishen P. The natural history of sciatica: a prospective radiological study. *Clin Radiol* 1992;46:7-12.
- [12] Cribb GL, Jaffray DC, Cassar-Pullicino VN. Observations on the natural history of massive lumbar disc herniation. *J Bone Jt Surgery Br Vol* 2007;89:782-4.
- [13] Komori H, Shinomiya K, Nakai O, Yamaura I, Takeda S, Furuya K. The natural history of herniated nucleus pulposus with radiculopathy. *Spine (Phila PA 1976)* 1996;21:225-9.
- [14] Masui T, Yukawa Y, Nakamura S, Kajino G, Matsubara Y, Kato F, et al. Natural history of patients with lumbar disc herniation observed by magnetic resonance imaging for minimum 7 years. *J Spinal Disord Tech* 2005;18:121-6.
- [15] Peul WC, van Houwelingen HC, van den Hout WB, Brand R, Eekhof JAH, Tans JTJ, et al. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med* 2007;356:2245-56.
- [16] Peul WC, van Houwelingen HC, van der Hout WB, Brand R, Eekhof JAH, Tans JTJ, et al. Prolonged conservative treatment or early surgery in sciatica caused by a lumbar disc herniation: rationale and design of a randomized trial ISRCT 26872154. *BMC Musculoskelet Disord* 2005;6:8.
- [17] Tubach F, Beauté J, Leclerc A. Natural history and prognostic indicators of sciatica. *J Clin Epidemiol* 2004;57:174-9.
- [18] Valat JP, Giraudeau B, Rozenberg S, Goupille P, Bourgeois P, Micheau-Beaugendre V, et al. Epidural corticosteroid injections for sciatica: a randomised, double blind, controlled clinical trial. *Ann Rheum Dis* 2003;62:639-43.
- [19] van den Hout WB, Peul WC, Koes BW, Brand R, Kievit J, Thomeer RTWM. Prolonged conservative care versus early surgery in patients with sciatica from lumbar disc herniation: cost utility analysis alongside a randomised controlled trial. *Br Med J* 2008;336:1351-4.
- [20] Vroomen PCAJ, De Krom M, Knottnerus JA. Predicting the outcome of sciatica at short-term follow-up. *Br J Gen Pract* 2002;52:119-23.
- [21] Weber H, Holme I, Amlie E. The natural course of acute sciatica with nerve root symptoms in a double-blind placebo-controlled trial evaluating the effect of piroxicam. *Spine (Phila PA 1976)* 1993;18:1433-8.
- [22] Buttermann GR. Treatment of lumbar disc herniation: epidural steroid injection compared with discectomy. A prospective, randomized study. *J Bone Jt Surg Am* 2004;86-A:670-9.
- [23] Lurie JD, Tosteson TD, Tosteson ANA, Zhao W, Morgan TS, Abdu WA, et al. Surgical versus nonoperative treatment for lumbar disc herniation: eight-year results for the spine patient outcomes research trial. *Spine (Phila PA 1976)* 2014;39:3-16.
- [24] Österman H, Seitsalo S, Karppinen J, Malmivaara A. Effectiveness of microdiscectomy for lumbar disc herniation: a randomized controlled trial with 2 years of follow-up. *Spine (Phila PA 1976)* 2006;31:2409-14.
- [25] Weber H. Lumbar disc herniation. A controlled, prospective study with ten years of observation. *Spine (Phila PA 1976)* 1983;8:131-40.
- [26] Weinstein JN, Lurie JD, Tosteson TD, Skinner JS, Hanscom B, Tosteson ANA, et al. Surgical vs nonoperative treatment for lumbar disc herniation: the Spine Patient Outcomes Research Trial (SPORT) observational cohort. *JAMA* 2006;296:2451-9.

- [27] Blazhevski B, Filipche V, Cvetanovski V, Simonovska N. Predictive value of the duration of sciatica for lumbar discectomy. *Prilozi* 2008;29:325-35.
- [28] Hurme F M., Alaranta H. Factors predicting the result of surgery for lumbar intervertebral disc herniation. *Spine (Phila PA 1976)* 1987;12:933-8.
- [29] Fisher C, Noonan V, Bishop P, Boyd M, Fairholm D, Wing P, et al. Outcome evaluation of the operative management of lumbar disc herniation causing sciatica. *J Neurosurg* 2004;100:317-24.
- [30] Nygaard ØP, Kloster R, Solberg T. Duration of leg pain as a predictor of outcome after surgery for lumbar disc herniation: a prospective cohort study with 1-year follow up. *J Neurosurg Spine* 2000;92:131-4.
- [31] Akagi R, Aoki Y, Ikeda Y, Nakajima F, Ohtori S, Takahashi K, et al. Comparison of early and late surgical intervention for lumbar disc herniation: is earlier better? *J Orthop Sci* 2010;15:294-8.
- [32] Gaetani P, Aimar E, Panella L, Debernardi A, Tancioni F, Baena RR. Surgery for herniated lumbar disc disease: factors influencing outcome measures. An analysis of 403 cases. *Funct Neurol* 2004;19:43-50.
- [33] Ng LCL, Sell P. Predictive value of the duration of sciatica for lumbar discectomy. A prospective cohort study. *J Bone Jt Surgery Br Vol* 2004;86:546-9.
- [34] Nygaard ØP, Romner B, Trumpy JH. Duration of symptoms as a predictor of outcome after lumbar disc surgery. *Acta Neurochir (Wien)* 1994;128:53-6.
- [35] Quon JA, Sobolev BG, Levy AR, Fisher CG, Bishop PB, Kopec JA, et al. The effect of waiting time on pain intensity after elective surgical lumbar discectomy. *Spine J* 2013;13:1736-48.
- [36] Rihn JA, Hilibrand AS, Radcliff K, Kurd M, Lurie J, Blood E, et al. Duration of symptoms resulting from lumbar disc herniation: effect on treatment outcomes analysis of the Spine Patient Outcomes Research Trial (SPORT). *J Bone Jt Surg* 2011;93:1906-14.
- [37] Schoeggl A, Maier H, Saringer W, Reddy M, Matula C. Outcome after chronic sciatica as the only reason for lumbar microdiscectomy. *J Spinal Disord Tech* 2002;15:415-9.
- [38] Suzuki A, Matsumura A, Konishi S, Terai H, Tsujio T, Dozono S, et al. Risk factor analysis for motor deficit and delayed recovery associated with L4/5 lumbar disc herniation. *J Spinal Disord Tech* 2011;24:1-5.
- [39] Ahn Y, Lee S-H, Park W-M, Lee H-Y, Shin S-W, Kang H-Y. Percutaneous endoscopic lumbar discectomy for recurrent disc herniation: surgical technique, outcome, and prognostic factors of 43 consecutive cases. *Spine (Phila PA 1976)* 2004;29:E326-32.
- [40] Ahn Y, Lee S-H, Lee JH, Kim JU, Liu WC. Transforaminal percutaneous endoscopic lumbar discectomy for upper lumbar disc herniation: clinical outcome, prognostic factors, and technical consideration. *Acta Neurochir (Wien)* 2009;151:199-206.
- [41] Lequin MB, Verbaan D, Jacobs WCH, Brand R, Bouma GJ, Vandertop WP, et al. Surgery versus prolonged conservative treatment for sciatica: 5-year results of a randomised controlled trial. *BMJ Open* 2013;3:e002534.
- [42] Krugluger J, Knahr K. Chemonucleolysis and automated percutaneous discectomy – a prospective randomized comparison. *Int Orthop* 2000;24:167-9.
- [43] Chatterjee S, Foy PM, Findlay GF. Report of a controlled clinical trial comparing automated percutaneous lumbar discectomy and microdiscectomy in the treatment of contained lumbar disc herniation. *Spine (Phila PA 1976)* 1995;20:734-8.
- [44] Haines SJ, Jordan N, Boen JR, Nyman JA, Oldridge NB, Lindgren BR. Discectomy strategies for lumbar disc herniation: results of the LAPDOG trial. *J Clin Neurosci* 2002;9:411-7.
- [45] Revel M, Payan C, Vallee C, Laredo JD, Lassale B, Roux C, et al. Automated percutaneous lumbar discectomy versus chemonucleolysis in the treatment of sciatica. A randomized multicenter trial. *Spine (Phila PA 1976)* 1993;18:1-7.
- [46] Sasani M, Ozer AF, Oktenoglu T, Canbulat N, Sarioglu AC. Percutaneous endoscopic discectomy for far lateral lumbar disc herniations: prospective study and outcome of 66 patients. *Minim Invasive Neurosurg* 2007;50:91-7.
- [47] Hanaoka T, Sawaumi A, Ohtsuka K, Nakata K, Taguchi K, Shimojima O. Percutaneous lumbar nucleotomy. *J Spinal Disord* 1996;9:257-64.
- [48] Degobbi A, Crucil M, Alberti M, Bortolussi A. A long-term review of 50 patients out of 506 treated with automated percutaneous nucleotomy according to Onik for lumbar-sacral disc herniation. *Acta Neurochir Suppl* 2005;92:103-5.
- [49] Liu W-G, Wu X-T, Guo J-H, Zhuang S-Y, Teng G-J. Long-term outcomes of patients with lumbar disc herniation treated with percutaneous discectomy: comparative study with microendoscopic discectomy. *Cardiovasc Intervent Radiol* 2010;33:780-6.
- [50] Schaufele MK. Single level lumbar disc herniations resulting in radicular pain: pain and functional outcomes after treatment with targeted disc decompression. *Pain Med* 2008;9:835-43.
- [51] Tullberg T, Isacson J, Weidenhielm L. Does microscopic removal of lumbar disc herniation lead to better results than the standard procedure? Results of a one-year randomized study. *Spine (Phila PA 1976)* 1993;18:24-7.
- [52] Mayer HM, Brock M. Percutaneous endoscopic discectomy: surgical technique and preliminary results compared to microsurgical discectomy. *J Neurosurg* 1993;78:216-25.
- [53] Lagarrigue J, Chaynes P. Comparative study of disk surgery with or without microscopy. A prospective study of 80 cases. *Neurochirurgie* 1994;40:116-20.
- [54] Henriksen L, Schmidt K, Eskesen V, Jantzen E. A controlled study of microsurgical versus standard lumbar discectomy. *Br J Neurosurg* 1996;10:289-93.
- [55] Hermantin FU, Peters T, Quartararo L, Kambin P. A prospective, randomized study comparing the results of open discectomy with those of video-assisted arthroscopic microdiscectomy. *J Bone Joint Surg Am* 1999;81:958-65.
- [56] Türeyen K. One-level one-sided lumbar disc surgery with and without microscopic assistance: 1-year outcome in 114 consecutive patients. *J Neurosurg* 2003;99:247-50.
- [57] Huang T-J, Hsu RW-W, Li Y-Y, Cheng C-C. Less systemic cytokine response in patients following microendoscopic versus open lumbar discectomy. *J Orthop Res* 2005;23:406-11.
- [58] Thomé C, Barth M, Scharf J, Schmiedek P. Outcome after lumbar sequestrectomy compared with microdiscectomy: a prospective randomized study. *J Neurosurg Spine* 2005;2:271-8.
- [59] Katayama Y, Matsuyama Y, Yoshihara H, Sakai Y, Nakamura H, Nakashima S, et al. Comparison of surgical outcomes between macro discectomy and micro discectomy for lumbar disc herniation: a prospective randomized study with surgery performed by the same spine surgeon. *J Spinal Disord Tech* 2006;19:344-7.
- [60] Righesso O, Falavigna A, Avanzi O. Comparison of open discectomy with microendoscopic discectomy in lumbar disc herniations: results of a randomized controlled trial. *Neurosurgery* 2007;61:545-9. discussion 549.
- [61] Shin DA, Kim KN, Shin HC, Yoon DH. The efficacy of microendoscopic discectomy in reducing iatrogenic muscle injury. *J Neurosurg Spine* 2008;8:39-43.

- [62] Brock M, Kunkel P, Papaverio L. Lumbar microdiscectomy: subperiosteal versus transmuscular approach and influence on the early postoperative analgesic consumption. *Eur Spine J* 2008;17:518-22.
- [63] Ryang Y-M, Oertel MF, Mayfrank L, Gilsbach JM, Rohde V. Standard open microdiscectomy versus minimal access trocar microdiscectomy: results of a prospective randomized study. *Neurosurgery* 2008;62:174-81. discussion 181-2.
- [64] Franke J, Greiner-Perth R, Boehm H, Mahlfeld K, Grasshoff H, Allam Y, et al. Comparison of a minimally invasive procedure versus standard microscopic discotomy: a prospective randomised controlled clinical trial. *Eur Spine J* 2009;18:992-1000.
- [65] Arts MP, Brand R, van den Akker ME, Koes BW, Bartels RHMA, Peul WC. Tubular discectomy vs conventional microdiscectomy for sciatica: a randomized controlled trial. *JAMA* 2009;302:149-58.
- [66] Teli M, Lovi A, Brayda-Bruno M, Zagra A, Corriero A, Giudici F, et al. Higher risk of dural tears and recurrent herniation with lumbar micro-endoscopic discectomy. *Eur Spine J* 2010;19:443-50.
- [67] Barth M, Weiss C, Thomé C. Two-year outcome after lumbar microdiscectomy versus microscopic sequestrectomy: part 1: evaluation of clinical outcome. *Spine (Phila PA 1976)* 2008;33:265-72.
- [68] Barth M, Diepers M, Weiss C, Thomé C, Diepers M, Weiss C, et al. Two-year outcome after lumbar microdiscectomy versus microscopic sequestrectomy: part 2: radiographic evaluation and correlation with clinical outcome. *Spine (Phila PA 1976)* 2008;33:273-9.
- [69] Shamji MF, Bains I, Yong E, Sutherland G, Hurlbert RJ. Treatment of herniated lumbar disk by sequestrectomy or conventional discectomy. *World Neurosurg* 2014;82 (November (5)):879-83.
- [70] Fakouri B, Patel V, Bayley E, Srinivas S. Lumbar microdiscectomy versus sequestrectomy/free fragmentectomy: a long-term (>2 y) retrospective study of the clinical outcome. *J Spinal Disord Tech* 2011;24:6-10.
- [71] Faulhauer K, Manicke C. Fragment excision versus conventional disc removal in the microsurgical treatment of herniated lumbar disc. *Acta Neurochir (Wien)* 1995;133:107-11.
- [72] Fountas KN, Kapsalaki EZ, Feltes CH, Smisson HF, Johnston KW, Vogel RL, et al. Correlation of the amount of disc removed in a lumbar microdiscectomy with long-term outcome. *Spine (Phila PA 1976)* 2004;29:2521-4 [discussion 2525-6].
- [73] Wera GD, Dean CL, Ahn UM, Marcus RE, Cassinelli EH, Bohlman HH, et al. Reherniation and failure after lumbar discectomy: a comparison of fragment excision alone versus subtotal discectomy. *J Spinal Disord Tech* 2008;21:316-9.
- [74] McGirt MJ, Ambrossi GLG, Dato G, Scubba DM, Witham TF, Wolinsky J-P, et al. Recurrent disc herniation and long-term back pain after primary lumbar discectomy: review of outcomes reported for limited versus aggressive disc removal. *Neurosurgery* 2009;64:338-44. discussion 344-5.
- [75] MacKay MA, Fischgrund JS, Herkowitz HN, Kurz LT, Hecht B, Schwartz M. The effect of interposition membrane on the outcome of lumbar laminectomy and discectomy. *Spine (Phila PA 1976)* 1995;20:1793-6.
- [76] Gambardella G, Gervasio O, Zaccone C, Puglisi E. Prevention of recurrent radicular pain after lumbar disc surgery: a prospective study. *Acta Neurochir Suppl* 2005;92:151-4.
- [77] Kim KD, Wang JC, Robertson DP, Brodke DS, BenDebba M, Block KM, et al. Reduction of leg pain and lower-extremity weakness for 1 year with Oxiplex/SP gel following laminectomy, laminotomy, and discectomy. *Neurosurg Focus* 2004;17:ECP1.
- [78] Richter HP, Kast E, Tomczak R, Besenfelder W, Gaus W. Results of applying ADCON-L gel after lumbar discectomy: the German ADCON-L study. *J Neurosurg* 2001;95:179-89.
- [79] de Tribolet N, Porchet F, Lutz TW, Gratzl O, Brotchi J, van Alphen HA, et al. Clinical assessment of a novel antiadhesion barrier gel: prospective, randomized, multicenter, clinical trial of ADCON-L to inhibit postoperative peridural fibrosis and related symptoms after lumbar discectomy. *Am J Orthop (Belle Mead NJ)* 1998;27:111-20.
- [80] Bernsmann K, Krämer J, Ziozios I, Wehmeier J, Wiese M. Lumbar micro disc surgery with and without autologous fat graft. A prospective randomized trial evaluated with reference to clinical and social factors. *Arch Orthop Trauma Surg* 2001;121:476-80.
- [81] Jensen TT, Asmussen K, Berg-Hansen EM, Lauritsen B, Manniche C, Vinterberg H, et al. First-time operation for lumbar disc herniation with or without free fat transplantation. Prospective triple-blind randomized study with reference to clinical factors and enhanced computed tomographic scan 1 year after operation. *Spine (Phila PA 1976)* 1996;21:1072-6.
- [82] Gibson JNA, Waddell G. Surgical interventions for lumbar disc prolapse: updated Cochrane review. *Spine (Phila PA 1976)* 2007;32:1735-47.
- [83] Rönnberg K, Lind B, Zoega B, Gadeholt-Göthlin G, Halldin K, Gellerstedt M, et al. Peridural scar and its relation to clinical outcome: a randomised study on surgically treated lumbar disc herniation patients. *Eur Spine J* 2008;17: 1714-20.
- [84] Fransen P. Reduction of postoperative pain after lumbar microdiscectomy with DuraSeal Xact Adhesion Barrier and Sealant System. *Spine J* 2010;10:751-61.
- [85] Hellbusch LC, Helzer-Julian M, Doran SE, Leibrock LG, Long DJ, Puccioni MJ, et al. Single-dose vs multiple-dose antibiotic prophylaxis in instrumented lumbar fusion – a prospective study. *Surg Neurol* 2008;70:622-7 [discussion 627].
- [86] Petignat C, Francioli P, Harbarth S, Regli L, Porchet F, Reverdin A, et al. Cefuroxime prophylaxis is effective in noninstrumented spine surgery: a double-blind, placebo-controlled study. *Spine (Phila PA 1976)* 2008;33:1919-24.
- [87] Takahashi H, Wada A, Iida Y, Yokoyama Y, Katori S, Hasegawa K, et al. Antimicrobial prophylaxis for spinal surgery. *J Orthop Sci* 2009;14:40-4.
- [88] van Middendorp JJ, Pull ter Gunne AF, Schuetz M, Habil D, Cohen DB, Hosman AJF, et al. A methodological systematic review on surgical site infections following spinal surgery: part 2: prophylactic treatments. *Spine (Phila PA 1976)* 2012;37:2034-45.
- [89] Dobzyniak MA, Fischgrund JS, Hankins S, Herkowitz HN. Single versus multiple dose antibiotic prophylaxis in lumbar disc surgery. *Spine (Phila PA 1976)* 2003;28:E453-5.
- [90] Khan IU, Janjua MB, Hasan S, Shah S. Surgical site infection in lumbar surgeries, pre and postoperative antibiotics and length of stay: a case study. *J Ayub Med Coll Abbottabad* 2009;21:135-8.
- [91] Núñez-Pereira S, Pellisé F, Rodríguez-Pardo D, Pigrau C, Sánchez JM, Bagó J, et al. Individualized antibiotic prophylaxis reduces surgical site infections by gram-negative bacteria in instrumented spinal surgery. *Eur Spine J* 2011;20(Suppl. 3):397-402.
- [92] Ohtori S, Inoue G, Koshi T, Yamashita M, Yamauchi K, Suzuki M, et al. Long-term intravenous administration of antibiotics for lumbar spinal surgery prolongs the duration of hospital stay and time to normalize body temperature after surgery. *Spine (Phila PA 1976)* 2008;33:2935-7.

- [93] Matsunaga S, Sakou T, Taketomi E, Ijiri K. Comparison of operative results of lumbar disc herniation in manual laborers and athletes. *Spine (Phila PA 1976)* 1993;18:2222-6.
- [94] Fu T-S, Lai P-L, Tsai T-T, Niu C-C, Chen L-H, Chen W-J. Long-term results of disc excision for recurrent lumbar disc herniation with or without posterolateral fusion. *Spine (Phila PA 1976)* 2005;30:2830-4.
- [95] Takeshima T, Kambara K, Miyata S, Ueda Y, Tamai S. Clinical and radiographic evaluation of disc excision for lumbar disc herniation with and without posterolateral fusion. *Spine (Phila PA 1976)* 2000;25:450-6.
- [96] Vaughan PA, Malcolm BW, Maistrelli GL. Results of L4-L5 disc excision alone versus disc excision and fusion. *Spine (Phila PA 1976)* 1988;13:690-5.
- [97] Frymoyer JW, Hanley E, Howe J, Kuhlmann D, Matteri R. Disc excision and spine fusion in the management of lumbar disc disease. A minimum ten-year followup. *Spine (Phila PA 1976)* 1978;3:1-6.
- [98] Bärlocher C, Benini A. Long-term results of discectomy and primary spondylodesis in treatment of lumbar disk hernia. *Schweiz Arch Neurol Psychiatr* 1994;145:14-24.
- [99] Eie N. Comparison of the results in patients operated upon for ruptured lumbar discs with and without spinal fusion. *Acta Neurochir (Wien)* 1978;41:107-13.
- [100] Otani K, Kikuchi S, Sato K. Does the fusion of a lumbar disk herniation improve the clinical outcome? An investigation with a minimum 10-year follow-up. *J Spinal Disord Tech* 2014;27:196-201.
- [101] El Shazly AA, El Wardany MA, Morsi AM. Recurrent lumbar disc herniation: a prospective comparative study of three surgical management procedures. *Asian J Neurosurg* 2013;8:139-46.
- [102] Wang J, Zhou Y, Zhang ZF, Li CQ, Zheng WJ, Liu J. Minimally invasive or open transforaminal lumbar interbody fusion as revision surgery for patients previously treated by open discectomy and decompression of the lumbar spine. *Eur Spine J* 2011;20:623-8.
- [103] Chitnavis B, Barbagallo G, Selway R, Dardis R, Hussain A, Gullan R. Posterior lumbar interbody fusion for revision disc surgery: review of 50 cases in which carbon fiber cages were implanted. *J Neurosurg* 2001;95:190-5.
- [104] Niemeyer T, Halm H, Hackenberg L, Liljenqvist U, Bövingloh AS. Post-discectomy syndrome treated with lumbar interbody fusion. *Int Orthop* 2006;30:163-6.
- [105] Weber H. Lumbar disc herniation: a controlled, prospective study with ten years of observation. *Spine (Phila PA 1976)* 1983;8:131-40.
- [106] Osterman H, Seitsalo S, Karppinen J, Malmivaara A, Osterman H, Seitsalo S, et al. Effectiveness of microdiscectomy for lumbar disc herniation: a randomized controlled trial with 2 years of follow-up. *Spine (Phila PA 1976)* 2006;31:2409-14.
- [107] Loupasis GA, Stamos K, Katonis PG, Sapkas G, Korres DS, Hartofilakidis G. Seven- to 20-year outcome of lumbar discectomy. *Spine (Phila PA 1976)* 1999;24:2313-7.
- [108] Kavanagh BP. The GRADE system for rating clinical guidelines. *PLoS Med* 2009;6:e1000094.
- [109] Sullivan MJL, Bishop SR, Pivik J. The pain catastrophizing scale: development and validation. *Psychol Assess* 1995;7:432-524.
- [110] Mirza SK, Deyo RA. Systematic review of randomized trials comparing lumbar fusion surgery to nonoperative care for treatment of chronic back pain. *Spine (Phila PA 1976)* 2007;32:816-23.
- [111] Grob D, Porchet F, Kleinstück FS, Lattig F, Jeszenszky D, Luca A, et al. A comparison of outcomes of cervical disc arthroplasty and fusion in everyday clinical practice: surgical and methodological aspects. *Eur Spine J* 2010;19:297-306.
- [112] Kreiner DS, Hwang SW, Easa JE, Resnick DK, Baisden JL, Bess S, et al. An evidence-based clinical guideline for the diagnosis and treatment of lumbar disc herniation with radiculopathy. *Spine J* 2014;14:180-91.
- [113] North American Spine Society. NASS coverage recommendations; 2012.