



Research Articles

Positive Personality Model: Which Traits Relate to Complete Mental Health as Conceived by the Dual Factor Model?

Guadalupe de la Iglesia*^{abc}, Alejandro Castro Solano^{abc}

[a] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina. [b] Universidad de Palermo, Buenos Aires, Argentina. [c] Universidad de Buenos Aires, Buenos Aires, Argentina.

Abstract

The aim of this research was to study the association between the Positive Personality Model (PPM) traits and the state of complete mental health as conceived by the Dual Factor Model (DFM). The sample was composed of 1502 Argentinean adults from the general population (age: $M = 39.79$, $SD = 14.23$; gender: 50.1% male, 49.9% female). A cluster analysis replicated the four-group classification of the DFM: complete mental health, symptomatic but content, troubled and vulnerable. The complete mental health cluster showed a significantly higher presence of sprightliness, serenity, moderation and integrity than the other groups. As for humanity, the complete mental health and the symptomatic but content groups had the highest scores. Sprightliness and serenity increased the odds of belonging to the complete mental health group while integrity and humanity decreased these odds. Implications for the Argentinean population are discussed.

Keywords: positive traits, mental health, Dual Factor Model, Positive Personality Model, personality

Psychological Thought, 2019, Vol. 12(2), 202–213, <https://doi.org/10.5964/psyct.v12i2.377>

Received: 2019-05-10. Accepted: 2019-06-29. Published (VoR): 2019-12-09.

Handling Editors: Marius Drugas, University of Oradea, Oradea, Romania; Stanislava Stoyanova, South-West University "Neofit Rilski", Blagoevgrad, Bulgaria

*Corresponding author at: Mario Bravo 1259, C1175ABW, Ciudad Autónoma de Buenos Aires, Argentina. E-mail: gdelaignlesia@gmail.com



This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License, CC BY 4.0 (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The interest in finding factors related to the presence of *mental health* is mostly driven by the possibility of interventions in terms of prevention and promotion of health. The aim is to identify which variables function as risk or protective factors of mental health. Because the development of psychology as a science has been mainly guided by the medical model, whereby mental health is the absence of pathology (Millon, 1996), the progress in psychology has been focused mainly on diagnosing and treating mental disorders. However, the absence of psychological symptoms does not guarantee the presence of mental health, especially in terms of quality of life and optimal functioning (Seligman & Csikszentmihalyi, 2014; World Health Organization, 2014). Therefore, research aimed at describing mental health and identifying factors related to its presence should study mental health as a whole. To do this, analyses should include the study of psychopathological symptoms and well-being signs conjointly, that is, research should encompass the *complete state of mental health*.

Complete Mental Health

A proposal comprising mental health as a whole was postulated by Greenspoon and Saklofske (2001): the *Dual-Factor Model* (DFM). This model departs from the claim that a decrease in symptomatology does not necessarily imply an increment in well-being and vice versa. The DFM proposes a combined assessment of both aspects given by four possible scenarios: (1) subjects with low symptomatology and low well-being; (2) subjects with high symptomatology and low well-being; (3) subjects with low symptomatology and high well-being; and (4) subjects with high symptomatology and high well-being. Following Suldo and Shaffer (2008), these groups may be labelled (1) *vulnerable*, (2) *troubled*, (3) *complete mental health*—also called *flourishing* (Kelly, Hills, Huebner, & McQuillin, 2012) – and (4) *symptomatic but content* – also labelled *ambivalent* (Eklund, Dowdy, Jones, & Furlong, 2010).

A similar model was developed by Keyes (2005): the *Complete State Model of Health*. This author proposes that health and sickness are two correlated unipolar dimensions that together constitute a complete state of mental health. Keyes (2005) conceives mental health as a state of *flourishing* in which individuals do not exhibit psychopathology and have signs of well-being. By contrast, a state of *languishing* refers to the individuals who have a high level of psychopathology and a low level of well-being. All other subjects that do not fit into these two categories are considered to be in a moderate state of mental health. Research that incorporated this methodology evidenced the importance of analysing symptoms and well-being together (e.g. Góngora & Castro Solano, 2017).

It has been found that individuals who present complete mental health are less likely to have a cardiovascular disease (Keyes, 2004) and report having better physical health (Suldo & Shaffer, 2008), better academic achievement (Antaramian, 2015; Antaramian, Huebner, Hills, & Valois, 2010; Guerra Vargas, 2017; Lyons, Huebner, & Hills, 2013; Suldo & Shaffer, 2008; Suldo, Thalji, & Ferron, 2011), fewer behaviour problems (Eklund et al., 2010), more hope and gratitude (Eklund et al., 2010) and less internalising behaviour (Smith, 2018). Although the greatest contrasts in these findings are mostly between the *complete mental health* group and the *troubled* group, Antaramian et al. (2010) stated that the *vulnerable* group—characterised by low symptomatology and low well-being—was as likely to have low academic achievement and to exhibit problematic behaviour as the *troubled* group. Among the investigations on the DFM, Trompetter, Lamers, Westerhof, Fledderus, and Bohlmeijer's (2017) research is interesting since they concluded that it is fundamental to measure progress in psychotherapy from a DFM perspective.

Different variables have been found to affect the state of complete mental health as conceived by the DFM. Social support, for example, has been reported to positively influence the presence of complete mental health (Kelly et al., 2012; Magalhães & Calheiros, 2017). Concerning personality traits, different studies using the Five Factor Model (FFM; Costa & McCrae, 1985) classification concluded that neuroticism increases the odds of belonging to the *troubled*, *symptomatic but content* and *vulnerable* groups and that extraversion increases the odds of belonging to the *complete mental health* group (Lyons, Huebner, Hills, & Shinkareva, 2012; MacMahan, 2013). The relative novelty of the DMF lies in the little empirical evidence of the association between personality traits and complete mental health. As personality traits are characterised as consistent patterns of behaviour, emotion and thought that are relatively stable over time and identifiable in different contexts (e.g., Allport, 1937; Cattell, 1965), they are central aspects of any individual and, therefore, determine many of his or her experiences. The state of complete mental health should not be an exception.

Positive Personality Model: Positive Personality Traits

De la Iglesia and Castro Solano (2018) have recently postulated a new personality trait model that assesses traits in their positive versions: the *Positive Personality Model* (PPM). This sanity model is associated with the *pathological traits model* proposed in Section III of the Diagnostic and Statistical Manual of Mental Disorders in its fifth edition (DSM-5; American Psychiatric Association, 2013) as well the FFM (Costa & McCrae, 1985). The PPM constitutes a positive pole in the personality traits *continuum*, completing the notion of dimensional continuity in the study of personality traits (e.g., Leary, 1957; Millon, 1996; Millon & Everly, 1994; Offer & Sabshin, 1991; Strack & Lorr, 1994).

The availability of a conceptual framework that integrates pathological and normal classifications has generated an increased interest in the scientific community and applied fields (e.g. Huppert & So, 2013; Leising, Rogers, & Ostner, 2009). The positive traits that make up the PPM are as follows: serenity (negative affectivity in the pathological version and emotional stability in the FFM), characterised by an almost imperturbable state of peace and calmness; humanity (detachment in the pathological version and extraversion in the FFM), implying a high contextual sensitivity and an orientation towards others in terms of solidarity and assistance; *integrity* (antagonism in the pathological version and agreeableness in the FFM), characterised by trust, honesty and humility; moderation (disinhibition in the pathological version and responsibility in the FFM), implying caution and reflection before action; and lastly, sprightliness (psychoticism in the pathological version and openness to experience in the FFM), related to having self-confidence and clear goals, being active and feeling satisfied. The ultimate aim of the PPM is to function as a sanity nosology that is updated and integrated into the latest notions in personality diagnosis.

The PPM has been empirically supported by the validation of its measurement instrument and other studies. Results showed that the positive traits are positively and partially related to FFM traits and negatively related to DSM-5 pathological traits. These relations place PPM traits in the positive pole of the traits' continuum. Additionally, when compared to normal traits (e.g. FFM), PPM positive traits showed evidence of incremental validity because they performed as better predictors of positive mental health in the general population (de la Iglesia & Castro Solano, 2018), of job performance and satisfaction (de la Iglesia, Lupano Perugini, & Castro Solano, 2019) and of academic adjustment and achievement (de la Iglesia & Castro Solano, 2019).

This research aimed at analysing PPM positive traits in relation to the DFM. The specific objectives were (a) to determine whether the positive personality traits are related to psychological symptoms and well-being; (b) to investigate whether the presence of positive traits differs across the different groups configured according to the DFM; and (c) to test the predictive power of PPM positive traits in the odds of belonging to the DFM groups.

Method

Participants

The sample was composed of 1502 Argentinean adults from the general population. The mean age was 39.79 ($SD = 14.23$, Min = 18, Max = 83, 50.1% male, 49.9% female). Most of them (66.5%) lived in Buenos Aires City, 26.4% ($n = 396$) in the greater metropolitan area of Buenos Aires City, and 7.1% in some of the country's provinces. As for their maximum educational level, 7.3% of the participants had not completed high-school, 14.7%

had finished high-school, 35.3% were attending college, 35.5% ($n = 533$) had completed college, 2.1% ($n = 31$) were attending a postgraduate course and 5.1% had obtained a postgraduate degree. With respect to their socioeconomic status (SES), most participants (67.8%) reported middle SES, 20.3% upper-middle SES, 10.3% lower-middle SES, 0.9% high SES and 0.8% low SES.

Materials

Positive Traits Inventory–5

The *Positive Traits Inventory–5* (PTI-5; de la Iglesia & Castro Solano, 2018) is a positive personality traits measure composed of 60 items that are answered in a six-point Likert-type scale ranging from 0 (*completely false*) to 6 (*completely true*). This test is a positive reverse version of the *Personality Inventory for DSM-5 – Adult* (PID-5; Krueger, Derringer, Markon, Watson, & Skodol, 2012), a scale of pathological traits included in DSM-5 (American Psychiatric Association, 2013). The positive traits assessed are sprightliness (17 items), integrity (13 items), serenity (13 items), humanity (8 items) and moderation (9 items). PTI-5 has sound psychometric properties regarding its factorial structure and internal consistency. In this sample, Cronbach's alpha was as follows: sprightliness $\alpha = .90$, integrity $\alpha = .85$, serenity $\alpha = .88$, humanity $\alpha = .80$ and moderation $\alpha = .84$.

Symptom Checklist 90 Revised

The *Symptom Checklist 90 Revised* (SCL-90-R; Derogatis, 1977), a 90-item scale, is used to measure the presence of psychological symptoms in the past seven days. The respondent is asked to indicate how much he or she was bothered by a number of symptoms on a five-point Likert scale (1 = *not at all* to 5 = *extremely*). It has nine subscales, but in this study only the Global Severity Index was used. This measure informs about the general psychological distress of the individual assessed. Psychometric studies in the Argentinean population have shown good psychometric properties (Casullo & Castro Solano, 1999; Sánchez & Ledesma, 2009). In this study, the internal consistency measured by Cronbach's alpha was .97.

Mental Health Continuum–Short Form

The *Mental Health Continuum–Short Form* (MHC-SF; Keyes, 2005) is a 14-item measure of well-being. Three subscales can be assessed: emotional, psychological and social. It uses a six-point Likert scale based on how often the respondent has felt different well-being states (0 = *never* to 5 = *everyday*). Its factorial structure was confirmed in its local adaptation, where evidence of convergent validity was also obtained (Lupano Perugini, de la Iglesia, Castro Solano, & Keyes, 2017). Cronbach's alpha for the total score in this sample was .88.

Results

Associations Among Positive Traits, Psychological Symptoms and Well-Being

Product-moment Pearson correlations were calculated in order to explore the associations between PPM positive traits and the presence of psychological symptoms and well-being. In the global severity index—a measure of psychological symptoms—all correlations were statistically significant and negative ($p < .01$). The strongest association was with sprightliness, $r(1500) = -.47$, $p < .01$. Moderate associations were found with serenity, $r(1500) = -.34$, $p < .01$, and moderation, $r(1500) = -.24$, $p < .01$, while the association with integrity, $r(1500) = -.19$, $p < .01$ and humanity $r(1500) = -.14$, $p < .01$ was weak. Regarding well-being, all associations with PPM

positive traits were statistically significant and positive ($p < .01$). Again, the strongest correlation was observed for sprightliness, $r(1500) = .52$, $p < .01$. The associations with serenity, $r(1500) = .35$, $p < .01$, moderation, $r(1500) = .20$, $p < .01$, integrity, $r(1500) = .29$, $p < .01$ and humanity, $r(1500) = .21$, $p < .01$ were all moderate.

DFM Clusters: Differences in Positive Traits

A hierarchical cluster analysis was used to identify groups according to DFM (combination of psychological symptoms and well-being). The method of choice was Ward, and the squared Euclidean distance was used to determine the adequate number of clusters that maximised the between-groups difference and minimised the within-groups difference. The conglomerate coefficient and the dendrogram suggested that a four-group solution was appropriate.

Then, four groups were classified in the k means cluster analysis. The ANOVAs showed that statistically significant differences existed in the presence of psychological symptoms and well-being regarding the cluster ($p < .01$). Table 1 includes the means for the presence of psychological symptoms (SCL-90-R's global severity index) and well-being (MHC's total score) for each cluster. It may be observed that cluster 1 scored high on psychological symptoms and well-being, corresponding to the *symptomatic but content* group. Cluster 2 scored low on psychological symptoms and well-being, corresponding to the *vulnerable* group. Cluster 3 scored high on psychological symptoms and low on well-being, corresponding to the *troubled* group. And finally, cluster 4 scored low on psychological symptoms and high on well-being, corresponding to the *complete mental health* group (see also Figure 1).

Table 1

Clusters According to DFM

DFM variable	Cluster				F	p	η^2
	1 (n = 241)	2 (n = 419)	3 (n = 153)	4 (n = 689)			
Global severity index (standardised score)	0.80	-0.20	2.14	-0.63	1063.72	< .001	.415
Well-being (standardised score)	0.22	-0.97	-1.28	0.74	1510.12	< .001	.502

Table 2

Dual Factor Model Clusters: Differences in Positive Traits

PPM trait	Cluster				F	p	η^2
	1 (n = 241)	2 (n = 419)	3 (n = 153)	4 (n = 689)			
Sprightliness	3.68 _b	3.54 _c	3.03 _d	4.07 _a	191.25	< .001	.27
Integrity	4.12 _b	3.97 _c	3.82 _d	4.24 _a	33.39	< .001	.06
Serenity	3.22 _b	3.26 _b	2.87 _c	3.70 _a	77.75	< .001	.13
Moderation	3.43 _b	3.46 _b	3.16 _c	3.71 _a	28.91	< .001	.05
Humanity	3.77 _a	3.42 _b	3.37 _b	3.82 _a	36.99	< .001	.06

Note. Different letters indicate statistically significant differences between groups.

Then, a MANOVA was run to compare the presence of positive traits in DFM clusters (Table 2). Statistically significant differences were found in all positive traits regarding the cluster, Wilks' $\lambda = .677$,




Figure 1. Dual-Factor Model clusters.

$F(15, 4124.67) = 41.698, p < .001, \eta^2 = .12$. Post-hoc Bonferroni tests showed that individuals grouped in the *complete mental health* cluster had higher sprightliness, integrity, serenity and moderation. The cluster with the lowest presence of these traits was the *troubled* cluster. As for humanity, the *complete mental health* and *symptomatic but content* clusters had a higher presence of this trait than those who belonged to the *troubled* and *vulnerable* clusters.

Prediction of the Odds of Belonging to the DFM Cluster and Precision in the Classification

Following the analysis strategy of Lyons et al. (2012), a multinomial logistic regression was calculated to determine whether there were statistically significant changes in the odds of belonging to one cluster in comparison to another. In all cases, the comparison was with the *complete mental health* cluster as follows: (1) *complete mental health* versus *symptomatic but content*, (2) *complete mental health* versus *vulnerable*, (3) *complete mental health* versus *troubled*.

The model was statistically significant and showed a good overall fit, $\chi^2(15) = 567.01, p < .001$, Cox-Snell $R^2 = .314$. All positive traits, except for moderation ($p > .05$), were statistically significant predictors ($p < .01$). Table 3 displays the estimated parameters for each comparison. In the comparison with the *symptomatic but content* cluster, sprightliness and serenity increased the odds of belonging to the *complete mental health* group, whereas integrity and humanity decreased these odds. The same pattern was observed in the comparison with the *vulnerable* cluster. Finally, in the comparison with the *troubled* cluster, sprightliness was the positive trait that increased the odds of belonging to the *complete mental health* cluster and integrity was the one that decreased these odds. The percentage of correct classification was 55%, that is, the model was able to

Table 3

Model Parameters of the Multinomial Logistic Regression: Predictive Model of DFM Clusters by PPM Traits

PPM trait	B (SE)	95% CI OR		
		LL	OR	UL
Symptomatic but content vs. Complete Mental Health				
Intercept	3.68 (0.70)			
Sprightliness	-1.61 (0.19)**	0.13	0.19	0.29
Integrity	0.68 (0.19)**	1.36	1.98	2.89
Serenity	-0.71 (0.12)**	0.38	0.48	0.63
Moderation	-0.09 (0.12)	0.71	0.91	1.16
Humanity	0.39 (0.13)**	1.12	1.47	1.93
Vulnerable vs. Complete Mental Health				
Intercept	6.53 (0.60)			
Sprightliness	-1.92 (0.17)**	0.10	0.14	0.20
Integrity	0.58 (0.16)**	1.30	1.80	2.47
Serenity	-0.41 (0.11)**	0.52	0.66	0.83
Moderation	0.17 (0.11)	0.96	1.19	1.47
Humanity	-0.35 (0.10)**	0.56	0.69	0.86
Troubled vs. Complete Mental Health				
Intercept	9.22 (0.81)**			
Sprightliness	-3.49 (0.25)**	0.01	0.03	0.04
Integrity	0.99 (0.22)**	1.73	2.69	4.19
Serenity	-0.61 (0.16)**	0.39	0.54	0.74
Moderation	-0.11 (0.15)	0.65	0.89	1.22
Humanity	0.08 (0.17)	0.77	1.08	1.52

Note. CI = confidence interval; LL = lower limit of CI; UL = upper limit of CI.

* $p < .05$. ** $p < .01$.

predict the odds of belonging to one of the DMF clusters with a significance level better than chance (c -statistic = .688, $p < .05$).

Discussion

The main objective of this research was to study the role that PPM traits play in the state of complete mental health. To begin with, the link between positive personality traits and mental health was explored by analysing their associations with a general measure of symptoms and one of well-being. As expected, positive correlations with the well-being measure and negative correlations with the psychopathological variable were observed. In agreement with a previous study on the PPM model (de la Iglesia & Castro Solano, 2018), positive traits were found to predict the presence of mental health. No other previous research that specifically used PPM may be used to compare these results since PPM postulation is relatively new. However, given the notion of a continuum with FFM, some results may be extrapolated and compared. Neuroticism is known to positively correlate with psychological symptoms, while conscientiousness, agreeableness and extraversion are known to negatively correlate with psychological symptoms (e.g. Habibi, Sadeghi, Haghrrangbar, Madanipour, & Azarnoosh, 2013; Kotov, Gamez, Schmidt, & Watson, 2010; Malouff, Thorsteinsson, & Schutte, 2005). The op-

posite pattern has been found to occur between FFM traits and well-being (Joshani & Nosratabadi, 2009; Lamers, 2012; Lamers, Westerhof, Kovács, & Bohlmeijer, 2012). Therefore, the association pattern of PPM traits with psychological symptoms and well-being does not come as a surprise since it intrinsically shows that the presence of these traits is accompanied by the presence of well-being and the absence of psychological symptoms. Therefore, PPM traits probably function as protective factors of mental health.

However, this research emphasised the importance of studying mental health from a model that assesses both psychopathological symptoms and well-being signs conjointly. To do this, different clusters of mental health were obtained by combining the symptoms with the well-being measure. The four isolated groups replicated those proposed by the DFM (Greenspoon & Saklofske, 2001): *complete mental health*, *symptomatic but content*, *vulnerable* and *troubled*. In other words, the same configuration that was found in many other research studies (e.g. Eklund et al., 2010; Lyons et al., 2012; Suldo & Shaffer, 2008) was replicated here, thus giving the DFM more empirical evidence.

Then, the results of the MANOVAs showed that the groups differed in the presence of positive traits. As expected, the *complete mental health* group had a greater presence of sprightliness, integrity, serenity and moderation than the other groups. This is in agreement with previous studies, where the *complete mental health* group showed better results in different outcome variables (Antaramian, 2015; Antaramian et al., 2010; Eklund et al., 2010; Guerra Vargas, 2017; Keyes, 2004; Lyons et al., 2012; Smith, 2018; Suldo & Shaffer, 2008; Suldo et al., 2011). It also supports the main hypothesis of the PPM that the presence of these positive traits is associated with higher mental health, and therefore, the model can be used as a sanity nosology.

Finally, the multinomial logistic regression indicated that the PPM traits sprightliness and serenity increased the odds of belonging to the *complete mental health* group while humanity and integrity decreased these odds. This goes in line with the findings regarding FFM (Lyons, Huebner, Hills, & Shinkareva, 2012; MacMahan, 2013), where neuroticism (serenity in PPM) decreased the odds of belonging to the *complete mental health* group, but it contradicts the precedent of extraversion (humanity in PPM) increasing these odds. On the one hand, sprightliness and serenity continue to prevail as the most important traits of PPM in terms of mental health. On the other hand, integrity—a trait involving honest, sincere, reliable and humble behaviour—and humanity—a trait involving sensitivity to the individual's context, a tendency to detect others' needs and act in order to help them—were less likely to belong to the *complete mental health* group. This finding raises the question of the cultural aspects that may moderate the relation of these traits to positive outcomes. Does this phenomenon occur only in Argentina? In their pursuit of a better life in Argentina, do individuals commit dishonest and selfish acts? Are dishonesty and selfishness rewarded in in this country? Or are honest and altruistic acts not rewarded at all or not rewarded enough? This research suggests that Argentineans who act with integrity and humanity are less likely to have *complete mental health*. Future research should investigate whether this pattern replicates in other cultures.

This research does not go without limitations. The conceptualisation of traits may suggest some sort of casual thinking between the variables studied, but the cross-sectional design of the research prevents us from engaging in any possible causal interpretations. In addition, the sample mostly represents educated and middle-SES subjects and this hampers the generalisation of the study. Finally, the novelty of both PPM and DFM proposals is reflected in the few precedents in the literature and the abundant parallelism with similar or disaggregated models.

To conclude, this research gives further support to two important ideas. First, the study of mental health should include both psychopathological symptoms and well-being signs. The results obtained by the initial bivariate correlations are similar but not equal to the results obtained in the conjoint analyses. Second, positive traits continue to show an important role in different life outcomes. Personality assessment should most definitely include a sanity nosology.

Funding

This work was supported by the National Council for Scientific and Technical Research (CONICET) under grant PIP CONICET 11220150100381CO and by the University of Buenos Aires under grant UBACyT 20020150100037BA.

Competing Interests

The authors have declared that no competing interests exist.

Acknowledgments

The authors have no support to report.

References

- Allport, G. W. (1937). *Personality: A psychological interpretation*. New York, NY, USA: Holt Rinehart & Winston.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC, USA: Author.
- Antaramian, S. (2015). Assessing psychological symptoms and well-being: Application of a dual-factor mental health model to understand college student performance. *Journal of Psychoeducational Assessment*, 33(5), 419-429. <https://doi.org/10.1177/0734282914557727>
- Antaramian, S. P., Huebner, E. S., Hills, K. J., & Valois, R. F. (2010). A dual-factor model of mental health: Toward a more comprehensive understanding of youth functioning. *The American Journal of Orthopsychiatry*, 80(4), 462-472. <https://doi.org/10.1111/j.1939-0025.2010.01049.x>
- Casullo, M. M., & Castro Solano, A. (1999). Síntomas psicopatológicos en estudiantes adolescentes argentinos. Aportaciones del SCL-90-R [Psychological symptoms in Argentinean adolescent students. Contributions of SCL-90-R]. *Anuario de Investigaciones (Facultad de Psicología. Universidad de Buenos Aires)*, 7, 147-157.
- Cattell, R. B. (1965). *The scientific analysis of personality*. Baltimore, MD, USA: Penguin.
- Costa, P. T., & McCrae, R. R. (1985). *The NEO Personality Inventory Manual*. Odessa, FL, USA: Psychological Assessment Resources.
- de la Iglesia, G., & Castro Solano, A. (2018). The Positive Personality Model (PPM): A new conceptual framework for personality assessment. *Frontiers in Psychology*, 9, Article 2025. <https://doi.org/10.3389/fpsyg.2018.02027>
- de la Iglesia, G., & Castro Solano, A. (2019). *Academic achievement in college students: What role does the Positive Personality Model play?* Manuscript submitted for publication.

- de la Iglesia, G., Lupano Perugini, M. L., & Castro Solano, A. (2019). Positive Personality Model: Su asociación al funcionamiento óptimo en trabajadores activos. *Revista de Psicología*, *37*(2), 425-449. <https://doi.org/10.18800/psico.201902.003>
- Derogatis, L. R. (1977). *SCL-90-R: Administration, scoring and procedures manual*. Baltimore, MD, USA: Clinical Psychometric Research.
- Eklund, K., Dowdy, E., Jones, C., & Furlong, M. (2010). Applicability of the Dual-Factor Model of mental health for college students. *Journal of College Student Psychotherapy*, *25*, 79-92. <https://doi.org/10.1080/87568225.2011.532677>
- Góngora, V., & Castro Solano, A. (2017). Pathological personality traits (DSM-5), risk factors, and mental health. *SAGE Open*, *7*(3), 1-10. <https://doi.org/10.1177/2158244017725129>
- Greenspoon, P. J., & Saklofske, D. H. (2001). Toward an integration of subjective well-being and psychopathology. *Social Indicators Research*, *54*, 81-108. <https://doi.org/10.1023/A:1007219227883>
- Guerra Vargas, S. N. (2017). *El modelo de salud mental dual y su relación con el rendimiento académico de estudiantes universitarios* [Dual Factor Model and its relation to academic achievement in college students] (Unpublished master's thesis). Universidad Autónoma de Nuevo León, Mexico.
- Habibi, Z., Sadeghi, H., Haghrrangbar, F., Madanipour, K., & Azarnoosh, A. (2013). The study of personality characteristics and mental health in addicts. *Procedia: Social and Behavioral Sciences*, *84*, 509-513. <https://doi.org/10.1016/j.sbspro.2013.06.594>
- Huppert, F. A., & So, T. T. C. (2013). Flourishing across Europe: Application of a new conceptual framework for defining well-being. *Social Indicators Research*, *110*, 837-861. <https://doi.org/10.1007/s11205-011-9966-7>
- Joshanloo, M., & Nosratabadi, M. (2009). Levels of mental health continuum and personality traits. *Social Indicators Research*, *90*(2), 211-224. <https://doi.org/10.1007/s11205-008-9253-4>
- Kelly, R. M., Hills, K. J., Huebner, E. S., & McQuillin, S. D. (2012). The longitudinal stability and dynamics of group membership in the Dual-Factor Model of Mental Health: Psychosocial predictors of mental health. *Canadian Journal of School Psychology*, *27*(4), 337-355. <https://doi.org/10.1177/0829573512458505>
- Keyes, C. L. K. (2004). The nexus of cardiovascular disease and depression revisited: The complete mental health perspective and the moderating role of age and gender. *Aging & Mental Health*, *8*(3), 266-274. <https://doi.org/10.1080/13607860410001669804>
- Keyes, C. L. M. (2005). Mental illness and/or mental health? Investigating axioms of the complete state model of health. *Journal of Consulting and Clinical Psychology*, *73*, 539-548. <https://doi.org/10.1037/0022-006X.73.3.539>
- Kotov, R., Gamez, W., Schmidt, F., & Watson, D. (2010). Linking "big" personality traits to anxiety, depressive, and substance use disorders: A meta-analysis. *Psychological Bulletin*, *136*(5), 768-821. <https://doi.org/10.1037/a0020327>
- Krueger, R. F., Derringer, J., Markon, K. E., Watson, D., & Skodol, A. E. (2012). *The Personality Inventory for DSM-5*. Retrieved from PsycTESTS. <https://doi.org/10.1037/t30042-000>

- Lamers, S. M. A. (2012). *Positive Mental Health: Measurement, relevance and implications* (Doctoral dissertation, University of Twente, Twente, The Netherlands). Retrieved from <https://research.utwente.nl/en/publications/positive-mental-health-measurement-relevance-and-implications>
- Lamers, S. M. A., Westerhof, G. J., Kovács, V., & Bohlmeijer, E. T. (2012). Differential relationships in the association of the Big Five personality traits with positive mental health and psychopathology. *Journal of Research in Personality, 46*(5), 517-524. <https://doi.org/10.1016/j.jrp.2012.05.012>
- Leary, T. (1957). *Interpersonal diagnosis of personality*. New York, NY, USA: Ronald Press.
- Leising, D., Rogers, K., & Ostner, J. (2009). The undisordered personality: Normative assumptions underlying personality disorder diagnoses. *Review of General Psychology, 13*, 230-241. <https://doi.org/10.1037/a0017139>
- Lupano Perugini, M. L., de la Iglesia, G., Castro Solano, A., & Keyes, C. L. M. (2017). The Mental Health Continuum—Short Form (MHC—SF) in the Argentinean context: Confirmatory factor analysis and measurement invariance. *Europe's Journal of Psychology, 13*, 93-108. <https://doi.org/10.5964/ejop.v13i1.1163>
- Lyons, M. D., Huebner, E. S., & Hills, K. J. (2013). The Dual-Factor Model of Mental Health: A short-term longitudinal study of school-related outcomes. *Social Indicators Research, 114*(2), 549-565. <https://doi.org/10.1007/s11205-012-0161-2>
- Lyons, M. D., Huebner, E. S., Hills, K. J., & Shinkareva, S. V. (2012). The Dual-Factor Model of Mental Health: Further study of the determinants of group differences. *Canadian Journal of School Psychology, 27*(2), 183-196. <https://doi.org/10.1177/0829573512443669>
- MacMahan, M. M. (2013). *A longitudinal examination of high school students' group membership in a Dual-Factor Model of Mental Health: Stability of mental health status and predictors of change* (Unpublished doctoral dissertation). University of South Florida, Tampa, FL, USA.
- Magalhães, E., & Calheiros, M. M. (2017). A dual-factor model of mental health and social support: Evidence with adolescents in residential care. *Children and Youth Services Review, 79*, 442-449. <https://doi.org/10.1016/j.childyouth.2017.06.041>
- Malouff, J. M., Thorsteinsson, E. B., & Schutte, N. S. (2005). The relationship between the Five-Factor Model of personality and symptoms of clinical disorders: A meta-analysis. *Journal of Psychopathology and Behavioral Assessment, 27*(2), 101-114. <https://doi.org/10.1007/s10862-005-5384-y>
- Millon, T. (1996). *Disorders of personality*. New York, NY, USA: John Wiley & Sons.
- Millon, T., & Everly, G. (1994). *La Personalidad y sus Trastornos* [Personality and its disorders]. Barcelona, Spain: Martínez Roca.
- Offer, D., & Sabshin, M. (1991). *The diversity of normal behavior*. New York, NY, USA: Basic Books.
- Sánchez, R. O., & Ledesma, R. D. (2009). Análisis psicométrico del Inventario de Síntomas Revisado (SCL-90-R) en población clínica [Psychometric study of the Symptom Checklist Revised (SCL-90-R) in clinical population]. *Revista Argentina de Clínica Psicológica, 18*, 265-274.

- Seligman, M. E. P., & Csikszentmihalyi, M. (2014). Positive psychology: An introduction. In M. Csikszentmihalyi (Ed.), *Flow and the foundations of Positive Psychology* (pp. 279–298). New York, NY, USA: Springer.
https://doi.org/10.1007/978-94-017-9088-8_18
- Smith, N. D. W. (2018). *An Application of the Dual Factor Model of Mental Health in elementary school students: Implications for social functioning and psychopathology* (Doctoral dissertation, University of South Florida, Tampa, FL, USA). Retrieved from <https://scholarcommons.usf.edu/etd/7231>
- Strack, S., & Lorr, M. (1994). *Differentiating normal and abnormal personalities*. New York, NY, USA: Springer.
- Suldo, S. M., & Shaffer, E. J. (2008). Looking beyond psychopathology: The Dual-Factor Model of Mental Health in youth. *School Psychology Review*, 37(1), 52-68.
- Suldo, S., Thalji, A., & Ferron, J. (2011). Longitudinal academic outcomes predicted by early adolescents' subjective well-being, psychopathology, and mental health status yielded from a dual factor model. *The Journal of Positive Psychology*, 6(1), 17-30. <https://doi.org/10.1080/17439760.2010.536774>
- Trompeter, H. R., Lamers, S. M. A., Westerhof, G. J., Fledderus, M., & Bohlmeijer, E. T. (2017). Both positive mental health and psychopathology should be monitored in psychotherapy: Confirmation for the Dual-Factor Model in Acceptance and Commitment Therapy. *Behaviour Research and Therapy*, 91, 58-63. <https://doi.org/10.1016/j.brat.2017.01.008>
- World Health Organization. (2014). *Basic documents* (48th ed.). Geneva, Switzerland: WHO Press.

About the Authors

Guadalupe de la Iglesia has a PhD in Psychology (University of Buenos Aires, Argentina), and a degree of Specialization in Statistics for Social Sciences (University of Buenos Aires). Currently works as a researcher in CONICET (National Council of Scientific and Technological Investigations) and is a professor at the University of Buenos Aires and University of Palermo.

Alejandro Castro Solano has a PhD in Psychology (Complutense University of Madrid, Spain). Currently is the Director of the Doctoral Program in Psychology from the University of Palermo, works as a researcher in CONICET (National Council of Scientific and Technological Investigations) and is a professor at the University of Buenos Aires and University of Palermo.