



Fairfield University
DigitalCommons@Fairfield

Mathematics Faculty Publications

Mathematics Department

2005

Markov extensions for dynamical systems with holes: an application to expanding maps of the interval

Mark Demers

Fairfield University, mdemers@fairfield.edu

Follow this and additional works at: <https://digitalcommons.fairfield.edu/mathandcomputerscience-facultypubs>

Copyright 2005 Springer Verlag.

The final publication is available at www.springerlink.com

Peer Reviewed

Repository Citation

Demers, Mark, "Markov extensions for dynamical systems with holes: an application to expanding maps of the interval" (2005). *Mathematics Faculty Publications*. 40.

<https://digitalcommons.fairfield.edu/mathandcomputerscience-facultypubs/40>

Published Citation

Mark Demers, "Markov extensions for dynamical systems with holes: an application to expanding maps of the interval," *Israel Journal of Mathematics* 146:1 (2005), 189-221.

This Article is brought to you for free and open access by the Mathematics Department at DigitalCommons@Fairfield. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of DigitalCommons@Fairfield. For more information, please contact digitalcommons@fairfield.edu.

Markov Extensions for Dynamical Systems with Holes: An Application to Expanding Maps of the Interval

Mark F. Demers*

March 3, 2005

Abstract

We introduce the Markov extension, represented schematically as a tower, to the study of dynamical systems with holes. For tower maps with small holes, we prove the existence of conditionally invariant probability measures which are absolutely continuous with respect to Lebesgue measure (abbreviated a.c.c.i.m.). We develop restrictions on the Lebesgue measure of the holes and simple conditions on the dynamics of the tower which ensure existence and uniqueness in a class of Holder continuous densities. We then use these results to study the existence and properties of a.c.c.i.m. for $C^{1+\alpha}$ expanding maps of the interval with holes. We obtain the convergence of the a.c.c.i.m. to the SRB measure of the corresponding closed system as the measure of the hole shrinks to zero.

1 Introduction

The study of dynamical systems with holes was launched by Pianigiani and Yorke in their seminal paper [PY]. In it, they posed the following still open question.

Consider a particle on a billiard table with convex boundaries so that the dynamics of the particle are hyperbolic, i.e. the trajectories are unstable with respect to initial conditions. Suppose a small hole is made in the table. What are the statistical properties of the trajectories in this system? If p_n is the probability that a trajectory remains on the table until time n , what is the decay rate of p_n ? More generally, we can place a particle randomly on the table according to an initial distribution μ_0 . If μ_n represents its normalized distribution at time n (assuming the particle has not escaped by time n), does μ_n converge to some μ independent of μ_0 ? Such a measure μ is a *conditionally invariant measure* for the open billiard system.

*Department of Mathematics, Georgia Institute of Technology, Atlanta GA 30332. demers@math.gatech.edu.

Considering the billiard table with a small hole as a perturbation of the billiard table with no holes, we can pose a related question in terms of the stability of the closed system: does the conditionally invariant measure of the open system converge to the invariant measure of the closed system as the size of the hole tends to zero?

Although these questions remain open, much progress has been made in understanding the existence and properties of conditionally invariant measures for dynamical systems with holes. More generally, the problem can be stated as follows.

Let \hat{T} be a piecewise differentiable map of a Riemannian manifold \hat{M} . We take the hole to be an open set H in \hat{M} and keep track of the iterates of points until they reach the hole. Once a point enters H , it is not allowed to return.

Let $M = \hat{M} \setminus H$ and let $T = \hat{T}|(M \cap \hat{T}^{-1}M)$. A probability measure μ on M is said to be a conditionally invariant measure with respect to T if μ satisfies

$$\frac{\mu(T^{-1}A)}{\mu(T^{-1}M)} = \mu(A) \tag{1}$$

for any Borel subset A of M . The measure is called an *absolutely continuous conditionally invariant measure* (abbreviated a.c.c.i.m.) if its conditional distributions on unstable leaves are absolutely continuous with respect to the Riemannian volume.

The quantity $\lambda = \mu(T^{-1}M)$ is called the eigenvalue of the measure and $-\log \lambda$ represents the exponential rate at which mass escapes from the system. We call μ a *trivial* conditionally invariant measure if $\lambda = 0$. From the point of view of physical observables, we are interested in conditionally invariant measures whose escape rate indicates the rate at which the Riemannian volume escapes from the system. For this reason, we will restrict our attention to the existence and properties of nontrivial absolutely continuous conditionally invariant measures in this paper.

In this context, Pianigiani and Yorke [PY] studied C^2 expanding maps which admit a finite Markov partition after the introduction of holes. Their work was extended to smooth Smale horseshoes by Čencova ([C1], [C2]) and used in [CMS1] and [CMS2] to obtain a natural invariant measure on the fractal set of non-wandering points of the system. In [LM] and [R], the authors achieved similar results for open billiards satisfying a non-eclipsing condition. Recently, Chernov and Markarian ([CM1], [CM2]) studied Anosov diffeomorphisms with holes which were elements of a finite Markov partition. In [CMT1] and [CMT2], the Markov restriction on the holes was relaxed, but the results still used strongly the Markov partitions associated with Anosov diffeomorphisms.

In low-dimensional settings, efforts to drop the Markov requirements on both the map and the holes have had some success for expanding maps of the interval. A spectral analysis of the transfer operator was performed in [BK] and the stability of the spectrum was established in [KL] for perturbations of expanding maps

including small holes. More constructive techniques using bounded variation and contraction mapping arguments have been used in [BC] and [LiM] to prove the existence and properties of a.c.c.i.m. All these results assume that the potential associated with the transfer operator has bounded variation.

This brief survey highlights the classes of systems with holes which have been studied to date: expanding maps in one dimension; and in higher dimensions, systems which admit finite Markov partitions. These systems are all uniformly hyperbolic. The purpose of this paper is to develop a method to study dynamical systems with holes which relies on neither finite Markov partitions nor uniform hyperbolicity. The Markov extension is a flexible tool which does precisely this.

The systematic application of Markov extensions represented schematically by tower maps is due to Young who has used this method to study a variety of closed dynamical systems including Axiom A diffeomorphisms, piecewise hyperbolic maps, Hénon maps, logistic maps, and a class of scattering billiards, among others ([Y1],[Y2], [BY]). Chernov has recently extended the method to study the statistical properties of other chaotic systems ([Ch1],[Ch2],[Ch3]). The flexibility of the Markov extension stems from the fact that the dynamical system in question need not be uniformly hyperbolic. What matters is the average behavior of the map T between returns to a reference set Λ . This is what allows the method to be applied to Hénon maps and the logistic family. We briefly describe the main ideas of the construction.

Given a dynamical system $T : M \curvearrowright$, we choose a reference set $\Lambda \subset M$ and run the system, waiting for a forward iterate of Λ to make a “good” (Markov) return to Λ . In the context of expanding systems, a good return is one which completely covers Λ . A stopping time is then declared on the piece of Λ which makes the good return and we continue to iterate the rest until another good return is made. In this way we generate a countable partition \mathcal{Z} of Λ and a stopping time R which is constant on elements of \mathcal{Z} . Schematically, we consider Λ as the base Δ_0 of a tower and each subsequent level Δ_l is identified with the part of $T^l(\Lambda)$ which has not made a good return to Λ by time l . The related system (F, Δ) is an extension of $T : \cup_{i \geq 0} T^i(\Lambda) \curvearrowright$ for which there is a projection $\pi : \Delta \rightarrow \cup_{i \geq 0} T^i(\Lambda)$ which satisfies $\pi \circ F = T \circ \pi$.

There are three basic steps which are required for this method to yield results.

- (1) Given a dynamical system $T : M \curvearrowright$, we construct a Markov extension $F : \Delta \curvearrowright$;
- (2) we prove results about (F, Δ) using its simpler properties: namely, controlled hyperbolicity and a countable Markov structure with a certain decay rate in the measure of the elements of the partition;
- (3) we pass these results back to the original system (T, M) .

In Section 2, we focus on (2) for a tower with holes. In Section 3, we apply the abstract model to expanding maps of the interval by proving (1) and (3). The

author has also used Markov extensions to study logistic maps with holes. These results appear in a separate paper [D].

1.1 Summary of Main Results

Since important definitions and assumptions have not yet been introduced, here we state our results in general terms. Precise statements are made in Sections 2 and 3.

Our principal result concerning Markov extensions is the following:

- (1) Given a tower (F, Δ) with exponential decay in the levels and sufficiently small holes, the tower map F admits a conditionally invariant measure whose density is Hölder continuous on elements of the Markov partition on Δ . If in addition F is transitive on components of Δ , then the density is unique in the space of Hölder continuous functions and is bounded away from zero on Δ .

This result is stated precisely as Theorem 2.1 in Section 2.1.5.

Remark. Because Δ is not compact, the exponential decay in the levels of the tower is essential to the discussion of meaningful conditionally invariant measures. Without it the escape rate of Lebesgue measure from Δ may not be exponential, thus making the conditionally invariant measure a poor indicator of the escape dynamics of the system. The exponential decay together with the transitivity assumption assures that the exponential escape rate of Lebesgue measure is well-defined.

In Section 3, we apply this result to expanding maps to obtain the following:

- (2) Let T be a $C^{1+\alpha}$ expanding map of the unit interval with sufficiently small holes. Then T admits an absolutely continuous conditionally invariant measure whose density is bounded above. If $\alpha = 1$, then the density is of bounded variation. If in addition T satisfies a transitivity condition, the density is bounded away from zero on the complement of the hole, thus making the escape rate of Lebesgue measure equal to the escape rate of the a.c.c.i.m. Moreover, the eigenvalue λ satisfies the Lipschitz condition $1 - \lambda \leq \text{const} \cdot m(H)$.
- (3) If the measure of the holes shrinks to zero while the number of holes remains bounded, then the a.c.c.i.m. converges weakly to the SRB measure of the expanding map without holes (here we assume the expanding map without holes has a unique a.c.i.m.). If $\alpha = 1$, the convergence is in $L^1([0, 1])$.

These results are stated with full details as Theorems 3.2 and 3.3, respectively, in Section 3.1.3.

Remark. The results of Theorem 3.2 are proved under more technical assumptions in [BC] and [LiM] for expanding maps having potentials of bounded variation. We do not use bounded variation arguments and only require that the potential be Hölder continuous. Our only restriction on the hole is on its size, not its placement.

2 An a.c.c.i.m. for a Tower with Holes

We begin by establishing some basic notation and definitions regarding the tower map, following [Y1]. An important difference between the tower we shall describe here and that in [Y1] is that we allow multiple intervals in the base. The returns are still Markov with respect to these intervals, but need not cover the entire base. Since we will apply these results in Section 3 to expanding maps of the interval, we formulate our definitions for one-dimensional towers, but it is clear that our arguments can be extended to higher dimensional expanding towers as well.

2.1 Setting and Assumptions

2.1.1 Tower with multiple bases

The base of the tower is an interval $\hat{\Delta}_0$ which is taken to be the union of finitely many closed unit intervals whose interiors are pairwise disjoint, $\hat{\Delta}_0 = \bigcup_{i=1}^N \hat{\Delta}_0^{(i)}$. We denote one-dimensional Lebesgue measure by m . Let \mathcal{Z} be a countable partition of $\hat{\Delta}_0$ each element of which is a subinterval of one of the $\hat{\Delta}_0^{(i)}$. Given a return time function $R : \hat{\Delta}_0 \rightarrow \mathbb{Z}^+$ which is constant on each element of \mathcal{Z} , we define a tower $(\hat{\Delta}, \hat{F}, m)$ over $\hat{\Delta}_0$ by

$$\hat{\Delta} := \{(z, n) \in \hat{\Delta}_0 \times \mathbb{N} \mid n < R(z)\}.$$

We call the l^{th} level of the tower $\hat{\Delta}_l := \hat{\Delta}|_{n=l}$ and $\hat{\Delta}_l^{(i)}$ is the part of $\hat{\Delta}_l$ directly over $\hat{\Delta}_0^{(i)}$. Note that each level $\hat{\Delta}_l$ is simply a union of disjoint intervals; it is not necessarily connected since some subintervals will have returned to the base by time l . We let $\hat{\Delta}^{(i)} = \bigcup_{l=0}^{\infty} \hat{\Delta}_l^{(i)}$.

The action of $\hat{F} : \hat{\Delta} \rightarrow \hat{\Delta}$ is characterized by

$$\begin{aligned} \hat{F}(z, l) &= (z, l+1) && \text{if } l+1 < R(z) \text{ and} \\ \hat{F}^{R(z)}(\mathcal{Z}(z)) &= \hat{\Delta}_0^{(i)} && \text{for some } 1 \leq i \leq N \end{aligned}$$

where $\hat{F}^{R(z)}|_{\mathcal{Z}(z)}$ is continuous and one-to-one and $\mathcal{Z}(z)$ is the element of \mathcal{Z} containing z . We adopt a slight abuse of notation by referring to $\hat{F}(x, l)$ as $\hat{F}(x)$ and Δ_l will be made clear by context.

As mentioned in the remark in Section 1.1, an essential requirement is an exponential decay in the measure of the levels of the tower.

(H1) There exist $A > 0$ and $0 < \theta < 1$ such that $m(\hat{\Delta}_l) \leq A\theta^l$ for $l \geq 0$.

2.1.2 Introduction of holes and regularity of \hat{F}

The hole \tilde{H} in $\hat{\Delta}$ is a union of open intervals $\tilde{H}_{l,j}^{(i)}$. We set $\bigcup_j \tilde{H}_{l,j}^{(i)} =: \tilde{H}_l^{(i)} \subset \hat{\Delta}_l^{(i)}$ and require that there be only finitely many $\tilde{H}_{l,j}^{(i)}$ per level l of the tower. We define $\tilde{H}_l = \bigcup_{i=1}^N \tilde{H}_l^{(i)}$. We also require that each $\tilde{H}_{l,j}^{(i)} = \hat{F}^l(\omega)$, where ω is a union of elements of \mathcal{Z} , thus preserving the Markov structure of the returns to $\hat{\Delta}_0$. If $\hat{F}^l(\omega) = \tilde{H}_{l,j}^{(i)}$, then the intervals on all levels of the tower directly above $\tilde{H}_{l,j}^{(i)}$ are deleted since once \hat{F} maps a point into \tilde{H} , it disappears forever. The interval ω does not return to $\hat{\Delta}_0$.

Let $\Delta = \hat{\Delta} \setminus \tilde{H}$ and $\Delta_l = \hat{\Delta}_l \setminus \tilde{H}$ with analogous definitions for $\Delta^{(i)}$ and $\Delta_l^{(i)}$. We assume the existence of a countable Markov partition $\{\Delta_{l,j}^{(i)}\}$ with $\bigcup_j \Delta_{l,j}^{(i)} = \Delta_l^{(i)}$ for each i and l . Each $\Delta_{l,j}^{(i)}$ is an interval comprised of countably many elements of the form $\hat{F}^l(\omega)$, $\omega \in \mathcal{Z}$, and $\hat{F}|_{\Delta_{l,j}^{(i)}}$ is one-to-one.

In applications, $\{\Delta_{l,j}^{(i)}\}$ will be dynamically defined during the construction of the tower and its elements will be the maximal intervals which project onto the iterated pieces of the reference set Λ at time l . Uniformly hyperbolic systems, such as the expanding maps of Section 3, will typically have finitely many $\Delta_{l,j}^{(i)}$ per level l . We do not impose a finiteness requirement here, however.

We denote by $\Delta_{l,j}^{(i)*}$ those $\Delta_{l,j}^{(i)}$ whose image returns to the base, i.e. such that $\hat{F}(\Delta_{l,j}^{(i)}) = \bigcup_{k=k_1}^{k_2} \Delta_0^{(k)}$ for some $1 \leq k_1 \leq k_2 \leq N$ and set $\Delta^* = \bigcup \Delta_{l,j}^{(i)*}$.

The map \hat{F} has the following properties with respect to the partition:

- (a) \hat{F} is $C^{1+\alpha}$ on each $\Delta_{l,j}^{(i)}$ for a fixed $\alpha > 0$.
- (b) There exist $\gamma > 1$ and $\beta > 0$ such that on $\Delta_{l,j}^{(i)*}$, $|\hat{F}'| \geq \gamma e^{\beta l}$. Elsewhere, $|\hat{F}'| = 1$.
- (c) *Bounded Distortion*. There exists $C > 0$ such that for any $x, y \in \hat{\Delta}_0^{(i)}$ and $x', y' \in \Delta_{l,j}^{(k)*}$ such that $\hat{F}(x') = x$ and $\hat{F}(y') = y$ we have

$$\left| \frac{\hat{F}'(x')}{\hat{F}'(y')} - 1 \right| \leq C|x - y|^\alpha. \quad (2)$$

In applications, property (b) follows quite naturally once bounded distortion and the exponential decay of the levels are proven. In fact, $e^\beta > \frac{1}{\theta}$ and a variant of (b) is often used to prove (H1) in the construction. For uniformly hyperbolic systems it is immediate, but even for nonuniformly hyperbolic systems such as logistic maps, it is easy to obtain once (c) and (H1) are established (see [D]).

We require an additional property of the map \hat{F} . Taking C , γ and α as defined above in (a), (b) and (c), we require:

$$(H2) \quad \frac{1+C}{\gamma^\alpha} < 1.$$

This assumption says that the non-linearity of \hat{F} should not be large compared to its minimum expansion. It is controlled by shrinking the size of the reference set in the underlying dynamical system.

Remark. If $\alpha = 1$, then assumption (H2) is not necessary. The justification for this is contained in the remark at the end of Section 2.2.2.

Let $F = \hat{F}|(\Delta \setminus \hat{F}^{-1}\tilde{H})$. Note that F^m is only defined on $\bigcap_{k=0}^m \hat{F}^{-k}\Delta$, the set of points which has not fallen in the hole by time m . We say F is *transitive on components* if for all pairs i, j , $1 \leq i, j \leq N$, there exists an m such that $F^m(\Delta_0^{(i)} \cap \bigcap_{k=0}^m \hat{F}^{-k}\Delta) \supseteq \Delta_0^{(j)}$. Note that if $N = 1$, then transitivity on components is automatic as long as the hole allows at least one return to the base.

2.1.3 Definition of a combined Hölder- L^∞ Functional

The last element of our setting is a function space X on Δ in which we will seek our conditionally invariant density. The Perron-Frobenius operator associated with F acts on $L^1(\Delta)$ by

$$\mathcal{P}f(x) = \sum_{y \in F^{-1}x} \frac{f(y)}{|F'(y)|}.$$

We define $\mathcal{P}_1 f = \mathcal{P}f / |\mathcal{P}f|_{L^1}$ and seek a fixed point for the operator \mathcal{P}_1 . A fixed point for \mathcal{P}_1 is a conditionally invariant density for F .

Choose $\xi > 0$ small enough that $e^{-\xi} > \max\{\theta, e^{-\alpha\beta}\}$. This is the only restriction we will put on ξ . Given $f \in L^1(\Delta)$, let $f_{l,j}^{(i)} = f|_{\Delta_{l,j}^{(i)}}$. Let $\|f\|_\infty$ denote the L^∞ norm of f and define

$$\begin{aligned} \|f_{l,j}^{(i)}\|_\infty &= e^{-\xi l} \|f_{l,j}^{(i)}\|_\infty, \\ \|f_{l,j}^{(i)}\|_r &= \sup_{\substack{x,y \in \Delta_{l,j}^{(i)} \\ f(x) \neq 0}} \frac{|f(x) - f(y)|}{|x - y|^\alpha |f(x)|} e^{-\xi l}. \end{aligned}$$

Then define

$$\|f\| = \max\{\|f\|_\infty, \|f\|_r\}$$

where $\|f\|_\infty = \sup_{i,l,j} \|f_{l,j}^{(i)}\|_\infty$ and $\|f\|_r = \sup_{i,l,j} \|f_{l,j}^{(i)}\|_r$. It should be noted that while $\|\cdot\|_\infty$ is a norm, $\|\cdot\|_r$ is not; however, $\|\cdot\|_r$ satisfies a convex-like

inequality on a subset of X defined in Section 2.2.1. We set

$$X = \{f : \Delta \rightarrow \mathbb{R} \mid \|f\| < \infty\}.$$

2.1.4 Condition on the Holes

We formulate a single condition involving the measure of the holes which guarantees the existence of an a.c.c.i.m. in X .

Let $a := \max\{e^{-\xi}, \frac{1+C}{\gamma^\alpha}\}$ and $b := 1+C$. Note that $a < 1$ by assumption (H2). The required condition on the holes is:

$$(H3) \quad Dm\tilde{H}_0 + \sum_{l \geq 1} e^{\xi(l-1)} m\tilde{H}_l \leq \frac{(1-a)^2}{4b}$$

where $D = (1+C) \sum_{\Delta_{l,j}^{(k)*}} e^{\xi l} m\Delta_{l,j}^{(k)*}$. In typical applications of the tower constructed from a dynamical system, there will be no holes in the base of the tower. In this case $m\tilde{H}_0 = 0$, the proof simplifies somewhat and assumption (H3) reduces to

$$(H3') \quad \sum_{l \geq 1} e^{\xi(l-1)} m\tilde{H}_l \leq \frac{(1-a)^2}{b}.$$

2.1.5 Statement of results

We are now ready to state our main result of this section.

Theorem 2.1 *Given a tower with holes (Δ, F, m) with properties (a)–(c) and under assumptions (H1)–(H3), there exists a probability density φ in X such that $\mathcal{P}_1\varphi = \varphi$. If in addition F is transitive on components, then φ is the unique nontrivial conditionally invariant density in X and φ is bounded away from zero on Δ .*

2.2 Proof of Theorem 2.1

Our proof takes the following steps:

1. define a convex, compact subset X_M of X ;
2. derive Lasota-Yorke type inequalities for the operator \mathcal{P} ;
3. formulate the condition on \tilde{H} which guarantees the invariance of X_M under \mathcal{P}_1 ;
4. use transitivity to prove the uniqueness of the invariant density in X .

2.2.1 Properties of the space X and a convex, compact subset

First note that $X \subset L^1(\Delta)$. For if we let $\|\cdot\|_1$ denote the norm in $L^1(\Delta)$, then for $f \in X$ we have

$$\|f\|_1 = \sum_{i,l,j} \int_{\Delta_{l,j}^{(i)}} |f| dm \leq \sum_{i,l,j} \|f\|_{l,j}^{(i)} m(\Delta_{l,j}^{(i)}) \leq \|f\|_\infty \sum_{i,l,j} m(\Delta_{l,j}^{(i)}) e^{\xi l},$$

which is finite by the assumption on ξ . We also record for future use that if $f \in X$, then for $x, y \in \Delta_{l,j}^{(i)}$

$$\|f\|_{l,j}^{(i)} \geq \frac{|f(x) - f(y)|}{|x - y|^\alpha |f(x)|} e^{-\xi l} = \frac{\left| \frac{f(y)}{f(x)} - 1 \right|}{|x - y|^\alpha} e^{-\xi l}.$$

This in turn yields

$$\sup_{\Delta_{l,j}^{(i)}} |f| \leq (1 + \|f\|_r e^{\xi l}) \inf_{\Delta_{l,j}^{(i)}} |f| \quad (3)$$

so that f is either identically zero on $\Delta_{l,j}^{(i)}$ or f is bounded away from zero on $\Delta_{l,j}^{(i)}$.

Now let

$$X_M = \{f \in X \mid \|f\| \leq M, f \geq 0, \int_\Delta f dm = 1\}$$

where $M > 0$ has yet to be determined.

Proposition 2.2 X_M is a convex, compact subset of $L^1(\Delta)$.

Proof. We first show that X_M is convex. Let $f, g \in X_M$. Then $\|sf + (1-s)g\|_\infty \leq s\|f\|_\infty + (1-s)\|g\|_\infty \leq M$ for $s \in [0, 1]$ since $\|\cdot\|_\infty$ is a norm.

To show that $\|\cdot\|_r$ satisfies a similar convexity property, we use the fact that for any positive numbers a_1, a_2, b_1, b_2 ,

$$\frac{a_1 + a_2}{b_1 + b_2} \leq \max \left\{ \frac{a_1}{b_1}, \frac{a_2}{b_2} \right\}. \quad (4)$$

If $f \equiv 0$ on $\Delta_{l,j}^{(i)}$, then clearly $\|(sf + (1-s)g)_{l,j}^{(i)}\|_r = \|g_{l,j}^{(i)}\|_r \leq M$. So let us assume that both f and g are positive on $\Delta_{l,j}^{(i)}$. Then

$$\begin{aligned} \|(sf + (1-s)g)_{l,j}^{(i)}\|_r &\leq \sup_{x,y \in \Delta_{l,j}^{(i)}} \frac{s|f(x) - f(y)| + (1-s)|g(x) - g(y)|}{|x - y|^\alpha (sf(x) + (1-s)g(x))} e^{-\xi l} \\ &\leq \max\{\|f_{l,j}^{(i)}\|_r, \|g_{l,j}^{(i)}\|_r\} \leq M \end{aligned}$$

where we have used equation (4) for the second inequality.

Thus $sf + (1-s)g \in X_M$ whenever $f, g \in X_M$, so X_M is convex. We now show that X_M is compact as well.

Let $\{f_n\}_{n=0}^\infty$ be a sequence in X_M . For fixed i, l and j ,

$$\sup_{x, y \in \Delta_{l,j}^{(i)}} \frac{|f_n(x) - f_n(y)|}{|x - y|^\alpha} \leq \|f_n\|_r \|f_n\|_\infty e^{2\xi l}.$$

This makes $\{f_n|_{\Delta_{l,j}^{(i)}}\}$ a uniformly Hölder continuous and thus equicontinuous family of functions. It is also bounded in L^∞ . By the Ascoli-Arzelà theorem, there exists a subsequence which converges pointwise uniformly on $\Delta_{l,j}^{(i)}$ to a function $f_{*l,j}^{(i)}$. By the uniformity of the convergence, $f_{*l,j}^{(i)}$ has the same $\|\cdot\|_r$ bound as the $f_n|_{\Delta_{l,j}^{(i)}}$. It also has the same bound in L^∞ .

Using Cantor diagonalization, we obtain a subsequence which converges pointwise on all of Δ . Call this sequence $\{f_{n_k}\}$ and the limit f_* . Since $|f_{n_k}|$ is dominated by $Me^{\xi l}$ on each Δ_l , we conclude that $f_{n_k} \rightarrow f_*$ in $L^1(\Delta)$ as well.

Note that our choice of ξ already limits the size of \tilde{H} by limiting the eigenvalue of the invariant measure. Suppose we find a nontrivial conditionally invariant density $\varphi \in X_M$ and let $\varphi dm = d\mu$. If $\lambda := \mu(F^{-1}\Delta) < 1$, then we must have $\varphi|_{\Delta_{l+1}} = \frac{1}{\lambda}\varphi|_{F^{-1}(\Delta_{l+1})}$ by the conditional invariance property. Since $\varphi \in X_M$ and in particular $\|\varphi\|_\infty$ is finite, we must have $\lambda \geq e^{-\xi}$.

2.2.2 Estimates on $\|\mathcal{P}f\|$

Our goal is to show that there exists a choice of M such that \mathcal{P}_1 takes X_M into itself. We do this by first deriving Lasota-Yorke type inequalities for the operator \mathcal{P} . Note that by definition, $\mathcal{P}f(x) = f(F^{-1}x)$ for $x \in \Delta_l$, $l \geq 1$.

For $\Delta_{l,j}^{(i)}$, $l \geq 1$, we have the estimate,

$$\begin{aligned} \|\mathcal{P}f_{l,j}^{(i)}\|_\infty &:= \sup_{x \in \Delta_{l,j}^{(i)}} |\mathcal{P}f(x)| e^{-\xi l} = \sup_{x \in \Delta_{l,j}^{(i)}} |f(F^{-1}x)| e^{-\xi l} \\ &= \sup_{y \in F^{-1}\Delta_{l,j}^{(i)}} |f(y)| e^{-\xi(l-1)} e^{-\xi} \leq \|f\|_\infty e^{-\xi}. \end{aligned}$$

And similarly,

$$\begin{aligned} \|\mathcal{P}f_{l,j}^{(i)}\|_r &= \sup_{x, y \in \Delta_{l,j}^{(i)}} \frac{|f(F^{-1}x) - f(F^{-1}y)|}{|x - y|^\alpha |f(F^{-1}x)|} e^{-\xi l} \\ &= \sup_{w, z \in F^{-1}\Delta_{l,j}^{(i)}} \frac{|f(w) - f(z)|}{|w - z|^\alpha |f(w)|} e^{-\xi l} \leq \|f\|_r e^{-\xi}. \end{aligned}$$

For $\Delta_0^{(i)}$ the estimates are more involved. First note that for $x, y \in \Delta_{l,j}^{(k)*}$ with

$F(x), F(y) \in \Delta_0^{(i)}$, equation (2) yields

$$\left| \frac{1}{|F'(x)|} - \frac{1}{|F'(y)|} \right| \leq C |F(x) - F(y)|^\alpha \frac{1}{|F'(y)|} \quad (5)$$

so that

$$\frac{1}{|F'(x)|} \leq (1 + C)m\Delta_{l,j}^{(k)*} \quad (6)$$

and also

$$\left| \frac{F'(y)}{F'(x)} \right| \leq 1 + C. \quad (7)$$

Estimating $\|\mathcal{P}f_{\mathbf{0},j}^{(i)}\|_\infty$.

For $x \in \Delta_0^{(i)}$ we have $\mathcal{P}f(x) = \sum_{\substack{y \in \Delta_{l,j}^{(k)*} \\ F(y)=x}} \frac{f(y)}{|F'(y)|}$. For a fixed $\Delta_{l,j}^{(k)*}$, let $\bar{a} \in \Delta_{l,j}^{(k)*}$ be

such that $f(\bar{a}) = \frac{1}{m\Delta_{l,j}^{(k)*}} \int_{\Delta_{l,j}^{(k)*}} f dm$. Then

$$|f_{l,j}^{(k)}|_\infty \leq \frac{1}{m\Delta_{l,j}^{(k)*}} \int_{\Delta_{l,j}^{(k)*}} f dm + \sup_{x \in \Delta_{l,j}^{(k)*}} |f(x) - f(\bar{a})|. \quad (8)$$

We use (6) to estimate the first term,

$$\frac{1}{|F'(y)|m\Delta_{l,j}^{(k)*}} \int_{\Delta_{l,j}^{(k)*}} f dm \leq (1 + C) \int_{\Delta_{l,j}^{(k)*}} f dm. \quad (9)$$

We estimate the second term by,

$$\begin{aligned} \sup_{x \in \Delta_{l,j}^{(k)*}} |f(x) - f(\bar{a})| &= \sup_{x \in \Delta_{l,j}^{(k)*}} \frac{|f(x) - f(\bar{a})|}{|x - \bar{a}|^\alpha f(\bar{a})} |x - \bar{a}|^\alpha f(\bar{a}) \\ &\leq \|f_{l,j}^{(k)}\|_r e^{\xi l} (m\Delta_{l,j}^{(k)*})^\alpha \frac{1}{m\Delta_{l,j}^{(k)*}} \int_{\Delta_{l,j}^{(k)*}} f dm. \end{aligned}$$

We then use (6) and property (b) to split up $|F'|$,

$$\frac{1}{|F'(y)|} = \frac{1}{|F'(y)|^{1-\alpha}} \cdot \frac{1}{|F'(y)|^\alpha} \leq (m\Delta_{l,j}^{(k)*})^{1-\alpha} (1 + C)^{1-\alpha} \frac{1}{\gamma^\alpha} e^{-\alpha\beta l}$$

so that

$$\frac{\sup_{x \in \Delta_{l,j}^{(k)*}} |f(x) - f(\bar{a})|}{|F'(y)|} \leq \|f_{l,j}^{(k)}\|_r \frac{(1 + C)^{1-\alpha}}{\gamma^\alpha} \int_{\Delta_{l,j}^{(k)*}} f dm. \quad (10)$$

where we have used the fact that $e^\varepsilon \leq e^{\alpha\beta}$. Combining (8), (9) and (10) we obtain:

$$\begin{aligned}
\|\mathcal{P}f_{0,j}^{(i)}\|_\infty &= |\mathcal{P}f_{0,j}^{(i)}|_\infty = \sup_{x \in \Delta_{0,j}^{(i)}} \sum_{\substack{y \in \Delta_{l,j}^{(k)*} \\ F(y)=x}} \frac{f(y)}{|F'(y)|} \\
&\leq \sum_{\Delta_{l,j}^{(k)*}} (1+C) \int_{\Delta_{l,j}^{(k)*}} f dm + \|f_{l,j}^{(k)}\|_r \frac{(1+C)^{1-\alpha}}{\gamma^\alpha} \int_{\Delta_{l,j}^{(k)*}} f dm \\
&\leq (1+C) \int_{\Delta^*} f dm + \|f\|_r \frac{(1+C)^{1-\alpha}}{\gamma^\alpha} \int_{\Delta^*} f dm \quad (11) \\
&\leq \frac{(1+C)^{1-\alpha}}{\gamma^\alpha} \|f\|_r + 1 + C.
\end{aligned}$$

Estimating $\|\mathcal{P}f_{0,j}^{(i)}\|_r$.

Let $x, z \in \Delta_{0,j}^{(i)}$.

$$\begin{aligned}
|\mathcal{P}f(x) - \mathcal{P}f(z)| &\leq \sum_{\substack{y, w \in \Delta_{l,j}^{(k)*} \\ F(y)=x, F(w)=z}} \left| \frac{f(y)}{|F'(y)|} - \frac{f(w)}{|F'(w)|} \right| \\
&\leq \sum_{\substack{y, w \in \Delta_{l,j}^{(k)*} \\ F(y)=x, F(w)=z}} \left| \frac{f(y)}{|F'(y)|} - \frac{f(y)}{|F'(w)|} \right| + \left| \frac{f(y)}{|F'(w)|} - \frac{f(w)}{|F'(w)|} \right| \\
&=: \sum_{\Delta_{l,j}^{(k)*}} A_{l,j}^k + B_{l,j}^k.
\end{aligned}$$

We use the following extension of equation (4): given two convergent series, $\sum a_i$ and $\sum b_i$, $a_i \geq 0$, $b_i > 0$, then

$$\frac{\sum a_i}{\sum b_i} \leq \sup_i \frac{a_i}{b_i}. \quad (12)$$

Note that if $f(y) = 0$, then $f(w) = 0$ by (3) since y and w are in the same $\Delta_{l,j}^{(k)}$. So dropping terms where $f(y) = 0$, we may use (12) to estimate

$$\begin{aligned}
\frac{|\mathcal{P}f(x) - \mathcal{P}f(z)|}{|x - z|^\alpha |\mathcal{P}f(x)|} &\leq \frac{\sum_{\Delta_{l,j}^{(k)*}} A_{l,j}^k + \sum_{\Delta_{l,j}^{(k)*}} B_{l,j}^k}{|x - z|^\alpha \sum_{\substack{y \in \Delta_{l,j}^{(k)*} \\ F(y)=x}} \frac{f(y)}{|F'(y)|}} \\
&\leq \sup_{\substack{\Delta_{l,j}^{(k)*} \\ F(y)=x}} \frac{A_{l,j}^k}{|x - z|^\alpha \frac{f(y)}{|F'(y)|}} + \sup_{\substack{\Delta_{l,j}^{(k)*} \\ F(y)=x}} \frac{B_{l,j}^k}{|x - z|^\alpha \frac{f(y)}{|F'(y)|}}
\end{aligned}$$

For the first term,

$$\frac{A_{l,j}^k}{|x-z|^\alpha \frac{f(y)}{|F'(y)|}} = \frac{1}{|x-z|^\alpha} \left| \frac{F'(y)}{F'(w)} - 1 \right| \leq C$$

where we have used (2) for the inequality. For the second term,

$$\frac{B_{l,j}^k}{|x-z|^\alpha \frac{f(y)}{|F'(y)|}} = \frac{|f(y) - f(w)|}{|x-z|^\alpha f(y)} \cdot \frac{|F'(y)|}{|F'(w)|} \leq \|f_{l,j}^{(k)}\|_r \frac{1}{\gamma^\alpha} (1+C)$$

where we have used the fact that $|x-z| \geq \left(\inf_{\Delta_{l,j}^{(k)*}} |F'| \right) |y-w|$ in addition to equation (7) and property (b), Combining the estimates for $A_{l,j}^k$ and $B_{l,j}^k$ we see that

$$\|\mathcal{P}f_{0,j}^{(k)}\|_r \leq C + \|f\|_r \frac{1}{\gamma^\alpha} (1+C).$$

Recalling that $a = \max\{e^{-\xi}, \frac{1+C}{\gamma^\alpha}\}$ and $b = 1+C$ we have shown that

$$\|\mathcal{P}f\|_r \leq a\|f\|_r + b. \quad (13)$$

We also have

$$\|\mathcal{P}f_{0,j}^{(i)}\|_\infty \leq a\|f\|_r + b \quad \text{and} \quad \|\mathcal{P}f_{l,j}^{(i)}\|_\infty \leq a\|f\|_\infty \quad \text{for } l \geq 1 \quad (14)$$

where $a < 1$ by assumption (H2).

Remark. If $\alpha = 1$, i.e. F is C^2 on each $\Delta_{l,j}^{(i)}$, then we do not need assumption (H2). We define $\|f_{l,j}^{(i)}\|_r = e^{-\xi l} \sup_{x \in \Delta_{l,j}^{(i)}} \left\{ \frac{|f'(x)|}{|f(x)|} \mid f(x) \neq 0 \right\}$. Using this norm, we note that on $\Delta_{l,j}^{(i)}$,

$$|f(x) - f(\bar{a})| = \left| \int_{\bar{a}}^x f' dm \right| \leq \|f_{l,j}^{(i)}\|_r e^{\xi l} \int_{\Delta_{l,j}^{(i)}} f dm.$$

Using this estimate, equation (10) becomes

$$\frac{\sup_{x \in \Delta_{l,j}^{(k)*}} |f(x) - f(\bar{a})|}{|F'(y)|} \leq \|f_{l,j}^{(k)}\|_r \frac{1}{\gamma} \int_{\Delta_{l,j}^{(k)*}} f dm.$$

So $\|\mathcal{P}f_{0,j}^{(i)}\|_\infty \leq \frac{1}{\gamma} \|f\|_r + C + 1$.

The estimate on $\|\mathcal{P}f_{0,j}^{(i)}\|_r$ changes slightly as well. If we denote by $\psi_{l,j}^{(k)}$ the inverse of $F|_{\Delta_{l,j}^{(k)}}$, then for $x \in \Delta_0^{(i)}$ we have

$$\frac{|\mathcal{P}f'(x)|}{\mathcal{P}f(x)} = \frac{\sum_{\Delta_{l,j}^{(k)*}} |(f \circ \psi_{l,j}^{(k)}(x)) \cdot |(\psi_{l,j}^{(k)})'(x)||}{\sum_{\Delta_{l,j}^{(k)*}} f \circ \psi_{l,j}^{(k)}(x) \cdot |(\psi_{l,j}^{(k)})'(x)|}$$

$$\begin{aligned}
&\leq \sup_{\Delta_{l,j}^{(k)*}} \frac{|f'| \circ \psi_{l,j}^{(k)}(x) \cdot |(\psi_{l,j}^{(k)})'(x)|^2}{f \circ \psi_{l,j}^{(k)}(x) \cdot |(\psi_{l,j}^{(k)})'(x)|} + \sup_{\Delta_{l,j}^{(k)*}} \frac{f \circ \psi_{l,j}^{(k)}(x) \cdot |(\psi_{l,j}^{(k)})''(x)|}{f \circ \psi_{l,j}^{(k)}(x) \cdot |(\psi_{l,j}^{(k)})'(x)|} \\
&\leq \frac{1}{\gamma} \|f\|_r + C
\end{aligned}$$

where we have used (12) in the second step and (5) in the second term of the third. In this case, $a := \max\{e^{-\xi}, \frac{1}{\gamma}\}$ which is automatically less than one.

2.2.3 Choice of M and condition on \tilde{H}

Choose $M = \frac{2b}{1-a}$. We derive a condition on the size of the hole depending on a, b and ξ which guarantees that \mathcal{P}_1 will map X_M into itself. This will be precisely assumption (H3) introduced in Section 2.1.4.

Since $\|\cdot\|_r$ is invariant under constant multiples of a given function f , we have $\|\mathcal{P}_1 f\|_r = \|\mathcal{P}f\|_r$. Since $b = \frac{1-a}{2}M$, using equation (13) we have

$$\|\mathcal{P}_1 f\|_r = \|\mathcal{P}f\|_r \leq aM + b = aM + \frac{1-a}{2}M < M.$$

The $\|\cdot\|_\infty$ norm is not invariant under normalization, however, so we need to introduce a restriction on the size of the holes. First note that for $f \in X_M$

$$\begin{aligned}
|\mathcal{P}f|_1 &= \int_{\Delta \setminus (\hat{F}^{-1}\tilde{H})} f dm = 1 - \int_{\hat{F}^{-1}(\tilde{H}) \cap \Delta} f dm \\
&= 1 - \sum_{l \geq 1} \int_{\hat{F}^{-1}\tilde{H}_l} f dm - \int_{\hat{F}^{-1}(\tilde{H}_0) \cap \Delta} f dm \\
&\geq 1 - \sum_{l \geq 1} |f_{l-1}|_\infty m(\hat{F}^{-1}\tilde{H}_l) - \sum_{\Delta_{l,j}^{(k)*}} \int_{\hat{F}^{-1}(\tilde{H}_0) \cap \Delta_{l,j}^{(k)*}} f dm \\
&\geq 1 - \sum_{l \geq 1} \|f_{l-1}\|_\infty e^{\xi(l-1)} m\tilde{H}_l - \sum_{\Delta_{l,j}^{(k)*}} \|f_{l,j}\|_\infty e^{\xi l} m\tilde{H}_0 m\Delta_{l,j}^{(k)*} (1+C) \\
&\geq 1 - \|f\| \left(\sum_{l \geq 1} e^{\xi(l-1)} m\tilde{H}_l + Dm\tilde{H}_0 \right).
\end{aligned}$$

where $D = (1+C) \sum_{\Delta_{l,j}^{(k)*}} e^{\xi l} m\Delta_{l,j}^{(k)*}$ as before. To simplify this expression, let $q := Dm\tilde{H}_0 + \sum_{l \geq 1} e^{\xi(l-1)} m\tilde{H}_l$ and note that $q \rightarrow 0$ as $m\tilde{H} \rightarrow 0$. We require that \tilde{H} be small enough so that

$$q \leq \frac{1-a}{2M} = \frac{(1-a)^2}{4b}$$

which is precisely assumption (H3). Using this bound on \tilde{H} , we conclude that

$$\begin{aligned}\|\mathcal{P}_1 f\|_\infty &\leq \frac{a\|f\| + b}{|\mathcal{P}f|_1} \leq \frac{a\|f\| + b}{1 - q\|f\|} \\ &\leq \frac{aM + \frac{1-a}{2}M}{1 - \frac{1-a}{2}M} = M.\end{aligned}$$

Note that this estimate implies that $\lambda \geq \frac{1+a}{2}$.

In the case when there are no holes in the base, we may use the simpler assumption (H3'). Now choose $M = \frac{b}{1-a}$. Repeating our previous argument, we have $\|\mathcal{P}_1 f\|_r \leq M$ when $\|f\|_r \leq M$.

For the $\|\cdot\|_\infty$ norm, note that equation (11) yields

$$\|\mathcal{P}f_{0,j}^{(i)}\|_\infty \leq (a\|f\|_r + b) \int_{\Delta \setminus (\hat{F}^{-1}\tilde{H})} f dm$$

since $\Delta^* \subseteq \Delta \setminus (\hat{F}^{-1}\tilde{H})$ when there are no holes in the base. But $|\mathcal{P}f|_1 = \int_{\Delta \setminus (\hat{F}^{-1}\tilde{H})} f dm$ so we have $\|\mathcal{P}_1 f_{0,j}^{(i)}\|_\infty \leq aM + b \leq M$.

All that remains is the estimate on $\|\mathcal{P}_1 f_{l,j}^{(i)}\|_\infty$ for $l \geq 1$.

$$\|\mathcal{P}_1 f_{l,j}^{(i)}\|_\infty \leq \frac{aM}{|\mathcal{P}f|_1} \leq \frac{aM}{1 - qM}.$$

This is less than or equal to M if $q \leq \frac{1-a}{M} = \frac{(1-a)^2}{b}$, which is precisely assumption (H3').

Since \mathcal{P}_1 takes X_M into itself and X_M is a convex, compact subset of $L^1(\Delta)$, we can apply the Schauder-Tychonov Theorem to conclude that \mathcal{P}_1 has a fixed point $\varphi \in X_M$ once we demonstrate that \mathcal{P}_1 is a continuous map on X_M . This follows from the estimate

$$\begin{aligned}|\mathcal{P}_1 f - \mathcal{P}_1 g|_1 &\leq \left| \frac{\mathcal{P}f}{|\mathcal{P}f|_1} - \frac{\mathcal{P}g}{|\mathcal{P}g|_1} \right|_1 + \left| \frac{\mathcal{P}g}{|\mathcal{P}f|_1} - \frac{\mathcal{P}g}{|\mathcal{P}g|_1} \right|_1 \\ &\leq \frac{|f - g|_1}{|\mathcal{P}f|_1} + \frac{||\mathcal{P}g|_1 - |\mathcal{P}f|_1|}{|\mathcal{P}f|_1} \\ &\leq \frac{|f - g|_1}{|\mathcal{P}f|_1} + \frac{|\mathcal{P}f - \mathcal{P}g|_1}{|\mathcal{P}f|_1} \leq 2 \frac{|f - g|_1}{|\mathcal{P}f|_1}\end{aligned}$$

together with the fact that $|\mathcal{P}f|_1 \geq 1 - qM$ for $f \in X_M$. Setting $d\mu = \varphi dm$, we have a measure that is conditionally invariant with respect to F and absolutely continuous with respect to Lebesgue measure. The eigenvalue $\lambda = |\mathcal{P}\varphi|_1$ satisfies $\lambda \geq 1 - qM$ by our estimates above and so $\lambda \rightarrow 1$ as $m\tilde{H} \rightarrow 0$.

2.2.4 Uniqueness of the conditionally invariant measure

We now prove that φ is the unique nontrivial conditionally invariant density in the function space X if F is transitive on components. Recall from (3) that if $f \in X$, then on each $\Delta_{l,j}^{(i)}$, $\sup_{\Delta_{l,j}^{(i)}} |f| \leq (1 + \|f\|e^{\xi l}) \inf_{\Delta_{l,j}^{(i)}} |f|$, so that $f_{l,j}^{(i)}$ is either identically zero or bounded away from zero. For a conditionally invariant density φ with eigenvalue $0 < \lambda \leq 1$, if $\varphi \equiv 0$ on some $\Delta_{0,j}^{(i)}$, then $\varphi \equiv 0$ on Δ by transitivity. And since if $\hat{F}(\Delta_{l,j}^{(k)*})$ intersects $\Delta_0^{(i)}$, it crosses it completely, we must have $\inf_{\Delta_0} \varphi = \delta$ for some $\delta > 0$. But by the conditional invariance property, $\varphi|_{\Delta_{l+1}} = \frac{1}{\lambda} \varphi|_{F^{-1}(\Delta_{l+1})}$ for $l \geq 0$. This makes $\inf_{\Delta} \varphi = \inf_{\Delta_0} \varphi$.

Proposition 2.3 *If $\varphi \in X$ is a nontrivial conditionally invariant density for F , then $\varphi \in X_M$.*

Proof. From equation (13), we have $\|\mathcal{P}_1 \varphi\|_r \leq a \|\varphi\|_r + b$ which implies

$$\|\varphi\|_r = \|\mathcal{P}_1^n \varphi\|_r \leq a^n \|\varphi\|_r + \frac{b}{1-a} \text{ for all } n.$$

Thus $\|\varphi\|_r \leq \frac{b}{1-a}$.

For the $\|\cdot\|_\infty$ norm, the conditional invariance property implies that

$$\|\varphi_{l,j}^{(i)}\|_\infty = e^{-\xi l} |\varphi_{l,j}^{(i)}|_\infty \leq e^{-\xi l} \frac{1}{\lambda^l} |\varphi|_{\Delta_0}|_\infty \leq \|\varphi|_{\Delta_0}\|_\infty$$

since $\lambda \geq e^{-\xi}$.

From equation (14), we have $\|\mathcal{P} \varphi_{0,j}^{(i)}\|_\infty \leq a \|\varphi\|_r + b$.

If there are holes in the base, then $\lambda \geq \frac{1+a}{2}$ so

$$\|\mathcal{P}_1 \varphi_{0,j}^{(i)}\|_\infty = \frac{\|\mathcal{P} \varphi_{0,j}^{(i)}\|_\infty}{\lambda} \leq \frac{a \frac{b}{1-a} + b}{\lambda} \leq \frac{2b}{1-a}.$$

If there are no holes in the base, then equation (11) yields

$$\|\mathcal{P}_1 \varphi_{0,j}^{(i)}\|_\infty \leq \frac{(a \|\varphi\|_r + b)}{\lambda} \int_{\Delta \setminus \hat{F}^{-1} \hat{H}} \varphi dm \leq a \frac{b}{1-a} + b = \frac{b}{1-a}.$$

So whether there are holes in the base and we define $M = \frac{2b}{1-a}$ or there are no holes in the base and we define $M = \frac{b}{1-a}$, we have $\varphi \in X_M$. \square

Now assume that there are two different nontrivial conditionally invariant densities φ_1 and φ_2 in X . Suppose they both have the same eigenvalue λ . For $s \in \mathbb{R}$, let $\varphi_s = s\varphi_1 + (1-s)\varphi_2$. Then φ_s is a conditionally invariant density for each s as long as $\varphi_s > 0$. This is an open condition since $\inf_{\Delta} \varphi_s = \inf_{\Delta_0} \varphi_s$ and will certainly be true for $s \in [0, 1]$ since X_M is convex. Let s_0 be the first $s > 1$

such that $\inf_{\Delta} \varphi_{s_0} = 0$. For each $s < s_0$, $\varphi_s \in X_M$ by Proposition 2.3 so that $\varphi_{s_0} \in X_M$ as well since X_M is closed. But then we must have φ_{s_0} identically equal to zero which is impossible since $|\varphi_{s_0}|_1 = 1$.

Now suppose that φ_1 and φ_2 have different eigenvalues $\lambda_1 > \lambda_2$. Since φ_2 is bounded away from zero and φ_1 is bounded above on Δ_0 , there exists a constant L such that $L\varphi_2 > \varphi_1$ on Δ_0 . We also know that $\varphi_1|_{\Delta_{i+1}} = \frac{1}{\lambda_1} \varphi_1|_{F^{-1}(\Delta_{i+1})}$ by the conditional invariance property and similarly for φ_2 . Since $\frac{1}{\lambda_2} > \frac{1}{\lambda_1}$, then $L\varphi_2 > \varphi_1$ on all of Δ . So by the positivity of the operator \mathcal{P} , we must have $\mathcal{P}^n(L\varphi_2) > \mathcal{P}^n\varphi_1$ for each $n \geq 0$. But this implies that $\lambda_2^n L\varphi_2 > \lambda_1^n \varphi_1$ for each n which is impossible since $\lambda_1 > \lambda_2$.

3 Expanding Maps with Holes: An Application

We describe an application of the tower model with holes to a class of open chaotic dynamical systems. By constructing a tower with multiple bases and applying the results of the previous section, we study the existence and properties of an absolutely continuous conditionally invariant measure for $C^{1+\alpha}$ expanding maps of the interval with holes.

3.1 Setting and Statement of Results

3.1.1 Properties of the Expanding Map

Let \hat{T} be an expanding map of the unit interval, $\hat{I} = [0, 1]$. Denote by \hat{I}_j , $1 \leq j \leq \hat{K}$, the intervals of monotonicity for \hat{T} . We assume that

- (a) \hat{T} is $C^{1+\alpha}$ on each \hat{I}_j .
- (b) $|\hat{T}'| \geq \mu > 2$.

Property (a) implies that there exists $\hat{C} > 0$ such that $|\hat{T}'(x) - \hat{T}'(y)| \leq \hat{C}|x - y|^\alpha$ whenever x, y are in the same \hat{I}_j . We let m denote Lebesgue measure on the tower and on \hat{I} interchangeably.

Now let \hat{I}_j^n be the intervals of monotonicity for \hat{T}^n . Let $\psi_{n,j}$ denote the inverse of \hat{T}^n acting on \hat{I}_j^n . Then $|\psi'_{n,j}| \leq 1/\mu^n$ so that

$$|x - y| \leq \frac{1}{\mu^n} |\hat{T}^n(x) - \hat{T}^n(y)|$$

whenever x, y are in the same \hat{I}_j^n . This gives rise to the following familiar fact.

Lemma 3.1 (Distortion Bounds). *Let $x, y \in \hat{I}_j^n$. Then*

$$\left| \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} - 1 \right| \leq \tilde{C} |\hat{T}^n(x) - \hat{T}^n(y)|^\alpha$$

where $\tilde{C} = \exp(\frac{\hat{C}}{\mu(\mu^\alpha - 1)}) - 1$.

Proof.

$$\begin{aligned}
\log \left| \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} \right| &\leq \sum_{i=0}^{n-1} |\log |\hat{T}'(\hat{T}^i x)| - \log |\hat{T}'(\hat{T}^i y)|| \\
&\leq \sum_{i=0}^{n-1} \frac{1}{\mu} |\hat{T}'(\hat{T}^i x) - \hat{T}'(\hat{T}^i y)| \\
&\leq \sum_{i=0}^{n-1} \frac{\hat{C}}{\mu} |\hat{T}^i x - \hat{T}^i y|^\alpha \\
&\leq \sum_{i=0}^{n-1} \frac{\hat{C}}{\mu} \frac{1}{\mu^{\alpha(n-i)}} |\hat{T}^n x - \hat{T}^n y|^\alpha
\end{aligned}$$

□

3.1.2 Introduction of Holes

A hole H in $[0,1]$ is a finite union of open intervals H_l , $1 \leq l \leq L$. Let $I = [0, 1] \setminus H$. For each n , let $I^n = \cap_{i=0}^n T^{-i} I$ and let $T^n = \hat{T}^n|_{I^n}$. Let K be the number of intervals of monotonicity for T and let $\mathcal{Q} = \{I_1, \dots, I_K\}$ be the partition of I into those intervals. Denote by $I_{j,n}$ the part of I_j which remains outside the hole for the first n iterates of the map \hat{T} , i.e. $I_{j,n} = I_j \cap I^n$. Note that $(T^n)'$ satisfies the same distortion bounds as $(\hat{T}^n)'$ wherever it is defined on each \hat{I}_j^n .

Let d be the minimum length of the I_j and D their maximum length. Define $C := \tilde{C}(2\delta)^\alpha$ and choose δ small enough that

$$\frac{2^\alpha(1+C)}{\mu^\alpha} < 1.$$

We do this so that the tower we construct will satisfy the nonlinearity condition (H2). If necessary, reduce δ further so that $\delta \leq d$. (If \hat{T} is C^2 , then we simply take $\delta = d$ since (H2) is not needed.) δ represents the length scale of our reference intervals which we use to construct the tower Δ . Let $a := \max \left\{ \sqrt{\frac{2}{\mu}}, \frac{2^\alpha(1+C)}{\mu^\alpha} \right\}$. Our sole condition on the hole is that its total measure must satisfy:

$$(A1) \quad mH \leq \frac{(1-a)^2 \mu \delta^2 \left(1 - \sqrt{\frac{2}{\mu}}\right)}{1+C}.$$

This assumption ensures that the hole in the tower we construct satisfies condition (H3'). (We do not need to satisfy (H3) since there will be no holes in the base of the tower we shall construct.)

Assumption (A1) also implies the following bound, which we will use to construct the tower,

$$h \leq \delta \frac{\mu - 2}{2} \tag{15}$$

where h is the maximum length of the intervals H_l . To see that equation (15) is implied by (A1), we note that $\delta \leq \frac{1}{\mu}$ and write

$$h \leq mH \leq \frac{(1-a)^2 \mu \delta^2 \left(1 - \sqrt{\frac{2}{\mu}}\right)}{1+C} \leq \delta \left(1 - \frac{2}{\mu}\right) < \delta \left(\frac{\mu}{2} - 1\right).$$

3.1.3 Statement of Results

We first state our result on the existence of an a.c.c.i.m.

Theorem 3.2 *Let \hat{T} be an expanding map of the interval with properties (a) and (b) of Section 3.1.1. Suppose H is a hole satisfying (A1). Then there exists a nontrivial absolutely continuous conditionally invariant measure for T whose density is bounded above. If $\alpha = 1$, the density is of bounded variation.*

Suppose in addition that T satisfies the following transitivity property: for each interval I_j , there exists an integer n_j such that

$$I_j \cup TI_{j,1} \cup T^2 I_{j,2} \cup T^3 I_{j,3} \cup \dots \cup T^{n_j} I_{j,n_j} = I;$$

then the conditionally invariant density is bounded away from zero on I .

Remark. We shall show in Section 3.2.5 that the eigenvalue of the a.c.c.i.m. satisfies the Lipschitz bound $1 - \lambda \leq C_0 mH$ where C_0 depends only on the map \hat{T} and the lengthscale δ .

This theorem implies that Lebesgue measure has a well-defined exponential escape rate given by the a.c.c.i.m. The escape rate relates to the question of uniqueness which is subtle for open systems. It is noted in [BC] that C^2 expanding maps may have families of conditionally invariant densities of bounded variation which share the same support, but have different eigenvalues, even when those maps are transitive on components and the holes are elements of a finite Markov partition. Nevertheless, since the spectrum of the transfer operator on BV is stable in the presence of small holes ([LiM]), we are assured when $\alpha = 1$ that we have found the unique density in BV whose eigenvalue gives the escape rate of the reference measure. When generalizing to $\alpha < 1$, our densities may no longer be in BV and it is not clear in what sense we can expect uniqueness, although we still have the correct escape rate with respect to the reference measure.

Our next result concerns the convergence as $mH \rightarrow 0$ of the conditionally invariant measures to an absolutely continuous invariant measure for \hat{T} with no holes.

Let \hat{T} and H satisfy the hypotheses of Theorem 3.2. Since we are interested in arbitrarily small holes, we may further assume that either none of the closures of the holes H_l contains an endpoint of one of the \hat{I}_j , or if one does, that H_l shrinks to this point in the limit. (If H does not satisfy this property, we may take a smaller hole which does.) In this way, we assure that no new intervals I_j are created as we take the limit $mH \rightarrow 0$. In fact, each interval I_j will either grow or remain the same as the hole shrinks.

We need the following assumption on the dynamics of \hat{T} . We assume that for each interval I_j , there exists an m_j such that

$$I_j \cup \hat{T}I_j \cup \hat{T}^2I_{j,1} \cup \hat{T}^3I_{j,2} \cup \dots \cup \hat{T}^{m_j}I_{j,m_j-1} = [0, 1]. \quad (16)$$

This means that the images of I_j eventually cover $[0,1]$, but we do not allow a part that has fallen into the hole to be considered in the future - it is only considered to cover that part of H on which it lands. This assumption is obviously stronger than the transitivity assumption of Theorem 3.2.

Note that this assumption on the dynamics of \hat{T} , in addition to the fact that \hat{T} is expanding, implies that the map \hat{T} with no holes has the covering property and so has a unique a.c.i.m. whose density is bounded away from zero on $[0,1]$.

Theorem 3.3 *Let \hat{T} and H satisfy the hypotheses of Theorem 3.2. Define $H_t = H$ and let $\{H_s\}$ for $s \in [0, t]$ be a sequence of holes with the following properties:*

- (1) $mH_s \leq s$, $H_s \subset H_t$ and each component of H_t contains at most one component of H_s ;
- (2) either the closure of H_t contains none of the endpoints of the \hat{I}_j , or if one endpoint is contained in the closure of H_t , then that endpoint is contained in the closure of H_s for all $s \in [0, t]$;
- (3) equation (16) is satisfied by \hat{T} acting on the intervals of monotonicity with respect to the hole H_t .

Let ν_s denote the a.c.c.i.m. corresponding to H_s obtained from Theorem 3.2. Then the ν_s converge weakly to ν as s goes to zero, where ν is the unique absolutely continuous invariant measure for \hat{T} with no holes. If $\alpha = 1$, then the densities converge in $L^1([0, 1])$ as well.

3.2 Proof of Theorem 3.2

Our proof takes the following steps:

1. prove a growth lemma for intervals of length at least δ ;
2. use the growth lemma to define a return time function and partition of I in order to build a tower;

3. verify that the tower has the desired properties and satisfies conditions (H1)-(H3') to conclude the existence of an a.c.c.i.m.;
4. project the measure on the tower to a measure on I and show it has the properties we claim.

3.2.1 A Growth Lemma

Although we have already fixed δ small enough to control the nonlinearities in the tower, it should be noted that the results of Lemma 3.4 hold for any δ' and h satisfying

$$\frac{2h}{\mu - 2} \leq \delta' \leq d$$

We will continue to use δ as the notation throughout this section, however.

Any interval Ω of length at least δ lying entirely in one of the I_j will have at least one subinterval which grows to cover one of the intervals of monotonicity exponentially fast. This is because if $\hat{T}\Omega$ does not cover one of the I_j , then $\mathcal{Q}|\hat{T}\Omega$ can have at most two components, and $\hat{T}\Omega$ can intersect at most one hole. The measure of what does not fall into the hole is

$$m(T(\Omega \cap I^1)) \geq \mu\delta - h \geq \mu\delta - \delta \left(\frac{\mu - 2}{2} \right) = \delta \left(\frac{\mu + 2}{2} \right),$$

where we have used equation (15) in the second inequality. So at least one of the two components of $\mathcal{Q}|\hat{T}\Omega$ must have length greater than or equal to $\delta(\frac{\mu+2}{4}) > \delta$ since $\frac{\mu+2}{4} > 1$. In fact, if we follow this component, call it ω_1 , we see that its image must also either cover one of the I_j or else $\mathcal{Q}|\hat{T}\omega_1$ has at most two components, one of whose lengths is at least $\delta(\frac{\mu+2}{4})^2$. Always following the larger component, we obtain a sequence of intervals ω_n of length at least $\delta(\frac{\mu+2}{4})^n$ which must eventually cover one of the intervals of monotonicity of T in exponential time depending only on μ , δ and D .

Lemma 3.4 *Let Ω be any interval of length at least δ lying entirely in one of the I_j . There exists a countable partition \mathcal{Z} of Ω and a stopping time $S(x)$ such that for each $\omega \in \mathcal{Z}$,*

- (a) S is constant on ω ;
- (b) \hat{T} is $C^{1+\alpha}$ on $\hat{T}^S\omega$;
- (c) Either $\omega \subset I^S$ and $T^S\omega = I_j$ for some $1 \leq j \leq K$ or $\omega \subset I^{S-1}$ and $\hat{T}^S\omega \subset H$;
- (d) $m\{x \in \Omega \mid S(x) > n\} \leq D(\frac{2}{\mu})^n$, i.e. the elements of \mathcal{Z} either fall into H or grow at an exponential rate;

$$(e) \ m\{x \in \Omega \mid \hat{T}^{S(x)}(x) \in H\} \leq \frac{mH}{\mu-2}.$$

Proof. Let $\Omega_0 := \Omega$. Given Ω_{n-1} , which consists of a finite number of subintervals of Ω , we show how to form the set Ω_n inductively. Ω_n represents those points in Ω which have not yet been assigned a stopping time by time n .

We proceed one element of Ω_{n-1} at a time. Fix $\omega \in \Omega_{n-1}$. We consider two cases depending on $\hat{T}^n \omega$.

Case 1: $\hat{T}^n \omega$ does not cover any of the I_j .

Then $\mathcal{Q}|\hat{T}^n \omega$ has at most two components and $\hat{T}^n \omega$ intersects at most one H_l . Set $S(x) = n$ on $\omega \cap \hat{T}^{-n} H$ and enter this interval as an element of the partition \mathcal{Z} . Note that this interval has length less than or equal to $\frac{h}{\mu^n}$. Put the (at most two) components of $\hat{T}^{-n} \mathcal{Q}|\omega$ into the new set Ω_n . The stopping time S has not yet been defined on the elements of Ω_n .

Case 2: $\hat{T}^n \omega$ covers at least one of the I_j .

Set $S(x) = n$ on each of the components of $\hat{T}^{-n} \mathcal{Q}|\omega$ whose image equals one of the I_j . Each of these components is entered as an element of the partition \mathcal{Z} . $\hat{T}^n \omega \cap H$ has at most L components. Set $S(x) = n$ on each of the intervals $\omega \cap \hat{T}^{-n} H_l$ and enter these as elements of the partition \mathcal{Z} . Note that $m(\omega \cap \hat{T}^{-n} H) \leq \frac{mH}{\mu^n}$. There are at most two intervals in $\hat{T}^{-n} \mathcal{Q}|\omega$ left over where S has not been defined. Include these intervals as elements of Ω_n . Note that their images under T^n each have length less than D .

By construction, Ω_n has at most 2^n components, each of length less than $\frac{D}{\mu^n}$. This implies

$$m\{x \in \Omega \mid S(x) > n\} = m\Omega_n \leq D \left(\frac{2}{\mu}\right)^n$$

so that (d) is satisfied. (a),(b) and (c) are clear from the construction of Ω_n .

To check (e), we note that the most Lebesgue measure that can fall into H at the n^{th} step is $\frac{mH}{\mu^n}$ from each component of Ω_{n-1} . This means at most $mH \frac{2^{n-1}}{\mu^n}$ falls in at each step. So the total measure of those elements of \mathcal{Z} which fall in the hole before they can grow to cover one of the intervals of monotonicity is

$$m\{x \in \Omega \mid \hat{T}^{S(x)}(x) \in H\} \leq \sum_{n=1}^{\infty} \frac{mH}{\mu} \left(\frac{2}{\mu}\right)^{n-1} = \frac{mH}{\mu-2}$$

□

3.2.2 Building the Tower

We show how to build a tower with multiple bases from the growth lemma. We partition I into N intervals of length between δ and 2δ . Call them $\Lambda^{(1)}, \dots, \Lambda^{(N)}$.

Since each $\Lambda^{(i)}$ has length at least δ , each will have a partition $\mathcal{Z}^{(i)}$ and a stopping time $S^{(i)}$ with properties (a)-(e) of Lemma 3.4.

Proceeding one $\Lambda^{(i)}$ at a time, we define below the return time function R and the partition \mathcal{Z} of I on whose elements R is constant. This will define the tower $\hat{\Delta}$. Fix k between 1 and N . On those $\omega \in \mathcal{Z}^{(k)}$ such that $\hat{T}^{S^{(k)}(\omega)}\omega \subset H$, we set $R = S^{(k)}$ and include ω as an element of \mathcal{Z} .

For those $\omega \in \mathcal{Z}^{(k)}$ with $\hat{T}^{S^{(k)}(\omega)}\omega = I_j$ for some j , we set $R = S^{(k)}$, but do not include ω as an element of \mathcal{Z} directly. I_j consists of a number of the $\Lambda^{(i)}$. We partition ω into elements of the form $\omega \cap T^{-S^{(k)}(\omega)}\Lambda^{(i)}$ and include each of these as elements of the partition \mathcal{Z} . We do this because each of the $\Lambda^{(i)}$ may have varying lengths and so their projections from the tower will have derivatives which vary accordingly. We subdivide the partition \mathcal{Q} into the intervals $\Lambda^{(i)}$ in order to have control over distortion bounds on the tower.

We arrive at a return time function R defined on I and a countable partition \mathcal{Z} which respects the boundaries of $\Lambda^{(1)}, \dots, \Lambda^{(N)}$ and satisfies the following properties for each $\omega \in \mathcal{Z}$,

- (a) R is constant on ω ;
- (b) \hat{T} is $C^{1+\alpha}$ on $\hat{T}^R\omega$;
- (c) Either $\hat{T}^R\omega \subset H$ or $\hat{T}^R\omega = \Lambda^{(i)}$ and T^R is defined on ω ;
- (d) $m\{x \in I \mid R(x) > n\} \leq ND(\frac{2}{\mu})^n$, i.e. the elements of \mathcal{Z} either fall into H or return at an exponential rate;
- (e) $m\{x \in I \mid \hat{T}^{R(x)}x \in H\} \leq N\frac{mH}{\mu-2}$

Lemma 3.1 implies that on each $\omega \in \mathcal{Z}$, we have the following distortion bound for $x, y \in \omega$ and $n \leq R(\omega)$.

$$\left| \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} - 1 \right| \leq \tilde{C}|\hat{T}^n x - \hat{T}^n y|^\alpha \quad (17)$$

We identify our reference intervals $\Lambda^{(1)}, \dots, \Lambda^{(N)}$ with N disjoint intervals of unit length, $\hat{\Delta}_0^{(1)}, \dots, \hat{\Delta}_0^{(N)}$, as follows. Let $\hat{\Delta}_0 = \bigcup_{i=1}^N \hat{\Delta}_0^{(i)}$ be as in Section 2.1.1. Define a bijection $\pi_0 : \hat{\Delta}_0 \rightarrow I$ as piecewise linear with $\pi_0(\hat{\Delta}_0^{(i)}) = \Lambda^{(i)}$ and $\pi_0'|\hat{\Delta}_0^{(i)} = |\Lambda^{(i)}|$. This makes $\delta \leq \pi_0' \leq 2\delta$ on $\hat{\Delta}_0$. The partition \mathcal{Z} and return time function R on I induce a partition and return time function on $\hat{\Delta}_0$. We call these \mathcal{Z} and R as well: they are the partition and return time function introduced in Section 2.1.1. We define as before

$$\hat{\Delta} = \{(x, n) \in \hat{\Delta}_0 \times \mathbb{N} : R(x) > n\}.$$

Recall the notation $\hat{\Delta}_l = \hat{\Delta}|_{n=l}$ for the l^{th} level of the tower, and let $\hat{\Delta}_l^{(i)}$ denote the l^{th} level above $\hat{\Delta}_0^{(i)}$. If $\hat{T}^R\omega \subset H$, then we put a hole $\tilde{H}_{R(\omega),j}$ in the $R(\omega)$

level of the tower above $\pi_0^{-1}\omega$ in $\hat{\Delta}_0$. This defines the hole \tilde{H} in the tower $\hat{\Delta}$. Let $\Delta = \hat{\Delta} \setminus \tilde{H}$ and in general $\Delta_l^{(i)} = \hat{\Delta}_l^{(i)} \setminus \tilde{H}$. Note that there are no holes in the base.

3.2.3 Properties of the Tower

Let \hat{F} be the tower map, $\hat{F} : \Delta \rightarrow \hat{\Delta}$. Define a projection $\pi : \hat{\Delta} \rightarrow [0, 1]$ such that $\pi \circ \hat{F} = \hat{T} \circ \pi$ on Δ . The elements of the Markov partition $\Delta_{l,j}^{(i)}$ are the maximal intervals on $\hat{\Delta}_l^{(i)}$ which project onto the dynamically defined elements of Ω_l in the construction of the return time function R above the reference interval $\Lambda^{(i)}$.

For $x \in \Delta_0$, we have $\pi' \circ \hat{F}^l(x) \cdot (\hat{F}^l)'(x) = (\hat{T}^l)' \circ \pi(x) \cdot \pi'(x)$ and if $R(x) > l$ then $(\hat{F}^l)'(x) = 1$ so that

$$\pi' \circ \hat{F}^l(x) = (\hat{T}^l)' \circ \pi(x) \cdot \pi'(x). \quad (18)$$

Now let $z \in \Delta_0^{(i)}$ with $R(z) = l + 1$. Let $\hat{F}^l z = x$. Then $\pi' \circ \hat{F}^{l+1}(z) \cdot (\hat{F}^{l+1})'(z) = (\hat{T}^{l+1})' \circ \pi(z) \cdot \pi'(z)$. But $\hat{F}^{l+1}(z) \in \Delta_0^{(j)}$ so that $\pi' \circ \hat{F}^{l+1}(z) = |\Lambda^{(j)}|$ and $(\hat{F}^{l+1})'(z) = \hat{F}'(\hat{F}^l z) \cdot (\hat{F}^l)'(z) = \hat{F}'(x)$. This yields

$$\hat{F}'(x) = (T^{l+1})'(\pi z) \frac{|\Lambda^{(i)}|}{|\Lambda^{(j)}|} \geq \frac{\mu^{l+1}}{2}. \quad (19)$$

We derive a distortion estimate for \hat{F} . Let $x, y \in \Delta_{l,j}^{(k)}$ be such that $\hat{F}(x), \hat{F}(y) \in \hat{\Delta}_0^{(i)}$. Using equations (17) and (19) we have

$$\begin{aligned} \left| \frac{\hat{F}'(x)}{\hat{F}'(y)} - 1 \right| &= \left| \frac{(\hat{T}^{l+1})'(\pi \circ \hat{F}^{-l}x)}{(\hat{T}^{l+1})'(\pi \circ \hat{F}^{-l}y)} - 1 \right| \\ &\leq \tilde{C} |\hat{T}^{l+1}(\pi \circ \hat{F}^{-l}x) - \hat{T}^{l+1}(\pi \circ \hat{F}^{-l}y)|^\alpha \\ &= \tilde{C} |\pi \circ \hat{F}(x) - \pi \circ \hat{F}(y)|^\alpha \\ &\leq \tilde{C} (2\delta)^\alpha |\hat{F}(x) - \hat{F}(y)|^\alpha \end{aligned}$$

Let $F = \hat{F}|(\Delta \setminus \hat{F}^{-1}\tilde{H})$. F also satisfies the relation $\pi \circ F = T \circ \pi$ on its domain, and so the above estimates hold for F .

3.2.4 (Δ, F, m) Satisfies Conditions (H1)-(H3')

Recall properties (a)-(c) required of the tower map in Section 2.1.2 as well as assumptions (H1)-(H3') of Section 2. It is clear from the discussion of Section 3.2.3 that \hat{F} has properties (a)-(c) of the tower map with $\gamma \geq \frac{\mu}{2}$, $\beta = \log \mu$, $C = \tilde{C}(2\delta)^\alpha$, and the same Hölder exponent α as the map \hat{T} .

(H1) is satisfied with $\theta = \frac{2}{\mu}$. This is because $m\hat{\Delta}_l = m\Delta_l + m\tilde{H}_l$ and we have bounds on these two quantities from the proof of Lemma 3.4. We have $m\Delta_l \leq \frac{ND}{\delta} (\frac{2}{\mu})^l$ and $m\tilde{H}_l \leq \frac{NmH}{\delta\mu} (\frac{2}{\mu})^{l-1}$. This yields the desired estimate $m\hat{\Delta}_l \leq \frac{N}{\delta} (\frac{2}{\mu})^l$.

(H2) is satisfied by choosing δ small enough.

Choose $\xi = \min\{\frac{1}{2} \log(\frac{\mu}{2}), \alpha \log \mu\}$. Then $e^\xi \leq \sqrt{\frac{\mu}{2}}$ so (H3') becomes

$$\sum_{l=1}^{\infty} \left(\frac{\mu}{2}\right)^{\frac{l-1}{2}} m\tilde{H}_l < \frac{(1-a)^2}{1+C}.$$

But $m\tilde{H}_l \leq \frac{NmH}{\delta\mu} \left(\frac{2}{\mu}\right)^{l-1}$. So

$$\begin{aligned} \sum_{l=1}^{\infty} \left(\frac{\mu}{2}\right)^{\frac{l-1}{2}} m\tilde{H}_l &\leq \sum_{l=1}^{\infty} \frac{NmH}{\delta\mu} \left(\frac{2}{\mu}\right)^{l-1} \left(\frac{\mu}{2}\right)^{\frac{l-1}{2}} \\ &= \frac{NmH}{\delta\mu} \frac{1}{1 - \sqrt{\frac{2}{\mu}}} \end{aligned}$$

Using this estimate, we see that (H3') will be satisfied if H satisfies

$$mH < \frac{(1-a)^2}{1+C} \cdot \frac{\mu\delta(1 - \sqrt{\frac{2}{\mu}})}{N}$$

which is slightly weaker than our assumption (A1).

3.2.5 Existence and properties of an a.c.c.i.m.

Since (Δ, F, m) satisfies (H1)-(H3'), we conclude using Theorem 2.1 that there exists a $\varphi \in X$ such that $d\tilde{\nu} := \varphi dm$ is a nontrivial a.c.c.i.m. with respect to F acting on Δ . Let λ be the eigenvalue of $\tilde{\nu}$. Now define a measure ν on I by

$$\nu(A) := \tilde{\nu}(\pi^{-1}A)$$

for any Borel subset A of I . Then it is clear that ν will be conditionally invariant with respect to T with the same eigenvalue λ since for any Borel $A \subset I$,

$$\begin{aligned} \nu(T^{-1}A) &:= \tilde{\nu}(\pi^{-1} \circ T^{-1}A) = \tilde{\nu}(F^{-1} \circ \pi^{-1}A) \\ &= \lambda \tilde{\nu}(\pi^{-1}A) =: \lambda \nu(A). \end{aligned}$$

We show that ν is absolutely continuous with respect to Lebesgue measure with a density that is bounded above. For this we will need to estimate the number of preimages that a Borel set A can have on each level of the tower. We consider the tower $\Delta^{(i)}$ above one of the $\Delta_0^{(i)}$. From the proof of Lemma 3.4 we know that the number of partition elements $\Delta_{l-1,j}^{(i)}$ which move up to $\Delta_l^{(i)}$ is at most 2^{l-1} (these are the preimages under π of the elements of Ω_{l-1}). Although each of these elements may be split into 2 or more pieces on $\Delta_l^{(i)}$, the map T is monotonic on each $\pi(\Delta_{l-1,j}^{(i)})$ and so $T \circ \pi(\Delta_{l-1,j}^{(i)})$ covers A at most once. Thus $\pi(\Delta_l^{(i)})$ covers A at most 2^{l-1} times.

This observation and equation (18) yield the following estimate.

$$\begin{aligned}
\nu(A) &= \sum_{k=1}^N \sum_{l=0}^{\infty} \sum_j \int_{\pi^{-1}A \cap \Delta_{l,j}^{(k)}} \varphi dm \\
&\leq \sum_{k=1}^N \sum_{l=0}^{\infty} \sum_j \sup_{\Delta_{l,j}^{(k)}} \varphi m(\pi^{-1}A \cap \Delta_{l,j}^{(k)}) \\
&\leq \sum_{k=1}^N \sum_{l=0}^{\infty} \sum_j \frac{1}{\lambda^l} \sup_{\Delta_0^{(k)}} \varphi \frac{1}{\delta \mu^l} m(A \cap \pi \Delta_{l,j}^{(k)}) \\
&\leq \sum_{k=1}^N \sum_{l=0}^{\infty} \frac{1}{\lambda^l} \sup_{\Delta_0^{(k)}} \varphi \frac{1}{\delta \mu^l} 2^{l-1} mA \\
&\leq \frac{NmA}{2\delta} \sup_{\Delta_0} \varphi \sum_{l=0}^{\infty} \left(\frac{2}{\mu}\right)^{\frac{l}{2}}
\end{aligned}$$

where we have used the fact that $\lambda \geq \sqrt{\frac{2}{\mu}}$ in the last line. This proves that $\nu \ll m$ with bounded density.

Let $d\nu = \psi dm$. Choose a $\Delta_0^{(i)}$ such that $\inf_{\Delta_0^{(i)}} \varphi = \rho > 0$. Then $\inf_{\Lambda^{(i)}} \psi \geq \frac{\rho}{2\delta}$. If T satisfies the transitivity property in the statement of Theorem 3.2, then there exists an n_0 such that the first n_0 images of $\Lambda^{(i)}$ under T cover I . Thus for any $x \in I$, there exists an $n \leq n_0$ and $z \in \Lambda^{(i)}$ such that $T^n z = x$.

$$\lambda^n \psi(x) = \mathcal{P}^n \psi(x) := \sum_{T^n y = x} \frac{\psi(y)}{|(T^n)'(y)|} \geq \frac{\psi(z)}{|(T^n)'(z)|}$$

Let η denote $\max_{[0,1]} |\hat{T}'|$. Then

$$\inf_I \psi \geq \frac{\rho}{2\delta \eta^{n_0}}.$$

Thus ν is proportional to Lebesgue measure on I .

Since $\varphi \in X_M$, we have $|\mathcal{P}\varphi|_1 = \lambda \geq 1 - qM$ from the estimates in Section 2.2.3. Recall that $M = \frac{b}{1-a}$ where a and b depend only on the map \hat{T} and the length scale δ . Similarly, $q = \sum_{\ell \geq 1} e^{\xi(\ell-1)} m \tilde{H}_\ell \leq \frac{mH}{\delta^2(\mu - \sqrt{2\mu})}$. From this we see that $\lambda \geq 1 - C_0 mH$, which is equivalent to the Lipschitz bound claimed in the remark after Theorem 3.2.

Estimating the variation of φ . If $\alpha = 1$, we can use the bound on $\|\varphi\|_r$ to estimate the variation of φ since on Δ_l , $|\varphi'| \leq \|\varphi\|_r \varphi e^{\xi l}$ by definition of $\|\cdot\|_r$. In fact, the following calculation shows that the projection of any function in X has bounded variation on I .

Let $\bigvee_J f$ denote the variation of f on the interval J . We estimate $\bigvee_I \psi$ using the following identity

$$\psi(x) = \sum_{i,l,j} \frac{\varphi(\pi_{l,j}^{(i)-1}x)}{\pi'(\pi_{l,j}^{(i)-1}x)}$$

where $\pi_{l,j}^{(i)-1}$ denotes the inverse of $\pi|_{\Delta_{l,j}^{(i)}}$. We estimate

$$\bigvee_I \psi \leq \sum_{i,l,j} \bigvee_{\pi(\Delta_{l,j}^{(i)})} \varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})' + 2|\varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})'|_\infty.$$

The second term can be estimated using equation (18) and the fact that $\varphi \in X_M$,

$$|\varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})'|_\infty \leq Me^{\xi l} \cdot \frac{1}{\delta\mu^l}. \quad (20)$$

To estimate the first term, we note that equation (18) and the distortion bounds of Lemma 3.1 imply that

$$\left| \frac{(\pi_{l,j}^{(i)-1})'(x)}{(\pi_{l,j}^{(i)-1})'(y)} - 1 \right| \leq \tilde{C}|x - y|$$

which in turn implies $\left| \frac{(\pi_{l,j}^{(i)-1})''}{(\pi_{l,j}^{(i)-1})'} \right| \leq \tilde{C}$. We use this to estimate

$$\begin{aligned} \bigvee_{\pi(\Delta_{l,j}^{(i)})} \varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})' &\leq \int_{\pi(\Delta_{l,j}^{(i)})} |(\varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})')'| dm \\ &\leq \int_{\pi(\Delta_{l,j}^{(i)})} |\varphi' \circ \pi_{l,j}^{(i)-1} \cdot ((\pi_{l,j}^{(i)-1})')^2| dm \\ &\quad + \int_{\pi(\Delta_{l,j}^{(i)})} \varphi \circ \pi_{l,j}^{(i)-1} |(\pi_{l,j}^{(i)-1})''| dm \\ &\leq \frac{1}{\delta\mu^l} \int_{\Delta_{l,j}^{(i)}} |\varphi'| dm + \tilde{C} \int_{\Delta_{l,j}^{(i)}} \varphi dm \\ &\leq \frac{Me^{\xi l}}{\delta\mu^l} \int_{\Delta_{l,j}^{(i)}} \varphi dm + \tilde{C} \int_{\Delta_{l,j}^{(i)}} \varphi dm. \end{aligned} \quad (21)$$

Putting together equations (20) and (21) we obtain

$$\begin{aligned} \bigvee_I \psi &\leq \sum_{i,l,j} \frac{Me^{\xi l}}{\delta\mu^l} \int_{\Delta_{l,j}^{(i)}} \varphi dm + \tilde{C} \int_{\Delta_{l,j}^{(i)}} \varphi dm + 2\frac{Me^{\xi l}}{\delta\mu^l} \\ &\leq \tilde{C} + \sum_{i,l} \frac{3Me^{\xi l}}{\delta\mu^l} 2^{l-1} \end{aligned}$$

$$\leq \tilde{C} + \frac{3NM}{2\delta} \sum_l \left(\frac{2}{\mu}\right)^{\frac{l}{2}} \quad (22)$$

where we have used $e^\xi \leq \sqrt{\frac{e}{2}}$ in the last line.

3.3 Proof of Theorem 3.3

We prove under the conditions stated in Theorem 3.3 that the conditionally invariant measures found via the tower construction will converge to the unique absolutely continuous invariant measure of the expanding map with no holes as $mH \rightarrow 0$. Our strategy will be to prove that the sequence of conditionally invariant densities generated by the sequence of holes is uniformly bounded above and below away from zero. The measures converge weakly to a density which satisfies the same bounds. This is an invariant measure for the expanding map with no holes and since it is equivalent to Lebesgue measure on $[0,1]$, it must be unique due to the transitivity property of \hat{T} (equation (16)). If $\alpha = 1$, we prove L^1 convergence of the densities as well.

Let \hat{T} and $\{H_s\}$ be as in the statement of Theorem 3.3. Let I_j^t be the intervals of monotonicity created by H_t and let $I^t = [0, 1] \setminus H_t$.

There exists a conditionally invariant measure φ_t on the tower Δ constructed as in Section 3.2.2 with eigenvalue λ_t (we suppress the dependence of Δ on H_t in our notation). This in turn induces a conditionally invariant density ψ_t for $T_t := \hat{T}|_{I^t} \cap \hat{T}^{-1}I^t$. Let δ from the proof be fixed and note that $N \leq \frac{1}{\delta}$ where N represents the number of $\Lambda^{(i)}$ needed to cover I^t . The constant δ does not change as the hole shrinks since the intervals of monotonicity of the map with holes can only grow and no new ones are created.

We essentially need to repeat the arguments given in Section 3.2.5, but taking care to show that the upper and lower bounds are uniform in s . This is trivial in the case of the upper bound since δ is fixed and $N \leq \frac{1}{\delta}$. Also $\sup_{\Delta_0} \varphi_s \leq M$ where $M = \frac{1+C}{1-a}$. All the constants here are independent of s . This also holds for the estimate on $\bigvee_I \varphi_s$.

We need to do the lower bound more carefully. We have

$$1 = \sum_{l=0}^{\infty} \int_{\Delta_l} \varphi_t dm \leq \sup_{\Delta_0} \varphi_t \sum_{l=0}^{\infty} \frac{1}{\lambda_t^l} m \Delta_l \leq \sup_{\Delta_0} \varphi_t \sum_{l=0}^{\infty} \frac{1}{\lambda_t^l} \frac{ND}{\delta} \left(\frac{2}{\mu}\right)^l.$$

Since $\lambda_t \geq \sqrt{\frac{2}{\mu}}$ and $D < 1$, this yields the inequality:

$$\sup_{\Delta_0} \varphi_t \geq \delta^2 \left(1 - \sqrt{\frac{2}{\mu}}\right) =: \kappa$$

So there exists a $\Delta_0^{(i)}$ such that the maximum of φ_t on that interval is at least

κ . The regularity of φ_t then forces $\inf_{\Delta_0^{(i)}} \varphi_t \geq \frac{\kappa}{1+M}$. This in turn implies that

$$\inf_{\Lambda^{(i)}} \psi_t \geq \frac{\kappa}{1+M}.$$

Now since $\Lambda^{(i)}$ grows (exponentially fast) to cover one of the intervals I_j^t , and each of these grows in at most m_j iterates to cover all of $[0,1]$, then there exists an m_0 such that $\Lambda^{(i)}$ grows to cover all of $[0,1]$ in at most m_0 iterates. So if \mathcal{P}_t is the Perron-Frobenius operator associated with T_t , then for any $x \in I^t$, there exists an $n \leq m_0$ and $z \in \Lambda^{(i)}$ such that $T_t^n z = x$.

$$\lambda_t^n \psi_t(x) = \mathcal{P}_t^n \psi_t(x) := \sum_{T_t^n y = x} \frac{\psi_t(y)}{|(T_t^n)'(y)|} \geq \frac{\psi_t(z)}{|(T_t^n)'(z)|}$$

Then

$$\inf_{I^t} \psi_t \geq \frac{\kappa}{(1+M)\eta^{m_0}}. \quad (23)$$

We are now ready to shrink the hole. Note that once the constants κ and M have been chosen for t , then they will work for each s as well. Also, m_0 can be chosen large enough to work for all $s \in [0, t]$ since the intervals I_j^s can only be larger than the intervals I_j^t . So we have the same lower bound from (23) for all the conditionally invariant densities ψ_s :

$$\inf_{I^s} \psi_s \geq \frac{\kappa}{(1+M)\eta^{m_0}}.$$

Now let ν_s be the measures conditionally invariant with respect to T_s having densities ψ_s and eigenvalues λ_s . Note also that by the proof of the existence of the conditionally invariant measures on the tower, we have $\lambda_s \rightarrow 1$ as $s \rightarrow 0$. Now take any sequence $\{s_i\}$ with $s_i \rightarrow 0$. The sequence of measures $\{\nu_{s_i}\}$ is precompact and so we may choose a subsequence $\{\nu_{s_k}\}$ such that the ν_{s_k} converge weakly to a probability measure ν on $[0,1]$. Now for any Borel subset A of $[0,1]$, we have

$$\frac{\kappa}{(1+M)\eta^{m_0}} m(A) \leq \nu(A) \leq \frac{M}{\kappa} m(A)$$

so that ν is equivalent to Lebesgue measure on $[0,1]$. Also

$$\begin{aligned} \nu(\hat{T}^{-1}A) &= \lim_{k \rightarrow \infty} \nu_{s_k}(\hat{T}^{-1}A) &= \lim_{k \rightarrow \infty} \nu_{s_k}(T_{s_k}^{-1}(A \setminus H_{s_k})) \\ &= \lim_{k \rightarrow \infty} \lambda_{s_k} \nu_{s_k}(A \setminus H_{s_k}) &= \lim_{k \rightarrow \infty} \lambda_{s_k} \nu_{s_k}(A) \\ &= \nu(A) \end{aligned}$$

so that ν is an invariant measure for \hat{T} . These facts, combined with the mixing property of \hat{T} , makes ν the unique absolutely continuous invariant measure for \hat{T} . Since ν is unique, the above argument forces the convergence of all the measures ν_s to ν as $mH_s \rightarrow 0$.

If $\alpha = 1$, we use a similar argument on the densities ψ_s since we have uniform bounds on their variations given by equation (22). We extend ψ_s to $L^1(\hat{I})$ by simply setting $\psi_s = 0$ on H_s . Since we have uniform upper bounds on ψ_s and the number of components of the hole does not increase as $s \rightarrow 0$, the variation of ψ_s will still be finite and uniformly bounded in s . Call this bound on the variations B .

Since the ψ_s lie in a compact subset of $L^1(\hat{I})$, we can choose a subsequence $\{\psi_{s_k}\}$ which converges in L^1 to a function ψ whose variation is at most B . ψ also has the same lower bound as the ψ_s . It only remains to show that ψ is an invariant density for \hat{T} .

Let $I_s = \hat{I} \setminus H_s$ and let $I_s^1 = I_s \cap T_s^{-1}I_s$. We have the relation $\mathcal{P}_s \psi_s = \lambda_s \psi_s$ for each s where \mathcal{P}_s is the Perron-Frobenius operator corresponding to T_s . Let $\hat{\mathcal{P}}$ be the Perron-Frobenius operator corresponding to \hat{T} and note that

$$\mathcal{P}_s f = \hat{\mathcal{P}}(f \cdot 1_{I_s^1}).$$

Noting that $|\psi_s|_\infty \leq 2 + B$, we are ready to estimate,

$$\begin{aligned} \int_{\hat{I}} |\hat{\mathcal{P}}\psi - \psi| &\leq \int_{\hat{I}} |\hat{\mathcal{P}}\psi - \hat{\mathcal{P}}\psi_s| + |\hat{\mathcal{P}}\psi_s - \psi_s| + |\psi_s - \psi| \, dm \\ &\leq 2|\psi_s - \psi|_1 + |\hat{\mathcal{P}}\psi_s - \hat{\mathcal{P}}(\psi_s \cdot 1_{I_s^1})|_1 + |\hat{\mathcal{P}}(\psi_s \cdot 1_{I_s^1}) - \psi_s|_1 \\ &\leq 2|\psi_s - \psi|_1 + |\psi_s \cdot 1_{\hat{I} \setminus I_s^1}|_1 + |\mathcal{P}_s \psi_s - \psi_s|_1 \\ &\leq 2|\psi_s - \psi|_1 + (2 + B)m(\hat{I} \setminus I_s^1) + (1 - \lambda_s). \end{aligned}$$

This last line approaches zero for $s = s_k$ as $k \rightarrow \infty$. We conclude that $\hat{\mathcal{P}}\psi = \psi$ so that ψ is the density for the unique SRB measure for \hat{T} . Since ψ is unique, the above argument forces the convergence in L^1 of all the densities ψ_s to ψ as $mH_s \rightarrow 0$.

Acknowledgments. The author is grateful to L.-S. Young for many helpful discussions and guidance during the writing of this paper.

References

- [BC] H. van den Bedem and N. Chernov, *Expanding maps of an interval with holes*, Ergod. Th. and Dynam. Sys. **22** (2002), 637-654.
- [BK] V. Baladi and G. Keller, *Zeta functions and transfer operators for piecewise monotonic transformations*, Comm. Math. Phys. **127** (1990), 459-477.
- [BY] M. Benedicks and L.-S. Young, *Markov extensions and decay of correlations for certain Hénon maps*, Asterisque **261** (2000), xi, 13-56.

- [C1] N. N. Čencova, *A natural invariant measure on Smale's horseshoe*, Soviet Math. Dokl. **23** (1981), 87-91.
- [C2] N. N. Čencova, *Statistical properties of smooth Smale horseshoes*, in Mathematical Problems of Statistical Mechanics and Dynamics, R. L. Dobrushin, ed. Reidel: Dordrecht, 1986, pp. 199-256.
- [Ch1] N. Chernov, *Statistical properties of piecewise smooth hyperbolic systems in high dimensions*, Discr. Cont. Dynam. Syst. **5** (1999), 425-448.
- [Ch2] N. Chernov, *Decay of correlations and dispersing billiards*, J. Stat. Phys. **94** (1999), 513-556.
- [Ch3] N. Chernov, *Sinai billiards under small external forces*, Annales Henri Poincaré **2** (2001), 197-236.
- [CM1] N. Chernov and R. Markarian, *Ergodic properties of Anosov maps with rectangular holes*, Bol. Soc. Bras. Mat. **28** (1997), 271-314.
- [CM2] N. Chernov and R. Markarian, *Anosov maps with rectangular holes. Nonergodic cases*, Bol. Soc. Bras. Mat. **28** (1997), 315-342.
- [CMS1] P. Collet, S. Martínez and B. Schmitt, *The Yorke-Pianigiani measure and the asymptotic law on the limit Cantor set of expanding systems*, Nonlinearity **7** (1994), 1437-1443.
- [CMS2] P. Collet, S. Martínez and B. Schmitt, *Quasi-stationary distribution and Gibbs measure of expanding systems*, in Instabilities and Nonequilibrium Structures V, E. Tirapegui and W. Zeller, eds. Kluwer: Dordrecht, 1996, pp. 205-19.
- [CMT1] N. Chernov, R. Markarian and S. Troubetskoy, *Conditionally invariant measures for Anosov maps with small holes*, Ergod. Th. and Dynam. Sys. **18** (1998), 1049-1073.
- [CMT2] N. Chernov, R. Markarian and S. Troubetskoy, *Invariant measures for Anosov maps with small holes*, Ergod. Th. and Dynam. Sys. **20** (2000), 1007-1044.
- [D] M. Demers, *Markov Extensions and Conditionally Invariant Measures for Certain Logistic Maps with Small Holes*, to appear in Ergod. Th. and Dynam. Sys.
- [KL] G. Keller and C. Liverani, *Stability of the spectrum for transfer operators*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **28** (1999), 141-152.
- [LiM] C. Liverani and V. Maume-Deschamps, *Lasota-Yorke maps with holes: conditionally invariant probability measures and invariant probability measures on the survivor set*, Annales de l'Institut Henri Poincaré Probability and Statistics, **39** (2003), 385-412.

- [LM] A. Lopes and R. Markarian, *Open Billiards: Cantor sets, invariant and conditionally invariant probabilities*, SIAM J. Appl. Math. **56** (1996), 651-680.
- [PY] G. Pianigiani and J. Yorke, *Expanding maps on sets which are almost invariant: decay and chaos*, Trans. Amer. Math. Soc. **252** (1979), 351-366.
- [R] P.A. Richardson, Jr., *Natural Measures on the Unstable and Invariant Manifolds of Open Billiard Dynamical Systems*, Doctoral Dissertation, Department of Mathematics, University of North Texas, 1999.
- [Y1] L.-S. Young, *Statistical properties of dynamical systems with some hyperbolicity*, Ann. of Math. **147** (1998), 585-650.
- [Y2] L.-S. Young, *Recurrence times and rates of mixing*, Israel J. of Math. **110** (1999), 153-188.