

Original Research Article

Prevalence and associated factors of oral Non-Candida albicans Candida carriage in Denture wearers in Sana'a City-Yemen.**Abstract**

The objective of this study was to contrast the prevalence and species of colonization of Non-*Candida albicans Candida* (NCAC) in the oral cavity of denture wearers and non-denture wearers and assess associated risk factors of their colonization.

A total of 208 subjects were studied: 104 denture wearers and 104 non-denture wearers, matched by age and sex, comprised the experimental and control groups, respectively.

Each subject was instructed to perform oral rinsing using a phosphate-buffered saline solution, which was expectorated and processed for the recovery of *Candida* species on Sabouraud dextrose agar. Isolates were identified by culturing on chromogenic *Candida* agar and noting species-specific colony characteristics.

There was a significant oral carriage rate of NCAC among denture wearers (5.83% versus 11.1% in controls) with associated risk factor (5.4) (PV<0.001). The most common isolated NCAC were *C.Krusi* and *C.tropicalis* with significant OR (5.5 and 4.7 respectively). When co-infections were considered, there was highly significant association of *C. albicans* + *C. krusi* oral colonization in cases (OR=4.56, PV<0.001). There was a significant oral carriage rate of NCAC among male denture wearers (36.9%, OR=6.6, PV<0.001). In addition, there was a significant rate of NCAC colonization with complete denture (rate= 50%, OR=2.4, PV= 0.02). While no significant increase associated with colonization of NCAC with partial, acrylic and/or chrome cobalt denture.

Conclusion

Based on the results of this study ability of NCAC were greater in denture wearers than non-denture wearers, also greater risk of NCAC were found with males, older ages, and complete denture.

Key words: Non-*Candida albicans Candida* (NCAC); carriage; denture; prevalence

Introduction

The preponderance oral yeast infections are caused by members of the genus *Candida*. Candidiasis is an opportunistic infection due to pathological changes in the surface of oral mucosal cavity.^{1, 2} Candidiasis patients may show various symptoms including painful sensation, burning, swallowing difficulty and change of taste, but most frequently are asymptomatic.³ The infection is generally treated with antifungal drugs, but in immune-compromised patients such as patients treated in intensive care units, cancer patients receiving radiation or chemotherapy, organ transplant patients and HIV-positive patients' return of infections may be problem. In the last twenty years, some *Candida species*, including *C. glabrata*, *C.tropicalis*, *C. krusei*, and *C. parapsilosis*, have been isolated with increasing frequency from cases of candidiasis.^{4,5,6} It is known that each species differs in the production of recognized virulence factors and sensitivity to antifungal agents. Isolation and identification the *Candida* spp. is useful in choosing correct treatment, as some species may be resistant to certain groups of antifungal drugs.⁷⁻⁹ Infection caused by NCAC, such as *C.glabrate*, *C.tropicalis*, and *C. krusei*, have been found to be less response to fluconazole.¹⁰ Moreover, there are reports of numerous cases

describing infection and colonization of immune-compromised patients on long-term treatments of oral antifungal agents, and drug resistant *C. glabrata* and *C. krusei* have been recovered from them.¹¹ As well as, host defenses have been reported to be less effective in patients infected by *C. glabrata* than *C. albicans*.¹¹ The newer triazoles, including posaconazole and voriconazole; and echinocandins, caspofungin, micafungin, and anidulafungin are antifungal drugs that show strong activity against *Candida* spp. On the other hand, echinocandins, appears to be less potent against some species, such as *C. guilliermondii* and *C. parapsilosis*.⁸ As well, *C. dubliniensis* is very similar to *C. albicans* has been reported to have low susceptibility to azole drugs.^{9,12}

The wearing of dentures are associated with overgrowth of *Candida* species, leading to denture stomatitis. Studies to identify *Candida* species in patients with denture stomatitis have produced contradictory results.⁸ A number of studies assertion that a single species was responsible for the infections¹³ while others isolated and accused multiple species of *Candida*.¹⁴ The objective of this study was to compare the prevalence and species colonization due to NCAC in the oral cavity of denture wearers and non-denture wearers and analysis associated risk factors of their colonization.

Subjects and Laboratory Methods

Subject selection

A total of two hundreds and eight persons, were included in this study, 104 of them were denture wearer patients (cases group) while the other 104 adults with natural teeth (controls group), whom been selected randomly from Al-Thawrah hospital, Al-Gumhory hospital and Dental centers in Sana'a city, Yemen. The duration of the study was six months period, started in August 2017 and ended in February 2018. Inclusion criteria for subject selection were healthy individuals with no clinical signs of *Candida* infection and no systemic disease. In addition, individuals who smoked, currently taking antifungal, steroids, antibiotics, or immunosuppressive drugs in the past 6 months were excluded.

Collection and identification of samples

Salivary samples were collected using the oral rinse technique.¹⁵ In brief, each subject was asked to rinse the mouth for 60 seconds with 10 ml of sterile phosphate-buffered saline (PBS; 0.01 M phosphate-buffered saline solution, pH 7.2) and expectorate the wash into a 15 ml sterile container.¹⁶ Individuals who had removable dentures were asked to take out the denture prior to samples collection. The samples were immediately transported on ice to the microbiology laboratory. Each oral rinse was centrifuged at 3500 rpm for 10 minutes, and then the supernatant was discarded. The pellet was re-suspended in 1ml sterile PBS. One hundred µl of the concentrated oral rinse was inoculated onto Sabouraudus dextrose agar and incubated at 37 °C for 48 hours. The lasting samples were stored at -20 °C. If *Candida* colonies appeared on the Sabouraud dextrose agar, then chromogenic *candida* agar was inoculated using 100 µl of the oral rinse supernatant and incubated for 48 hours for colonies study. *Candida* species were identified by the color of the colonies using the color reference guide supplied by the manufacturer. When color identification was unclear, fermentation assay of sucrose, maltose, glucose, lactose and galactose was done. The *Candida* species were also identified by the ability to produce chlamydo-spores on glutinous rice agar.¹⁷

Data analysis

Data were statistically analyzed using the EPI-Info program version 6. The difference in distribution of the *Candida species* between groups was based on comparison of frequency distributions by a chi-square test. A p value < 0.05 was considered significant.

Ethical approval

We obtained written consent from all cases. Assent was taken from participants before collecting the specimens. The study proposal was evaluated and approved by the Ethics Committee of Faculty of Medicine and Health Sciences, Sana'a University.

Results

There was a significant oral carriage rate of NCAC among denture wearers (cases) equaled to 5.83% comparing with only 11.1% among non-denture wearers (controls) with OR of association equaled to 5.4 times (PV<0.001). There was a significant oral carriage rate of *C. Krusei* among cases (30.7%,) comparing with only 7.4% in controls with 5.5 association (PV<0.001). A significant oral carriage rate of *C. tropicalis* among cases (15.4%,) comparing with only 3.7% in controls with 4.7 times of association (PV=0.003), while no significant association with *C. glabrata*. When co-infections were considered, there was highly significant association of *C. albicans* + *C. krusi* oral colonization in cases (OR=4.56, PV<0.001), but no significant association of *C. albicans* + *C. tropicalis* infection in the denture wearer (Table 1). There was a significant oral carriage rate of NCAC species among male denture wearers (36.9%) with significant association (OR=6.6, PV<0.001). While in the females group, there was only a slightly difference between oral carriage rate of NCAC among female denture wearers comparing with non-denture wearer females. There was a highly significant association between older age groups (>45 years) of cases and a contract of oral NCAC colonization with OR ranged from 1.4 to 5.4 times (table 2). When we considered type of dental prosthesis, there was a significant rate increase with complete denture. The rate was 50%, with a highly significant association between complete denture and a contract of oral NCAC colonization (OR=2.4, PV= 0.02) . While no significant increase associated with colonization of NCAC with partial, acrylic and/or chrome cobalt denture (table 3).

Discussion

NCAC strains, however, are isolated in growing numbers in medically compromised patients. These strains might cause systemic infections and are frequently resistant to commonly used antifungal agents such as fluconazole.^{18,19} In the present study, there was a significant oral carriage rate of NCAC among denture wearers (38.5%) comparing with only 11.1% among non-denture wearers. In addition, there was a highly significant association between denture wear and a contract of oral NCAC with OR equaled to 5.4 times (PV<0.001). When single species of NCAC was considered, there was a significant oral carriage rate of *C. Krusei* among cases (30.7%,) comparing with only 7.4% in controls with 5.5 association (PV<0.001) followed by *C. tropicalis* (15.4%) comparing with only 3.7% in controls with 4.7 times of association (PV=0.003), while no significant association with *C. glabrata* and absent of *C. parapsilosis* (Table 1). Our results were different from that reported elsewhere, in which *C. tropicalis* was the most common NCAC, followed by *C. glabrata*, while *C. parapsilosis* were rare isolated in healthy oral colonization or as a cause of illnesses in patients.¹⁹

When co-infections were considered, there was highly significant association of *C. albicans* + *C. krusi* oral colonization in cases (OR=4.56, PV<0.001), but no significant association of *C. albicans* + *C. tropicalis* infection in the denture wearer (Table 1). The

carriage rates of single and multiple *Candida* species were reported to be significantly higher in denture wearers.²⁰ As it is known, NCAC may be capable of metabolizing ethanol to carcinogenic acetaldehyde and can thus progress oral and upper gastrointestinal tract cancer. So, 38.5% of our studied individuals having dentures under possible risk of oral and upper gastrointestinal tract cancer. Consequently, more focus should be placed on diagnosis and treatment of oral *Candida* infections, also on other *Candida* species than *C. albicans* as it has been recommended.¹⁹⁻²²

The highly significant association between denture wear and a contract of oral NCAC with the high risk (OR equaled to 5.4 times) can be explained by the fact that one of the most important virulence factors of NCAC is its ability to form biofilms, which has an important clinical consequence, as it confers resistance to antifungal therapy and capacity for yeast cells within the biofilms to resist host immune defenses.²³⁻²⁶ A second explanation might be due to that changes in the oral environment affected by tooth loss or denture wearing can cause changes in oral microflora.²⁰

Also in this study, the data supported the rejection of the null hypothesis, which states that there would be no difference between male and female denture wearers in terms of the colonization by NCAC of the inner surfaces of dentures and the surrounding attachments. While there was a significant association between denture wearing in male group and a contract of oral carriage of NCAC with OR equaled to 6.6 times, (CI=2.7 -16.5, PV<0.001). Our result was different from that reported by Their²⁷ in which similar colonization rate was observed in both adult sexes.

Moreover, there was a trend of rising of oral NCAC colonization with increasing age in cases and controls (table 1), in which the highest rate of NCAC was in the older age groups. Our result, also, was similar to that reported by Chopde *et al.*²⁸ in which extremes of age is more susceptible to oral NCAC colonization. This result can be explained by that the elderly generally suffered from systemic illnesses, changes in nourishment and their salivary characteristics.^{28, 29} Advancing age was also a risk factor for denture stomatitis in the elderly, because cell-mediated immunity, which provides protection against *candidal* infection, declines with age. According to Dar-odeh and Shehabi³⁰, some oral environmental factors, such as un-stimulated salivary flow rate and age of subjects, were associated with higher numbers of microbes in the saliva of complete denture wearers. Those authors suggested that a reduction in the salivary flow rate with aging induced an increase in concentration of microbes in saliva.³¹

Conclusion

Based on the results of this study ability of NCAC were greater in denture wearers than non-denture wearers, also greater risk of NCAC were found with males, older ages, and complete denture. In addition, our results are important for the development of strategies to for eliminate these indicators of risk and significantly reduce NCAC colonization and oral *Candida* infections in denture wearers.

Acknowledgments:

Authors acknowledge the financial support of Sana'a University, Yemen.

Conflict of interest:

"No conflict of interest associated with this work".

References

- 1-Gonsalves WC, Chi AC, Neville BW. Common oral lesions: Part I. Superficial mucosal lesions. *Am Fam Physician* 2007;75:501-7.
- 2-Da Costa SC, De Resende MA, Lyon JP, Totti VMG, Munhoz MF. Predisposing conditions for *Candida* spp. carriage in the oral cavity of denture wearers and individuals with natural teeth. *Can J Microbiol* 2006; 52:462-7.
- 3-Samaranayake LP. Host factors and oral candidiasis. In: MacFarlane TW, Samaranayake LP, editors. *Oral candidosis*. London: Butterworth & Company Ltd; 1990. p.66-103.
- 4-Samaranayake LP. Superficial oral fungal infections. *Curr Opin Dent*. 1991; 1:415-22.
- 5-Scully C, el-Kabir M, Samaranayake LP. *Candida* and oral candidosis: a review. *Crit Rev Oral Biol Med* 1994;5:125-57.
- 6-Borg-von ZM, Kunz L, Rüchel R, Reichard U, Weig M, Gross U. Epidemiology and antifungal susceptibilities of *Candida* spp. to six antifungal agents: results from a surveillance study on fungaemia in Germany from July 2004 to August 2005. *J Antimicrob Chemother*. 2007;60:424-8.
- 7-Kuriyama T, Williams DW, Bagg J, Coulter WA, Ready D, Lewis MA. In vitro susceptibility of oral *Candida* to seven antifungal agents. *Oral Microbiol Immunol*. 2005; 20:349-53.
- 8-Moudgal V, Little T, Boikov D, Vazquez JA. Multiechinocandin-and Multiazole-resistant *Candida parapsilosis* isolates serially obtained during therapy for prosthetic valve endocarditis. *Antimicrob Agents Chemother* 2005; 49:767-9.
- 9-Sullivan DJ, Moran GP, Pinjon E, Al-Mosaid A, Stokes C, Vaughan C, *et al*. Comparison of the epidemiology, drug resistance mechanisms, and virulence of *Candida dubliniensis* and *Candida albicans*. *FEMS Yeast Res*. 2004; 4:369-76.
- 10-Sheehan DJ, Hitchcock CA, Sibley CM. Current and emerging azole antifungal agents. *Clin Microbiol Rev* 1999;12:40-79.
- 11-Mendling W, Koldovsky U. Immunological investigations in vaginal mycoses. *Mycoses* 1996;39: 177-83.
- 12-Schubert S, Rogers PD, Morschhäuser J. Gain-of-function mutations in the transcription factor *MRR1* are responsible for over expression of the *MDR1* efflux pump in fluconazole-resistant *Candida dubliniensis* strains. *Antimicrob Agents Chemother* 2008;52:4274-80.
- 13-Kreher JM, Graser GN, Handelman SL, Eisenberg AD. Oral yeasts, mucosal health, and drug use in an elderly denture-wearing population. *Spec Care Dentist*. 1991;11:222-6.
- 14-Cumming CG, Wight C, Blackwell CL, Wray D. Denture stomatitis in the elderly. *Oral Microbiol Immunol*. 1990;5:82-5.
- 15-Coulter WA, Kinirons MJ, Murray SD. The use of a concentrated oral rinse culture technique to sample oral *Candida* and lactobacilli in children, and the relationship between *Candida* and Lactobacilli levels and dental caries experience: A pilot study. *Int J Paediatr Dent*. 1993;3:17-21.
- 16-MacFarlane TW, Samaranayake LP, Williamson MI. Comparison of Sabouraud dextrose and Pagano-Levin agar media for detection and isolation of yeasts from oral samples. *J Clin Microbiol* 1987;25:162-4.
- 17-Staib P, Morschhäuser J. Chlamyospore formation in *Candida albicans* and *Candida dubliniensis* – an enigmatic developmental programme. *Mycoses*. 2007; 50:1-12.
- 18-Lalla, RV; Patton, LL; Dongari-Bagtzoglou, A. "Oral candidiasis: pathogenesis, clinical presentation, diagnosis and treatment strategies." *J. of the California Dental Association* 2013; 41 (4): 263–8.
- 19-Kourkoumpetis, Themistoklis. "Candida infection and colonization among non-trauma emergency surgery patients." *Virulence* 2010; 359-366.

20-Jenkinson HF, Douglas LJ. Interactions between *Candida* species and bacteria in mixed infections. In: Brogden KA, Guthmiller JM, editors. Polymicrobial diseases. Washington DC: ASM Press 2002; p. 357-73.

21-Manolakaki, D., Velmahos, G., Kourkoumpetis, T., Chang, Y., Alam, H. B., De Moya, M. M., & Mylonakis, E. *Candida* infection and colonization among trauma patients. *Virulence* 2010; 1 (5):367-375.

22-dEnfert C; Hube B (editors). *Candida: Comparative and Functional Genomics*. Caister Academic Press. ISBN 978-1-904455-13-4 (2007).

23-Pereira-Cenci T, Del Bel Cury AA, Crielaard W, Ten Cate JM. Development of *Candida*-associated denture stomatitis: new insights. *J. Appl. Oral Sci* 2008; 16:86-94.

24-Coogan MM, Fidel PL Jr, Komesu MC, Maeda N, Samaranayake LP. *Candida* and mycotic Infections. *Adv Dent Res* 2006; 19:130-8.

25-Costa CR, Cohen AJ, Fernandes OF, Miranda KC, Passos XS, Souza LK, *et al.* Asymptomatic oral carriage of *Candida* species in HIV-infected patients in the highly active antiretroviral therapy era. *Rev. Inst. Med. Trop. Sao Paulo* 2006; 48:257-61.

26-Jin YY, Yip HK, Samaranayake YH, Yau JY, Samaranayake LP. Biofilm-forming ability of *Candida albicans* is unlikely to contribute to high levels of oral yeast carriage in cases of human immunodeficiency virus infection. *J. Clin. Microbiol* 2003; 41:2961-7.

27-Thein ZM, Samaranayake YH, Samaranayake LP. *In vitro* biofilm formation of *Candida albicans* and non-albicans *Candida* species under dynamic and anaerobic conditions. *Arch. Oral Biol* 2007; 52:761-7.

28-Chopde N, Jawale B, Pharande A, Chaudhari L, Hiremath V, Redasani R. Microbial colonization and their relation with potential cofactors in patients with denture stomatitis. *J Contemp Dent Pract* 2012; 13:456-9.

29-Semlali A, Killer K, Alanazi H, Chmielewski W, Rouabhia M. Cigarette smoke condensate increases *C. albicans* adhesion, growth, biofilm formation, and EAP1, HWP1 and SAP2 gene expression. *BMC Microbiol* 2014; 14:61.

30-Dar-odeh NS, and Shehabi AA. Oral candidiasis in patients with removable dentures *Mycoses* 2003; 46:187-19.

31-Penha SS, Birman EG, Silveira FR, Paula CR. Frequency and enzymatic activity (proteinase and phospholipase) of *Candida albicans* from edentulous patients, with and without denture stomatitis. *Pesq Odontol Bras* 2000; 14:119-122.

Table 1: The yeast distribution in different sexes of the denture wearer (cases) and non denture wearer (controls) groups of the study populations:

Organisms	Cases (No.=104)		controls (No.=104)		OR	CI	χ^2	PV
	No.	%	No.	%				
<i>Total Non C. albicans Candida</i>	42	38.5	12	11.5	5	2.4-10.2	35	<0.001
<i>C. krusi</i>	32	30.7	8	7.6	5.5	2.4-12.7	18.8	<0.001
<i>C. tropicalis</i>	16	15.4	4	3.8	4.7	1.5-14.6	8.4	0.003
<i>C. glabrata</i>	2	1.9	2	1.9	1.03	0.14-7.5	0.96	0.6
<i>C. albicans + C. krusi</i>	22	21.1	6	5.8	4.56	1.7-11.7	22	<0.001
<i>C. albicans + C. tropicalis</i>	8	7.7	4	3.8	2.16	0.6-7.4	1.9	0.16
<i>C. albicans + C. krusi + C. tropicalis</i>	6	5.7	0	0.0	undefined		6.5	0.012

OR Odds ratio = Relative risk

95%CI 95% Confidence intervals

χ^2 Chi-square = 3.9 or more is significant

pv Probability value = 0.05 or less is significant

Table 2: The carriage rate of NCAC in cases and controls and associated OR for different sexes and age groups :

Factors		cases (n=104)		controls (n=104)		OR	CI	χ^2	PV
		No.	%	No.	%				
Male		36/92	39.1	8/92	8.7	6.6	2.7 -16.5	23	<0.001
Female		6/12	50	4/12	33.3	1.6	0.4-6.9	0.5	0.47
Age group	< 45 years	2/10	20	2/10	20	1.4	0.1- 10	0.0	0.97
	45-54 years	8/16	50	2/16	12.5	4.4	1.0 -30	4	0.04
	55 – 64 years	16/32	50	4/32	12.5	4.7	1.4-17.4	8.5	0.003
	≥ 65 years	18/46	39.1	4/46	8.7	5.4	1.6-19.8	10.5	0.001
Total		42/104	40.4	12/104	11.5	5.4	2.5-11.8	23.9	<0.001

OR Odds ratio = Relative risk

95%CI 95% Confidence intervals

χ^2 Chi-square = 3.9 or more is significant

pv Probability value = 0.05 or less is significant

Table 3: The type of dental prosthesis factors that associated with colonization of NCAC among denture wearer patients

Factors	Positive NCAC (n=42)		OR	CI	χ^2	PV
	No.	%				
Type of dental prosthesis						
complete n=46	24	52.2	2.4	1.0.- 5.8	4.7	0.02
Partial n=58	18	31	0.4	0.16- 0.9	5.4	0.02
Acrylic n=74	30	40.5	1.02	0.4-2.6	0.00	0.97
Chrome cobalt n=38	12	31.6	0.73	0.3-1.8	0.54	0.46

OR Odds ratio = Relative risk

95%CI 95% Confidence intervals

χ^2 Chi-square = 3.9 or more is significant

pv Probability value = 0.05 or less is significant