

1-17-2007

Finite-Size Effects in a Two-Dimensional Electron Gas with Rashba Spin-Orbit Interaction

C P. Moca
University of Oradea

D C. Marinescu
Clemson University, dcm@clemson.edu

Follow this and additional works at: https://tigerprints.clemson.edu/physastro_pubs

Recommended Citation

Please use publisher's recommended citation.

This Article is brought to you for free and open access by the Physics and Astronomy at TigerPrints. It has been accepted for inclusion in Publications by an authorized administrator of TigerPrints. For more information, please contact kokeefe@clemson.edu.

Finite-size effects in a two-dimensional electron gas with Rashba spin-orbit interaction

C. P. Moca^{1,2} and D. C. Marinescu³¹*Department of Physics, University of Oradea, 410087 Oradea, Romania*²*Institute of Physics, Technical University Budapest, Budapest H-1521, Hungary*³*Department of Physics and Astronomy, Clemson University, Clemson, South Carolina 29634, USA*

(Received 7 August 2006; revised manuscript received 5 December 2006; published 17 January 2007)

Within the Kubo formalism, we estimate the spin-Hall conductivity in a two-dimensional electron gas with Rashba spin-orbit interaction and study its variation as a function of disorder strength and system size. The numerical algorithm employed in the calculation is based on the direct numerical integration of the time-dependent Schrödinger equation in a spin-dependent variant of the particle-source method. We find that the spin-precession length L_s , controlled by the strength of the Rashba coupling, establishes the critical length scale that marks the significant reduction of the spin-Hall conductivity in bulk systems. In contrast, the electron mean free path, inversely proportional to the strength of disorder, appears to have only a minor effect.

DOI: [10.1103/PhysRevB.75.035325](https://doi.org/10.1103/PhysRevB.75.035325)

PACS number(s): 72.10.-d, 72.20.-i, 72.90.+y

I. INTRODUCTION

The physical phenomenon behind the spin-Hall effect^{1,2} (SHE) in two-dimensional (2D) systems is the flow of a pure spin current, spin polarized in a transverse direction, driven by a perpendicular electric field. Its existence is conditioned by the presence of a spin-orbit interaction (SOI), such as Rashba-Dresselhaus,^{3,4} in n -type two-dimensional systems or the spin-split band structure in p -type GaAs.⁵

In clean samples, the spin-Hall conductivity σ_{sH} was predicted to have a universal, constant value of $e/8\pi$; however, in the presence of disorder the resulting picture was less clear. It was pointed out that in 2D infinite systems, in the presence of short range scatterers, the vertex corrections provided the exact compensation to cancel the effect.⁶ Moreover, an argument was made that this cancellation occurs even for infinitesimal disorder potentials.⁷ These conclusions were challenged by analytic⁸ and numerical calculations⁹⁻¹¹ of the spin-Hall conductivity in one- and two-dimensional finite-size mesoscopic samples, performed within the Landauer-Büttiker formalism, where it was shown that the effect survives up to a critical disorder strength.

Even though the robustness of the spin-Hall effect in the presence of disorder seems to have been definitively confirmed by the angle-resolved optical detection of spin polarization at opposite edges of a two-dimensional hole layer,¹² a better understanding of the mechanism by which disorder and system size affect spin transport in systems with spin-orbit interaction warrants further investigation. We focus, therefore, on a study of the interplay between the disorder strength, embodied in the electron mean free path l and the spin precession length L_s proportional to the spin-orbit interaction, in determining the spin-transport regime in finite-size samples. Such an analysis is especially relevant in two dimensions where, in the absence of any additional interactions, the two lengths are independent of each other and, along with the Fermi energy, are the only relevant physical parameters of the system.

The relationship between l and L_s and the system size L determines the existence of four distinctive transport regimes. A semiclassical approximation is appropriate for L_s

$\ll L$, when the spin coherence is lost over the length of the sample, while $L_s \gg L$ corresponds to a mesoscopic regime. When $l \gg L$ the electron propagation is ballistic, while for $l \ll L$ multiple scattering events are assumed and the diffusive regime is present.

In the following analysis, we use the Kubo formula to estimate σ_{sH} as a function of system size and disorder in a two-dimensional electron system. The numerical formalism adopted here represents an extension to the spin-Hall problem of the particle-source method developed by Tanaka and Itoh.¹³ This algorithm is based on the direct integration of the time-dependent Schrödinger equation and allows the calculation of the matrix elements of Green's functions, linear response functions, or any combination of Green's function and quantum operators in a very efficient way.

The main result of this study is that the delimitation between the mesoscopic and semiclassical regimes, as reflected by the rapid decline of the spin-Hall conductivity, is established by L_s . For system sizes smaller than L_s , the spin-Hall conductivity increases monotonically with the system size while being weakly affected by disorder. When $L \gg L_s$, σ_{sH} decreases exponentially for any amount of disorder in the system. This result supports the conclusions of two previous reports by Sheng *et al.*¹⁴ and Nomura *et al.*,¹⁵ where it was found that σ_{sH} remains finite up to an unspecified characteristic length scale and vanishes in the thermodynamic limit for any small amounts of disorder in the system. Here, we identify this length as being determined by the spin-precession length. Our results reflect no qualitative modification of the overall behavior when the system evolves from the ballistic to the diffusive regimes, crossover controlled by the mean free path characteristic length scale. For a fixed Fermi energy and L_s , the spin-Hall conductivity decreases monotonically with disorder for any system size, as the system evolves from diffusive to ballistic regime.

II. THEORETICAL FRAMEWORK

A. Model

The single-particle Hamiltonian that describes the dynamics of an electron of momentum \mathbf{p} and effective mass m^* is

written, in terms of the Pauli matrices $\sigma_{x,y}$ and the Rashba coupling constant λ , as

$$\tilde{H} = \frac{\mathbf{p}^2}{2m^*} + \lambda(\sigma_x p_y - \sigma_y p_x). \quad (1)$$

The exact diagonalization procedure that can be performed on the Hamiltonian in the case of a clean system¹ becomes impossible when disorder is included in the form of an additional random scattering term. It is, therefore, more convenient for a numerical analysis to adopt the tight-binding approximation for the many-body Hamiltonian by employing a local-orbital basis associated with a virtual square $N \times N$ lattice of constant a_0 . In this model, the many-body Hamiltonian is

$$H = \sum_{i,\alpha} \varepsilon_i c_{i\alpha}^\dagger c_{i\alpha} - t \sum_{\langle i,j \rangle, \alpha} c_{i\alpha}^\dagger c_{j\alpha} + V_R \sum_{i,\delta_x,\delta_y} [(c_{i\uparrow}^\dagger c_{i+\delta_x,\downarrow} - c_{i\downarrow}^\dagger c_{i+\delta_x,\uparrow}) - i(c_{i\uparrow}^\dagger c_{i+\delta_y,\downarrow} + c_{i\downarrow}^\dagger c_{i+\delta_y,\uparrow})]. \quad (2)$$

In this expression, an electron with spin α at site i , created by $c_{i\alpha}^\dagger$, is subjected to a random on-site energy as in the Anderson model for disorder, generated by a box distribution $\varepsilon_i \in [-W/2, W/2]$. The electron transport is described by a sequence of discrete hopping events. Lateral transport, without spin flip, to an adjacent site occurs with probability $t = \hbar^2/2m^*a_0$, taken to be the unit of energy in our calculation, as described by the second term in Eq. (2). Propagation along the diagonal sites, driven by the spin-orbit interaction, occurs with a simultaneous spin flip, as in the last term of Eq. (2). The latter is the most important as it mixes the spin channels and leads to a finite spin-Hall conductivity and spin accumulations at the edges of sample. The Rashba coupling constant is renormalized by the lattice constant to $V_R = \hbar\lambda/a_0$.

The Kubo formula for the spin-Hall conductivity is written as

$$\sigma_{sH} = \frac{1}{2} \text{Tr} \int \frac{d\varepsilon}{2\pi} \left(-\frac{\partial f(\varepsilon)}{\partial \varepsilon} \right) \langle j_x^z [G_R(\varepsilon) - G_A(\varepsilon)] \times v_y G_A(\varepsilon) - j_x^z G_R(\varepsilon) v_y [G_R(\varepsilon) - G_A(\varepsilon)] \rangle. \quad (3)$$

The velocity operator is defined by the commutator: $i\hbar v_y = [y, H]$, while for the spin current we adopt a traditional expression,¹⁶ given in terms of the anticommutator between the velocity operator and the Pauli matrix σ_z : $j_x^z = \hbar\{\sigma_z, v_x\}/4$. $G_{R/A}(\varepsilon)$ represents the retarded/advanced Green's function. In Eq. (3), the integration over the energy is restricted over the Fermi surface due to the presence of $[-\partial f(\varepsilon)/\partial \varepsilon]$ factor.

In the tight-binding framework, the effect of disorder and spin-orbit interaction strength on the spin-Hall conductance was investigated previously, using the Landauer-Büttiker formalism.^{17,18} As will be discussed in the next section, in the present work we use a different approach in computing Green's function needed for the calculation of the spin-Hall conductivity.

B. Numerical algorithm

Since the purpose of this investigation is an analysis of the spin-Hall conductivity dependence on system size and disorder, we will apply the Kubo formula to large size systems for different values of the disorder potential W . The numerical algorithm that underlies this calculation has been introduced in Ref. 13 and represents an extension of the particle-source method combined with tight-binding formalism. This method was first applied to the calculation of Green's function, density of states, conductivity,¹⁹ and Hall conductivity.¹³ The main advantage is that one can evaluate both the diagonal and off-diagonal parts of Green's function and their products with other quantum operators with low computing effort. In principle, the computing effort for computing Green's function is $O(N^3)$ (Hamiltonian is expressed as an $N \times N$ matrix), while within the present algorithm only $O(N)$ computational effort is required for the same calculation. Here, we briefly outline the main features of the algorithm.

The central part of the method consists in solving the time-dependent Schrödinger equation with a single-frequency source term,

$$i\hbar \frac{d|\tilde{j}, t\rangle}{dt} = H|\tilde{j}, t\rangle + |j\rangle \theta(t) \exp^{-i(E+i\eta)t}, \quad (4)$$

where η is a finite small value and θ is the step function. The solution of the equation, with the initial condition $|\tilde{j}, t=0\rangle = 0$, becomes

$$|\tilde{j}, t\rangle = -i \int_0^t dt' e^{-iH(t-t')} |j\rangle e^{-i(E+i\eta)t'} = \frac{1}{E+i\eta-H} [e^{-i(E+i\eta)t} - e^{-iHt} |j\rangle]. \quad (5)$$

For sufficiently large amount of time, one can then write the solution to the Schrödinger equation in terms of Green's function acting on the "source" $|j\rangle$, with the relative accuracy $\delta = e^{-\eta T}$, as

$$|\tilde{j}, T\rangle = G(E+i\eta) |j\rangle e^{-i(E+i\eta)T}, \quad (6)$$

leading to Green's function operating on the ket $|j\rangle$,

$$G(E+i\eta) |j\rangle = \lim_{T \rightarrow \infty} |\tilde{j}, T\rangle e^{i(E+i\eta)T}. \quad (7)$$

The matrix element between states $\langle i|$ and $|j\rangle$ is then obtained as

$$\langle i|G(E+i\eta)|j\rangle = \lim_{T \rightarrow \infty} \langle i|\tilde{j}, T\rangle. \quad (8)$$

The matrix elements of a product including several Green's functions and other operators are obtained by choosing a new initial state, such as $|j'\rangle = AG(E+i\eta)|j\rangle$ in Eq. (4), and repeating the same procedure.

To calculate the matrix elements of Green's function at many different energy values, one solves Eq. (4) simultaneously for a source term with multiple frequencies, $|j\rangle (\sum_l e^{-i(E_l+i\eta)l}) \theta(t)$. Following the algorithm outlined above, one obtains as an approximate solution the ket,




FIG. 1. Spin-Hall conductivity as a function of Fermi energy in units of $e/8\pi$ for clean and disordered systems. System size is 30×30 . Spin-orbit interaction strength is fixed to $V_R=0.2t$. Average was done over 2000 random frequencies.

$$\langle \tilde{j}, T \rangle \simeq \sum_l G(E_l + i\eta) |j\rangle e^{-i(E_l + i\eta)T}. \quad (9)$$

The matrix element of Green's function between the states $\langle i|$ and $|j\rangle$ for a given energy is then easily obtained as

$$G_{ij}(E_l' + i\eta) = \langle i|G(E_l' + i\eta)|j\rangle = \frac{1}{T} \int_0^T dt' \langle i|\tilde{j}, t'\rangle e^{i(E_l' + i\eta)t'}, \quad (10)$$

where the terms involving transitions between different energies have been neglected with the relative accuracy $\delta = 1/T\Delta E$, with ΔE as the minimum increment in the energies E_l .

To obtain the time dependent ket $|\tilde{j}; T\rangle$, a direct numerical integration of the Schrödinger equation can be performed, as in the “leap-frog” algorithm.²⁰ This is a second order, symmetrized differencing scheme, accurate up to $(H\Delta t)^2$. In this form, Eq. (4) becomes

$$\begin{aligned} |\tilde{j}; t + \Delta t\rangle &= -2i\Delta t H |\tilde{j}; t\rangle + |\tilde{j}; t - \Delta t\rangle \\ &\quad - 2i\Delta t |j\rangle \sum_l e^{-i(E_l + i\eta)t} \theta(t), \end{aligned} \quad (11)$$

with a time step Δt determined by $\Delta t = \beta/E_{max}$, where E_{max} is the absolute value of the extreme eigenvalue and β is a parameter whose value is less than 1 in order for the solution to be stable.²¹

Estimating the trace in Eq. (3) requires a suitable basis set, such as the local orbital basis. It is more efficient, however, to choose a randomized version of this basis, described by a ket $|\phi\rangle = \sum_{n=1}^N |n\rangle \exp(-i\phi_n)$, where $|n\rangle$ are the tight-binding orbitals and ϕ_n are random numbers in the $[0, 2\pi]$ interval. For a given operator A , $\langle \phi|A|\phi\rangle \simeq \sum_n \langle n|A|n\rangle$ within the statistical errors of $1/\sqrt{N}$.

III. RESULTS AND DISCUSSION

In this section, we show the results of our computation based on the previously outlined algorithm. First, we study the Fermi energy dependence of the spin-Hall conductivity, presented in Fig. 1. Random Fermi energies in the interval $[-4t, 4t]$ are considered and we average over 2000 samples




FIG. 2. (Color online) Spin-Hall conductivity as a function of system size in units of $e/8\pi$ represented on a logarithmic scale. Fermi energy is fixed to $E_F = -3.0t$ and the spin precession length is $L_s \simeq 16a_0$, corresponding to $V_R = 0.2t$. Mean free path ranges from $l \simeq 45a_0$ for $W=0.6$ down to a few lattice constants when disorder increases. Averages are done over 10^3 disorder samples and for each sample, 200 initial state vectors are considered.

for each system size. For clean systems and for states in the band, σ_{SH} is close to 0.8 (in unit of $e/8\pi$), except at half filling where, due to electron-hole symmetry considerations, it vanishes. In the presence of disorder, the calculated value of σ_{SH} decreases, our results reproducing very well the known behavior previously obtained in the Landauer-Büttiker formalism^{9,10} or by the analytical Kubo formula.¹⁵

The dependence of the spin-Hall conductivity on the system size is shown in Fig. 2. One is interested in finding out whether the variation of σ_{SH} is dramatically changed by disorder and in determining the length scale at which this change occurs. For this, the two relevant parameters are the electronic mean free path l and the spin-precession length L_s . In a quasiclassical approximation $l = 12\hbar v_F a_0 / (2\pi N(E_F)W)^2$, where v_F is the Fermi velocity and $N(E_F)$ is the density of states at the Fermi energy measured from the bottom of the band. The spin-precession length is defined in terms of the Rasba coupling constant by $L_s = \pi t a_0 / V_R$.

The electronic mean free path is the length scale that separates the ballistic from the diffusive regimes, with a ballistic behavior for system sizes smaller than l and diffusive otherwise. We found that the crossover between these two regimes is smooth, without any dramatic change in the overall behavior of the spin-Hall conductivity. The only observable effect is a decrease of the spin-Hall conductivity when disorder increases. For example, when $W=0.6t$, $l \simeq 45a_0$, while for $W=1.0t$, $l \simeq 16a_0$, whereas, as can be seen in Fig. 2, the behavior of the spin-Hall conductivity remains unchanged. At the same time, for system sizes below L_s , σ_{SH} always monotonically increases, reaches a plateau between L_s and $2L_s$, and then decreases for large system sizes, and is expected to vanish in the thermodynamic limit, as in Refs. 14 and 15. The spin-precession length, therefore, is the characteristic length scale at which a crossover between the different regimes of the spin-Hall conductivity is expected. In the semiclassical regime, a scaling analysis is appropriate. We find that for a given Fermi energy, the size dependence of the spin-Hall conductivity can be very well fitted with an exponential function $\sigma_{SH} \simeq \exp(-L/\xi)$, where ξ is a characteristic length that depends on the disorder strength, which is diver-




FIG. 3. Disorder dependence of the characteristic length scale ξ for Fermi energy $E_F = -3.0t$ and spin-orbit interaction strength $V_R = 0.2t$.

gent in a clean system. In Fig. 3, we present the dependence of this characteristic length as a function of disorder which follows a power law $\xi \approx \xi_0 (t/W)^{0.57}$ behavior, with the best fit $\xi_0 \approx 46a_0$. We remark again that the system size where σ_{SH} starts to decrease is strongly conditioned by L_s , rather than l as can be inferred from the similarities between the behaviors drawn for different values of the disorder for the same L_s . In all our calculations, the decrease in the spin-Hall conductivity starts for system sizes $L \approx 2L_s$, regardless of the value of the electronic mean free path.

IV. CONCLUSIONS

In this work, we study the effect of the spin-precession length scale and of the electronic mean free path on the the spin-Hall conductivity in different regimes, by adapting the particle-source algorithm to spin transport in systems with SOI in the framework of the tight-binding approximation. The dependence of σ_{SH} on the Fermi energy is also investigated. Our main finding is that the spin precession length is the critical length scale for the spin-Hall behavior. For a system size smaller than L_s , the spin-Hall conductivity increases even in the presence of disorder, reaches a plateau between L_s and $2L_s$, and then, in the semiclassical limit (when $L \gg L_s$), decreases exponentially. In the thermodynamic limit, σ_{SH} is zero for any amount of disorder present in the system. We have also shown that the electronic mean free path does not play a fundamental role in the spin-Hall conductivity behavior.

ACKNOWLEDGMENTS

This research was supported by the Hungarian OTKA Grants Nos. NF061726 and T046303, and by the Romanian CNCSIS Grant 1.97.

- ¹J. Sinova, D. Culcer, Q. Niu, N. A. Sinitsyn, T. Jungwirth, and A. H. MacDonald, Phys. Rev. Lett. **92**, 126603 (2004).
- ²S. Murakami, Phys. Rev. B **69**, 241202(R) (2004).
- ³E. I. Rashba, Sov. Phys. Solid State **2**, 1109 (1960); M. I. Dzyanov and V. I. Perel, Sov. Phys. JETP **33**, 1053 (1971).
- ⁴G. Dresselhaus, Phys. Rev. **100**, 580 (1955).
- ⁵J. Schliemann and D. Loss, Phys. Rev. B **69**, 165315 (2004); B. A. Bernevig, J. P. Hu, E. Mukamel, and S.-C. Zhang, *ibid.* **70**, 113301 (2004); B. A. Bernevig and S.-C. Zhang, *ibid.* **72**, 115204 (2005); E. I. Rashba, *ibid.* **68**, 241315(R) (2003); L. Hu, J. Gao, and S.-Q. Shen, *ibid.* **68**, 153303 (2003); A. A. Burkov, A. S. Nunez, and A. H. MacDonald, *ibid.* **70**, 155308 (2004).
- ⁶J. I. Inoue, G. E. W. Bauer, and L. W. Molenkamp, Phys. Rev. B **70**, 041303(R) (2004).
- ⁷O. V. Dimitrova, Phys. Rev. B **71**, 245327 (2005).
- ⁸Naoyuki Sugimoto, Shigeki Onoda, Shuichi Murakami, and Naoto Nagaosa, Phys. Rev. B **73**, 113305 (2006).
- ⁹C. P. Moca and D. C. Marinescu, Phys. Rev. B **72**, 165335 (2005).
- ¹⁰B. K. Nikolić, S. Souma, L. R. Zárbo, and J. Sinova, Phys. Rev. Lett. **95**, 046601 (2005).
- ¹¹S.-Q. Shen, M. Ma, X. C. Xie, and F. C. Zhang, Phys. Rev. Lett. **92**, 256603 (2004).
- ¹²J. Wunderlich, B. Kaestner, J. Sinova, and T. Jungwirth, Phys. Rev. Lett. **94**, 047204 (2005).
- ¹³H. Tanaka and M. Itoh, Phys. Rev. Lett. **81**, 3727 (1998).
- ¹⁴D. N. Sheng, L. Sheng, Z. Y. Weng, and F. D. M. Haldane, Phys. Rev. B **72**, 153307 (2005).
- ¹⁵K. Nomura, J. Sinova, N. A. Sinitsyn, and A. H. MacDonald, Phys. Rev. B **72**, 165316 (2005); K. Nomura, J. Wunderlich, J. Sinova, B. Kaestner, A. H. MacDonald, and T. Jungwirth, *ibid.* **72**, 245330 (2005).
- ¹⁶K. Nomura, Jairo Sinova, T. Jungwirth, Q. Niu, and A. H. MacDonald, Phys. Rev. B **71**, 041304(R) (2005).
- ¹⁷E. M. Hankiewicz, L. W. Molenkamp, T. Jungwirth, and Jairo Sinova, Phys. Rev. B **70**, 241301(R) (2004).
- ¹⁸B. K. Nikolić, Liviu P. Zárbo, and S. Souma, Phys. Rev. B **72**, 075361 (2005).
- ¹⁹T. Iitaka, Riken Super Computing Prog. Rep. **1**, 1 (1996); T. Iitaka, S. Nomura, H. Hirayama, X. Zhao, Y. Aoyagi, and T. Sugano, Phys. Rev. E **56**, 1222 (1997); H. Tanaka, Phys. Rev. B **57**, 2168 (1998).
- ²⁰T. Iitaka, Phys. Rev. E **49**, 4684 (1994).
- ²¹A. Askar and A. S. Cakmak, J. Chem. Phys. **68**, 2794 (1978).