

**Clemson University**  
**TigerPrints**

---

All Theses

Theses

---

12-2015

# Weeds, nitrogen, and yield: Measuring the effectiveness of an organic no-till system

David Robb

Clemson University, [drobb@clemson.edu](mailto:drobb@clemson.edu)

Follow this and additional works at: [https://tigerprints.clemson.edu/all\\_theses](https://tigerprints.clemson.edu/all_theses)

 Part of the [Plant Sciences Commons](#)

---

## Recommended Citation

Robb, David, "Weeds, nitrogen, and yield: Measuring the effectiveness of an organic no-till system" (2015). *All Theses*. 2261.  
[https://tigerprints.clemson.edu/all\\_theses/2261](https://tigerprints.clemson.edu/all_theses/2261)

This Thesis is brought to you for free and open access by the Theses at TigerPrints. It has been accepted for inclusion in All Theses by an authorized administrator of TigerPrints. For more information, please contact [kokeefe@clemson.edu](mailto:kokeefe@clemson.edu).

WEEDS, NITROGEN, AND YIELD:  
MEASURING THE EFFECTIVENESS OF AN ORGANIC NO-TILL SYSTEM

---

A Thesis  
Presented to  
the Graduate School of  
Clemson University

---

In Partial Fulfillment  
of the Requirements for the Degree  
Master of Science  
Plant and Environmental Science

---

by  
David Charles Robb  
December 2015

---

Accepted by:  
Geoffrey W. Zehnder, Committee Chair  
William C. Bridges, Jr.  
Dara M. Park  
Robin W. Kloot

## ABSTRACT

Previous research in the mid-Atlantic and midwestern USA has identified advantages and drawbacks of “organic no-till” vegetable production, but few studies have been conducted in the warmer southeastern region. The purpose of this study was to examine the effects of tillage [no-till (NT) vs. conventional tillage (CT) of a cereal rye/crimson clover cover crop] and three nitrogen fertilization rates on organic tomato (*Solanum lycopersicum* L.) and summer squash (*Cucurbita pepo* L.) yield, weed suppression, and soil N dynamics in two years in a soil series in Clemson, SC. Squash yields were similar between tillage treatments in both years. NT tomato yields were 43% greater than CT yields in 2014, whereas CT tomato yields were 46% greater than NT yields in 2015. Squash and tomato yields per unit of management labor (time) were significantly greater in NT compared to CT treatments for both years. There were no statistical differences in squash and tomato yields between N fertilization treatments in either year. Pre-season soil N levels were significantly higher in NT tomato plots in 2014 but similar between tillage treatments in tomato plots in 2015 and in squash plots both years. Post-season soil N levels in tomato plots were similar between tillage treatments both years. Post-season soil N levels were significantly higher in NT squash plots in 2014 and in CT squash plots in 2015. Roller-crimped NT mulches provided adequate early-season weed suppression in both years and saved considerable weed management and seedbed preparation labor. Overall, the results demonstrated that organic no-till is a viable method for reduced tillage summer vegetable production in the South Carolina Piedmont region.

## **ACKNOWLEDGEMENTS**

I would like to thank my adviser, Dr. Geoff Zehnder, for his support and guidance over the many years of this study and the work leading up to it. I would also like to thank the rest of my committee: Dr. Billy Bridges for his help making sense of my data, Dr. Dara Park for her help with the experiment setup and literature review, and Dr. Robin “Buz” Kloot for his assistance with the soil analyses. Lastly, I would like to thank my dear wife, Cara, for her support throughout this entire process.

## TABLE OF CONTENTS

	Page
ABSTRACT.....	ii
ACKNOWLEDGEMENTS.....	iii
TABLE OF CONTENTS.....	iv
LIST OF TABLES.....	v
CHAPTER ONE: INTRODUCTION AND LITERATURE REVIEW.....	1
Organic Agriculture .....	1
Advantages of No-till Systems .....	3
Cover Cropping for Soil Health and Weed Management.....	8
Organic No-till.....	9
Research Objective .....	16
Literature Cited.....	16
CHAPTER TWO: WEEDS, NITROGEN, AND YIELD: MEASURING THE EFFECTIVENESS OF AN ORGANIC NO-TILL SYSTEM.....	24
Abstract.....	24
Introduction.....	25
Materials and Methods.....	27
Results.....	34
Discussion.....	36
Acknowledgements.....	40
Literature Cited.....	40

## LIST OF TABLES

	Page
Table 1. Total labor inputs by tillage treatment.....	45
Table 2. Effects of tillage (NT and CT) on average plant available N, total N, and CO <sub>2</sub> -C, pre-season analysis.....	46
Table 3a. Effects of tillage (NT and CT) and N fertilization (0, 58, and 116 kg ha <sup>-1</sup> ) on average plant available N, total N, and CO <sub>2</sub> -C, post-season analysis: tomato .....	47
Table 3b. Effects of tillage (NT and CT) and N fertilization (0, 58, and 116 kg ha <sup>-1</sup> ) on average plant available N, total N, and CO <sub>2</sub> -C, post-season analysis: squash.....	48
Table 4. Effects of tillage (NT and CT) and N fertilization (0, 58, 116 kg ha <sup>-1</sup> ) on average leaf tissue N (2015) .....	49
Table 5. Effects of tillage (NT and CT) and N fertilization (0, 58, 116 kg ha <sup>-1</sup> ) on average marketable vegetable yields per row .....	50
Table 6. Effects of tillage (NT and CT) on average marketable vegetable yield per unit of effort .....	51

## **CHAPTER ONE: INTRODUCTION AND LITERATURE REVIEW**

### **Organic Agriculture**

Representing just over four percent of overall U.S. food sales and roughly about 0.5% of total U.S. cropland, organic production is a small but steadily growing segment in U.S. agriculture (Greene et al., 2009; USDA ERS, 2015). Among organic food sales, fresh fruits and vegetables are a top selling category, accounting for 43% of organic food sales. (USDA ERS, 2015). Organic food sales quintupled between 1997 and 2007 with consumer demand for organics often outstripping supply (Greene et al., 2009). The growth in demand is largely driven by consumer concerns over the health and environmental impacts of food choices (USDA ERS, 2014). Similarly, organic agriculture has been lauded for pesticide reduction, improvements in soil health, carbon sequestration, and enhanced biodiversity (Greene et al., 2009). Concurrent to the rising popularity of organic foods is the growth of direct-to-consumer marketed foods, with a strong consumer preference for local organically produced foods (Low et al., 2015; USDA ERS, 2014). Forty percent of organic farms in the U.S. market food through direct-to-consumer channels (Low et al., 2015). Organic foods demand a higher price premium than conventionally produced food, which can be a strong marketing incentive for farmers to switch to organic practices (Greene et al., 2009; USDA ERS, 2014).

Organic farming, at its core, is soil-health centric (Kuepper, 2010). Early organic farmers, promoted traditional agricultural practices such as animal and green manuring,

composting, crop rotation and diversified cropping to build and feed the “soil food web” (Kuepper, 2010). These traditional practices are codified in the National Organic Program (NOP), the governing document for organic agriculture, which emphasizes soil health by requiring organic producers to use cover crops, crop rotations and application of plant and animal materials (NOP, 2015). Weed management, though, is one of the biggest challenges to organic crop production (Sooby et al., 2007; Riemens et al., 2007), and the prohibition of synthetic herbicides and lack of effective organic alternatives makes weed management more complex in organic systems compared to conventional ones (DeDecker et al., 2014).

Weed management strategies in organic farming include cultural practices such as cover cropping, crop rotations, diversified cropping as well and stale seedbed/delayed planting techniques and mechanical practices such as mowing and tilling (Morse and Creamer, 2006; Sooby et al., 2007; Schonbeck and Morse, 2007). Despite the widely known deleterious effects of soil disturbance on soil health organic farmers, out of necessity, rely heavily on soil tillage for weed management (Schonbeck and Morse, 2007; Morse and Creamer, 2006). Indeed, organic farmers are confounded by the seasonal paradox of both improving and damaging soil health (Schonbeck and Morse, 2007).

A growing number of organic farmers, though, have begun finding success managing weeds while concomitantly improving soil health by blending traditional cover cropping for weed management with reduced tillage techniques (Leavitt et al., 2011; Mirsky et al., 2013).



## **Advantages of No-till Systems**

### **Background**

No-till is a reduced tillage practice in which soil disturbance is all but eliminated between harvest and the subsequent season's planting (Köller, 2003). No-till and other reduced tillage practices spread gradually in the United States in the wake of technological innovations during and after World War II (Lal et al., 2007; Magdoff & Van Es, 2009). Chemical advancements such as the broadleaf herbicide 2-4,D (1945), and the non-selective herbicide paraquat (1962), created tillage alternatives for farmers who, prior to their invention, relied on primary (moldboard plow) and secondary tillage (chisel and/or disc plow) for weed management and seedbed preparation (Lal et al., 2007; Magdoff & Van Es, 2009; Coughenour and Chamala, 2000). Farmer adoption of reduced tillage practices spread gradually in the second half of the 20<sup>th</sup> century (Coughenour and Chamala, 2000). Currently, over 40% of agronomic crop acreage in the U.S. is managed using some form of conservation tillage, mostly no-till (CTIC, 2004). Several decades of no-till research, primarily regarding conventionally grown agronomic crops, has demonstrated a number of beneficial attributes associated with reduced tillage management.

### **Soil Organic Matter**

No-till practices leave abundant residue on the soil surface resulting in slower residue decomposition than when incorporated (Magdoff and Weil, 2004; Franzluebbers,

2004). Surface-concentrated organic matter is better protected within soil macro-aggregates from consumption by soil fauna and microbes (Franzluebbers, 2004). Eventually, all residues left in the field will decompose to some degree, losing the majority of their carbon (C) to the atmosphere via microbial respiration (Franzluebbers, 2004), but by slowing down the process of decomposition, the rate of C loss can be lessened and soil organic matter (SOM) better preserved (Magdoff and Weil, 2004). Over time, continually adding residues while limiting or eliminating tillage altogether (i.e. no-till) can gradually increase SOM levels (Franzluebbers, 2004). Decades of research have demonstrated that no-till increases SOM. Longer-term studies in different climatic zones in North America demonstrated higher SOM levels in no-till systems compared to conventionally tilled systems: barley (Arshad et al., 1990); maize and soybeans, (Dick, 1983); maize (Blevins et al., 1977); wheat (Bauer and Black, 1981); and maize and soybean (Edwards et al., 1992).

### Soil Biology

Microorganism diversity and functional diversity are greatly affected by soil disturbance (Kennedy et al., 2004; Altieri, 1999). Tillage disrupts at least the top 15-25cm of the soil profile and distributes organic matter more evenly throughout the plow zone creating a less stratified, more homogenous surface layer with reduced soil microbial biomass and soil microbial species density (Franzluebbers, 2004; Altieri, 1999; Seiter and Horwath, 2004). In contrast, no-till systems, which increase and concentrate SOM in the upper centimeters of the soil profile increase microbial biomass and diversity

in surface soils (Franzluebbers, 2004; Seiter and Horwarth, 2004) and generally have higher fungi-to-bacteria ratios (Frey et al., 1999; Andrade et al., 2003). A number of studies (Aslam et al., 1999; Hubbard et al., 1999; Jordan et al., 1997; Kladivko et al., 1997) also have demonstrated earthworm populations are generally larger in no-till compared to tilled systems.

Andrade et al. (2003) reviewed 40 short- and long-term no-till/conventional tillage comparison studies and reported that soil microbial biomass generally was higher in no-till systems regardless of study site and crop rotation. Further, they documented that differences between microbial biomass in the two tillage treatments became more pronounced as the duration of the studies increased (Andrade et al. 2003). Increased microbial biomass and SOM, in turn, increase the mineralization potential of the soil resulting in greater plant available nutrients (Seiter and Horwath, 2004). Over time, no-till systems, therefore, tend to accumulate greater biologically active fractions of SOM such as mineralizable C and nitrogen (N) than conventionally tilled systems (Pekrun et al., 2003; Kennedy et al., 2004; Seiter and Horwath, 2004). Overall, no-till imparts less stress on soil micro- and macrobiota and creates a soil environment closely resembling undisturbed, natural ecosystems (Andrade et al., 2003).

### Nutrient Cycling

No-till residues decompose slower than those incorporated into the soil through tillage, thus the timing of nutrient release can be different between tilled and no-till systems (Magdoff and Weil, 2004). Conventional tillage triggers oxidative processes,

which accelerate the mineralization of nutrients such as N, P, and S for plant uptake (Franzluebbbers, 2004; Magdoff and Weil, 2004). With no-till and other reduced tillage practices, mineralization of N generally is slower (Franzluebbbers, 2004). In the short term, after transitioning from intensive cultivation to no-till management practices, additional N fertilizer may be required to account for slower N mineralization (Pekrun et al., 2003). However, after a transition period – which could last several years to a decade depending on soil conditions – a new equilibrium can be reached in no-till systems wherein increased SOM and microbial biomass lead to greater mineralized N, reducing the need for additional N fertilizer amendments (Magdoff and Weil, 2004; Pekrun, 2003). Increases in SOM and microbial biomass in long-term no-till management, in essence, compensate for slower N mineralization and can lead to increased net N mineralization compared to conventionally tilled systems (Pekrun et al., 2003).

### Agronomic Crop Yields

Franzluebbbers (2004) reviewed over 80 agronomic crop studies comparing no-till, shallow tillage, and deep inversion tillage and found that no-till systems produced no consistent yield advantage compared to tilled systems – yields tended to be comparable between the systems. No-till has also been associated with yield reduction, particularly with direct-seeded crops in poorly drained soils that are slow to warm up in the spring (Cannell and Hawes, 1994; Dick et al., 1991; Van Doren et al., 1976; Lal et al., 1989). Yet, longer-term agronomic crop studies revealed more distinct yield advantages conferred by no-till management when compared to conventional tillage. Dick et al.

(1991) and Ismail et al. (1994), 25-yr and 20-yr maize studies, respectively, reported that yields tended to increase with time under no-till management compared with conventional tillage.

### Soil and Water Conservation

In general, returning plant residue to the soil surface tends to increase soil water entry and storage (Magdoff and Weil, 2004). Residues left on the soil surface also protect the soil from the erosive forces of rainfall and wind (Pimentel et al., 1995). Franzluebbbers et al. (1999) reported that no-till practices resulted in significant reductions in both water runoff and soil loss in land that had previously been under conventional tillage. SOM enhances the soil's ability to retain water by way of direct absorption and by promoting stronger aggregation (Magdoff and Weil, 2004). Thus, soils with a high SOM content have significantly higher available water capacity than soils with similar texture but with less SOM (Hudson, 1994). No-till practices, which maximize soil coverage with residues and increase SOM, tend to result in more erosion-resistant soils with greater soil water content compared conventional tillage (Blevins and Frye, 1993).

### Carbon Sequestration

Agricultural soil conservation practices such as no-till, which increases soil C long term, serve as a potential mitigation strategy to curb greenhouse gas emissions and thus combat accelerated global climate change (Lal, 2010; West and Post, 2002; Six et al., 2004). Estimates of how effective no-till farming could potentially be to reducing

global warming, though, are highly variable and complex due to variations in crop residue, climate, and soil properties (Lal, 2010). Franzluebbers (2010), reviewed 147 long-term no-till/conventional tillage comparison studies from the southeastern U.S. and reported that the agricultural soils in this region alone hold the potential to sequester 0.45 Mg C ha<sup>-1</sup> yr<sup>-1</sup> if converted to no-till management. Six et al. (2004) reported that an estimated 0.031 Mg C ha<sup>-1</sup> yr<sup>-1</sup> of greenhouse gas emissions could be realized by using no-till, mainly by way of reduced fuel consumption.

### **Cover Cropping for Soil Health and Weed Management**

Teasdale et al. (2007) defined cover crops as "a wide range of plants that are grown for various ecological benefits other than as a cash crop." Cover crops are commonly used to fill a between-season temporal and/or spatial niche that otherwise would be occupied by unmanageable weed populations in the absence of cash crops (Lal et al., 1991; Teasdale, 1996). Cover crops provide a number of beneficial ecosystem functions: cover crops add organic C and N to the soil, uptake nutrients and recycle them, improve soil aggregation, increase macropore formation, enhance water infiltration, conserve soil moisture, protect the soil surface from wind and water erosion, improve soil tilth, and, in general, enhance soil biological diversity (Seiter and Horwath, 2004; Lal et al., 1991; Schonbeck and Morse, 2006; Schonbeck and Morse, 2007, Clark, 2007; Creamer and Baldwin, 2000).

Regarding weed management, cover crops cover the soil surface with living plant tissue and outcompete weeds for sunlight, water, and nutrients (Mohler and Teasdale,

1993; Balkcom et al., 2007; Teasdale, 1996). A number of studies also have shown that cover crops suppress weeds to a varying degree through phytotoxic allelochemicals (e.g. phenolic acids, coumarins, benzoquinones, terpenoids, glucosinolates, and tannins) released into the soil (Reberg-Horton et al., 2005; Barnes and Putnam, 1983; Putnam and DeFrank, 1983; Chung et al. 2002; Seigler 1996; Swain, 1977). In addition to outcompeting weeds while growing, cover crops can be terminated and their residues left on the soil surface as an *in situ* mulch to suppress weed growth early in the growing season (5-7 weeks), commonly known as the “critical weed-free period”, when cash crops have yet to establish dense canopies to out-compete weeds (Schonbeck and Morse, 2007). Surface-placed residues suppress weeds by intercepting and reflecting light transmittance, which stymies weed seed germination (Mohler and Teasdale, 1993; Teasdale and Mohler, 2000). Further, the physical presence of mulch on the soil surface impedes the emergence of germinated weeds (Teasdale and Mohler, 1993). Mulch can also alter soil temperature conditions, delaying the emergence of weed seeds by keeping soil temperatures sub-optimal early in the season (Teasdale and Mohler, 1993).

## **Organic No-till**

### Background

Conventional no-till, an agricultural practice that relies heavily on chemically based weed suppression, is not practicable in organic production systems (Schonbeck and Morse, 2007; Moyer, 2011). Organic farmers are prohibited from using synthetic substances, such as herbicides, in organic production and handling (NOP, 2015)

However, in the last several decades, advancements in cover crop research and improvements in no-till equipment have led to a growing interest among organic farmers in developing no-till strategies that are compatible with organic systems (Moyer, 2011; Schonbeck, 2015; Schonbeck and Morse, 2007). “Organic no-till”, as it is known, is a cover crop centric reduced tillage system whereby an *in situ* cover crop mulch alone provides weed suppression and enables farmers to reduce tillage in organic systems (Moyer, 2011; Schonbeck, 2015; Schonbeck and Morse, 2007).

In an organic no-till system, a cover crop – normally a winter annual (e.g. cereal rye, hairy vetch, crimson clover) – is planted in the fall at a dense seeding rate and overwintered until mature in the spring (Moyer, 2011; Schonbeck and Morse, 2007, Schonbeck, 2015). Variations of this cropping schedule exist. For instance, faster maturing, short-season crops (e.g. cowpeas, millet, sorghum-Sudan grass, buckwheat and sunn hemp) can be planted in the summer for fall cropping or planted in the fall and “frost” killed for early spring cropping (Schonbeck and Morse, 2007, Schonbeck, 2015). However, most published research regarding organic no-till has focused on fall-planted over-wintering cover crops. Large amounts of plant biomass are favorable when using cover crop *in situ* mulches for in-season weed suppression (Schonbeck and Morse, 2007). Mohler and Teasdale (1993) demonstrated that normal cover crop biomass levels, approximately 3,000 kg ha<sup>-1</sup> or less, did not provide adequate weed suppression. They reported that biomass levels two and four times the natural amount were needed for adequate suppression of annual grasses and annual/perennial broadleaf weeds, respectively (Mohler and Teasdale 1993).



## No-till Cover Crops

Cereal rye (*Secale cereale* L.) is a fall-planted cover crop that is adaptable for most USDA plant hardiness zones (Clark, 2007). Rye, which has well-documented allelopathic properties, can produce a large amount of aboveground biomass ( $>9,000$  kg ha<sup>-1</sup>), and is an optimal cool-season cover crop for organic no-till cropping systems in the southeastern U.S. (Mirsky et al., 2009; Reberg-Horton et al., 2012; Schonbeck and Morse, 2006). Cereal rye, especially as it matures, has a high C/N ratio and is thus slower to decompose than other cover crops, such as legumes (Teasdale and Abdul-Baki, 1998; Mohler and Teasdale, 1993). Mohler and Teasdale (1993) demonstrated that cereal rye maintains weed suppression over a longer growing season compared to legume cover crops, which generally have lower C/N ratios. Rye also is a scavenger or “catch” crop for residual N in the fall and winter, reducing the quantity of leachable NO<sub>3</sub><sup>-</sup> in the soil (Clark, et al., 1994; Meisinger, 1991; Shipley, et al., 1992).

The high C/N ratio of mature cereal rye (25:1 to 55:1), which increases markedly later in the growing season as it matures, can lead to immobilization of N in the soil (Clark, et al., 1994; Creamer et al., 1997; Teasdale and Abdul-Baki, 1998). To reduce N immobilization while maintaining adequate biomass for weed suppression, rye is often planted with a companion winter legume such as crimson clover (*Trifolium incarnatum* L.) or hairy vetch (*Vicia villosa* Roth) (Schonbeck and Morse, 2007). Hairy vetch is better adapted to slightly cooler climates (USDA zones 5-7) than crimson clover and is winter hardy to USDA zones 3-4 (Clark, 2007). Compared to vetch, crimson clover is

better adapted to warmer U.S. climates (USDA zones 8-10) including much of the southeastern U.S. (Clark, 2007). Both hairy vetch and crimson clover can contribute considerable N to the soil (75-100 kg ha<sup>-1</sup> N fertilizer equivalence), which reduces N immobilization in rye cover crop systems (Fageria et al., 2005; Hargrove, 1986; Clark et al., 1994; Creamer et al., 1997). Another advantage of using a cover crop biculture (grass + legume) is that it can accumulate more “topgrowth” N than either cover crop grown alone because by depleting the soil of N, the grass, in essence, drives greater biological N fixation on behalf of the legume (Creamer et al., Clark et al., 1994).

### Cover Crop Termination

Proper mechanical termination of cover crops is necessary to make organic no-till systems work (Schonbeck and Morse, 2007; Creamer and Dabney, 2002; Moyer, 2011). Organic no-till cover crops can be killed mechanically with a number of farm implements (e.g. mowers, cultipackers, undercutters, rolling stalk choppers, crop rollers) but optimal results have been documented with specialized roller-crimping devices – heavy, often water-filled cylinders with welded steel blades – that crush cover crop tissue and press the crop residue flat onto the soil surface (Schonbeck and Morse, 2007; Moyer, 2011, Mischler, et al., 2010, Ashford and Reeves, 2003; Morse, 1999). Mischler et al. (2010) and Mirsky et al. (2009) demonstrated that roller-crimpers provide consistent cover crop control comparable to herbicides. Ashford and Reeves (2003) demonstrated that roller-crimping was more economical than chemical weed control.

Unlike mowing, which tends to shred residues finely and distribute them

unevenly, roller-crimping keeps residues intact and in place, leaving behind a thick weed barrier that persists longer in season than when mowed (Schonbeck and Morse, 2007; Mirsky et al. 2009, Creamer and Dabney, 2002; Moyer, 2011). Further, roller-crimped cover crop residue is uniformly pressed in the direction of travel, which reduces the risk of hairpins and clogged equipment if followed with no-till planting equipment (Moyer, 2011; Creamer and Dabney, 2002). Roller-crimpers also are more cost-effective compared to other methods of cover crop termination, such as mowing. Roller-crimpers can be operated at faster speeds than when mowing and do not require an energy-intensive power take-off (PTO) drive (Mirsky et al., 2013). Ashford and Reeves (2003) reported that energy requirements for roller-crimper operation were one tenth that of a rotary mower.

Timing of the cover crop termination is critical in organic no-till systems (Moyer, 2011; Mirsky et al., 2009; Mischler et al., 2010; Creamer and Dabney, 2002). The cover crop needs to have time to generate sufficient biomass – if killed too early, not enough biomass has accumulated for lasting weed suppression (Schonbeck and Morse, 2007; Moyer, 2011, Mirsky et al., 2013). In experiments examining cover crop kill-date effects, Clark et al. (1994), Clark et al. (1995), Wagger (1989), and Mirsky et al. (2011) demonstrated that significant increases in cover crop biomass could be achieved by delaying spring termination by just 2-4 weeks. In general, mechanical control of cover crops improves with increasing plant maturity (Mirsky et al., 2012). Conversely, if the cover crop is left in the field too long and viable seed is formed, the seed from the cover crop can create weed problems in future cropping seasons (Moyer, 2011).

For optimal biomass production with little risk of carry-over weed seed, mechanically killed cover crops for organic no-till are best terminated at anthesis or greater but before viable seed has been produced (Moyer, 2011; Schonbeck, 2015; Mirsky et al., 2009). In an organic no-till study examining roller-crimped cereal rye, Mirsky et al. (2009) found that cereal rye regrowth was consistently controlled by terminating the crop at anthesis (Zadoks growth stage 61) or greater. Ashford and Reeves (2003) had similar findings: 85% and 95% cover crop control when rye was terminated at anthesis and soft dough stages, respectively. Another concern regarding the timing of cover crop termination is regrowth of a cover crop that was too immature at time of termination. Cover crop regrowth can create a carryover weed problem in the field as lingering cover crops can compete with cash crops for water and nutrients much like weeds do (Moyer, 2011; Mischler et al., 2010). Waiting until anthesis or greater before terminating the cover crop diminishes the chance for regrowth of the cover crop (Mischler et al., 2010; Mirsky et al., 2009).

### Vegetable Yields

Similar to the aforementioned conventional no-till agronomic crop studies, yield results in organic no-till vegetable production have been mixed. Yield reductions associated with no-till rye and/or vetch mulches were documented in squash (Leavitt et al., 2011), bell pepper (Diaz-Perez et al., 2008; Leavitt et al., 2011), and tomato production (Leavitt et al., 2011). On the contrary, comparable or positive yield responses in organic no-till systems compared to conventionally tilled systems were reported in

tomatoes (Abdul-Baki et al., 1996; Madden et al., 2004; Delate et al., 2012). Abdul-Baki et al. (1996) documented higher tomato yields with lower N fertilizer inputs using mowed hairy vetch, crimson clover, and rye + hairy vetch mulches compared to black polyethylene mulch (plasticulture) treatments. They also demonstrated that using cover crop mulches instead of plastic mulches potentially could save growers an estimated \$1850/ha in reduced fertilizer, herbicide and equipment costs (Abdul-Baki et al., 1996). Indeed, one of the biggest potential advantages to using organic no-till practices is the potential for greater economic returns through lessened production and management costs (Moyer, 2011).

### Future Challenges

Organic no-till, though, is not without its share of pitfalls. Common problems researchers have identified regarding organic no-till production include: sub-optimal soil temperatures early in the season and decreased degree growing days due to the cooling effect of cover crop mulches; loss of earliness due to a lack of synchrony between cover crop maturity and optimal cash crop planting dates; N immobilization when using high C/N cover crops (i.e. rye); increased weed pressure particularly when cover crop stands are inadequate; and reduced N mineralization due to a lack of cover crop incorporation (Leavitt et al., 2011; Schonbeck and Morse, 2007; Schonbeck, 2015; Moyer, 2011; Creamer et al., 1997; Morse, 1999; Mirsky et al., 2011). To keep weeds (particularly perennial weeds) in check, continuous no-till is not recommended in organic no-till systems – eventually some tillage is required between cropping seasons to manage weed

pressure (Schonbeck and Morse, 2007). Therefore, organic no-till is viewed not as a “zero” tillage system but more of a rotational tillage method – a means to eliminate most, but not all, tillage events from crop production (Schonbeck and Morse, 2007; Mirsky et al., 2012).

### **Research Objective**

The purpose of this research is to evaluate organic no-till production for weed management, vegetable yield, and nutrient management compared to a conventionally tilled system.

### **Literature Cited**

- Abdul-Baki, A. A., & Teasdale, J. R. (2007.). Sustainable production of fresh market tomatoes with organic mulches. USDA Farmers' Bulletin 2280. Retrieved from: <http://www.ars.usda.gov/is/np/SustainableTomatoes2007/TomatoPub.pdf>
- Abdul-Baki, A., Stommel, J., Watada, A., Teasdale, J., and Morse, R. (1996). Hairy vetch mulch favorably impacts yield of processing tomato. *HortScience*, 31(338), 340.
- Altieri, M. A. (1999). The ecological role of biodiversity in agroecosystems. *Agriculture Ecosystems & Environment*, 74(1-3), 19-31.
- Arshad, M. A., Schnitzer, M., Angers, D. A., and Ripmeester, J. A. (1990). Effects of till vs no-till on the quality of soil organic matter. *Soil Biology and Biochemistry*, 22(5), 595-599.
- Ashford, D. L. and Reeves, D. W. (2003). Use of a mechanical roller-crimper alternative kill method for cover crops. *American Journal of Alternative Agriculture*, 18, 37-45.
- Aslam, T., Choudhary, M. A., and Saggarr, S. (1999). Tillage impacts on soil microbial biomass C, N and P, earthworms and agronomy after two years of cropping following permanent pasture in New Zealand. *Soil & Tillage Research*, 51(1-2), 103-111.

- Balkcom, K.S., Schomberg, H., Reeves, D. W. and Clark, A. J. (2007). Managing cover crops in conservation tillage systems. In A. Clark (Ed.), *Managing cover crops profitably* (3rd ed., pp. 44-61). College Park, MD: SARE.
- Barnes, J. P. and Putnam, A. R. (1983). Rye residues contribute weed suppression in no-tillage cropping systems. *Journal of Chemical Ecology*, 9(8), 1045-1057.
- Bauer, A., and Black, A. L. (1981). Soil carbon, nitrogen, and bulk density comparisons in two cropland tillage systems after 25 years and in virgin grassland. *Soil Science Society of America Journal*, 45, 1166-1170.
- Blevins, R. L. and Frye, W. F. (1993). Conservation tillage: an ecological approach to soil management. *Advances in Agronomy*, 51(34), 77.
- Blevins, R. L., Thomas, G. W., and Cornelius, P. L. (1977). Influence of no-tillage and nitrogen-fertilization on certain soil properties after 5 years of continuous corn. *Agronomy Journal*, 69(3), 383-386.
- Cannell, R. Q., and Hawes, J. D. (1994). Trends in tillage practices in relation to sustainable crop production with special reference to temperate climates. *Soil and Tillage Research*, 30(2-4), 245-282.
- Chung et al. (2002). Screening of allelochemicals on barnyardgrass (*Echinochloa crus-galli*) and identification of potentially allelopathic compounds from rice (*Oryza sativa*) variety hull extracts. *Journal of Crop Protection*, 21, 913-920.
- Clark, A. J. (Ed.). (2007). *Managing cover crops profitably* (3rd ed.). College Park, MD: SARE.
- Clark, A. J., Decker, A. M., and Meisinger, J. J. (1994). Seeding rate and kill date effects on hairy vetch cereal rye cover crop mixtures for corn production. *Agronomy Journal*, 86(6), 1065-1070.
- Clark, A. J., Decker, A. M., Meisinger, J. J. and McIntosh, M. S. (1997). Kill date of vetch, rye, and a vetch-rye mixture: I. cover crop and corn nitrogen. *Agronomy Journal*, 89(3), 427-434.
- Clark, A. J., Decker, A. M., Meisinger, J. J., Mulford, F. R., and McIntosh, M. S. (1995). Hairy vetch kill date effects on soil-water and corn production. *Agronomy Journal*, 87(3), 579-585.
- Conservation Technology Information Center (CTIC). (2004). National crop residue management survey. Retrieved from: <http://www.ctic.org/CRM/>

- Coughenour, C. M., and Chamala, S. (2000). Conservation tillage and cropping innovation: constructing the new culture of agriculture. Ames, Iowa: Iowa State University Press.
- Creamer, N. G. and Dabney, S. M. (2002). Killing cover crops mechanically: review of recent literature and assessment of new research results. *American Journal of Alternative Agriculture*, 17(01), 32-40.
- Creamer, N. G., Bennett, M. A., and Stinner, B. R. (1997). Evaluation of cover crop mixtures for use in vegetable production systems. *HortScience*, 32(5), 866-870.
- Creamer, N.G. and Baldwin, K. R. (2000). An evaluation of summer cover crops for use in vegetable production systems in North Carolina. *HortScience*, 35, 600-603.
- DeDecker, e. a. (2014). Weed management practice selection among midwest U.S. organic growers. *Weed Science*, Vol. 62(3), 520-531.
- Delate, K., Cwach, D., and Chase, C. (2012). Organic no-tillage system effects on soybean, corn and irrigated tomato production and economic performance in Iowa, USA. *Renewable Agriculture and Food Systems*, 27, 49-59.
- Diaz-Perez, J., Silvoy, J., Phatak, S., Ruberson, J., and Morse, R. (2008). Effect of winter cover crops and no-till on the yield of organically grown bell pepper (*Capsicum annuum* L.). In R. Prange and S. Bishop (eds.). (Ed.), □Proc. XXVII IHC-S11 Sustainability through integrated and organic horticulture. pp. 767.
- Dick, W. A. (1983). Organic-carbon, nitrogen, and phosphorus concentrations and pH in soil profiles as affected by tillage intensity. *Soil Science Society of America Journal*, 47(1), 102-107.
- Dick, W. A., McCoy, E. L., Edwards, W. M., and Lal, R. (1991). Continuous application of no-tillage to Ohio soils. *Agronomy Journal*, 83(1), 65-73.
- Edwards, J. H., Wood, C. W., Thurlow, D.L., and Ruf, M. E. (1992). Tillage and crop-rotation effects on fertility status of a Hapludult soil. *Soil Science Society of America Journal*, 56(5), 1577-1582.
- Fageria, N. K., Baligar, V. C., and Bailey, B. A. (2005). Role of cover crops in improving soil and row crop productivity. *Communications in Soil Science and Plant Analysis*, 36(19-20), 2733-2757.
- Franzluebbers, A. J. (2004). Tillage and residue management effects on soil organic matter. In R. Weil and F. Magdoff (Eds.), *Soil organic matter in sustainable agriculture* (pp. 227-268). Boca Raton, FL: CRC Press.



- Franzluebbers, A. J., Hons, F. M., and Zuberer, D. A. (1994). Long-term changes in soil carbon and nitrogen pools in wheat management-systems. *Soil Science Society of America Journal*, 58(6), 1639-1645.
- Frey, S. D., Elliott, E. T., and Paustian, K. (1999). Bacterial and fungal abundance and biomass in conventional and no-tillage agroecosystems along two climatic gradients. *Soil Biology and Biochemistry*, 31(4), 573-585.
- Greene et al. (2009). Emerging issues in the U.S. organic industry, No. EIB-55. U.S. Department of Agriculture, Economic Research Service. Retrieved from: <http://www.ers.usda.gov/publications/eib-economic-information-bulletin/eib55.aspx>
- Hargrove, W. L. (1986). Winter legumes as a nitrogen-source for no-till grain-sorghum. *Agronomy Journal*, 78(1), 70-74.
- Hubbard, V. C., Jordan, D., and Stecker, J. A. (1999). Earthworm response to rotation and tillage in a Missouri claypan soil. *Biology and Fertility of Soils*, 29(4), 343-347.
- Hudson, B. D. (1994). Soil organic matter and available water capacity. *Journal of Soil and Water Conservation*, 49(2), 89-194.
- Ismail, I., Blevins, R. L., and Frye, W. W. (1994). Long-term no-tillage effects on soil properties and continuous corn yields. *Soil Science Society of America Journal*, 58(1), 193-198.
- Jordan, D., Stecker, J. A., Cacio-Hubbard, V. N., Li, F., Gantzer, C. J., and Brown, J. R. (1997). Earthworm activity in no-tillage and conventional tillage systems in Missouri soils: a preliminary study. *Soil Biology and Biochemistry*, 29(3-4), 489-491.
- Kennedy, A. C., Stubbs, T. L., and Schillinger, W. F. (2004). Soil and crop management effects on soil microbiology. In R. Weil and F. Magdoff (Eds.), *Soil organic matter in sustainable agriculture* (pp. 295-326). Boca Raton, FL: CRC Press.
- Kladivko, E. J., Akhouri, N. M., and Weesies, G. (1997). Earthworm populations and species distributions under no-till and conventional tillage in Indiana and Illinois. *Soil Biology and Biochemistry*, 29(3-4), 613-615.
- Koller, K. (2003). Techniques of soil tillage. In A. El Titi (Ed.), *Soil tillage in agroecosystems* (pp. 1-27). Boca Raton, FL: CRC Press.

- Kuepper, G. (2010). A brief overview of the history and philosophy of organic agriculture. Kerr Center for Sustainable Agriculture.
- Lal, R. (2010). Managing soils and ecosystems for mitigating anthropogenic carbon emissions and advancing global food security. *Bioscience*, 60(9), 708-721.
- Lal, R. (2013). Enhancing ecosystem services with no-till. *Renewable Agriculture and Food Systems*, 28(02), 102-114.
- Lal, R., Logan, T. J., and Fausey, N. R. (1989). Long-term tillage and wheel traffic effects on a poorly drained mollic ochraqualf in northwest Ohio. 1. soil physical properties, root distribution and grain yield of corn and soybean. *Soil and Tillage Research*, 14(4), 341-358.
- Lal, R., Regnier, E., Eckert, D. J., Edwards, W. M., and Hammond, R. (1991). Expectations of cover crops for sustainable agriculture. In W. L. Hargrove (Ed.), *Cover crops for clean water* (pp. 1-10). Ankeny, IA: Soil and Water Conservation Society.
- Lal, R., Reicosky, D. C., and Hanson, J. D. (2007). Evolution of the plow over 10,000 years and the rationale for no-till farming. *Soil and Tillage Research*, 93(1), 1-12.
- Leavitt, M. J., Sheaffer, C. C., Wyse, D. L., and Allan, D. L. (2011). Rolled winter rye and hairy vetch cover crops lower weed density but reduce vegetable yields in no-tillage organic production. *HortScience*, 46(3), 387-395.
- Low, S. et. al. 2015. (2015). Trends in U.S. local and regional food systems, No. AP-068. U.S. Department of Agriculture, Economic Research Service. Retrieved from: <http://www.ers.usda.gov/media/1763057/ap068.pdf>
- Madden, N. M. et al. (2004). Evaluation of conservation tillage and cover crop systems for organic processing tomato production. *Horttechnology*, 14(2), 243-250.
- Magdoff, F. and Weil, R. (2004). Soil organic matter management strategies. In R. Weil and F. Magdoff (Eds.), *Soil organic matter in sustainable agriculture* (pp. 45-66). Boca Raton, FL: CRC Press.
- Meisinger, J. J., Hargrove, W. L., Mikkelsen, Jr., R. L., Williams, J. R., & Benson, V. W. (1991). Effects of cover crops on groundwater quality. In W. L. Hargrove (Ed.), *Cover crops for clean water* (pp. 57-68). Ankeny, Iowa: Soil and Water Conservation Society.

- Mirsky, S. B. et al. (2012). Conservation tillage issues: Cover crop-based organic rotational no-till grain production in the mid-atlantic region, USA. *Renewable Agriculture and Food Systems*, 27, 31-40.
- Mirsky, S. B. et al. (2013). Overcoming weed management challenges in cover crop-based organic rotational no-till soybean production in the eastern United States. *Weed Technology*, 27(1), 193-203.
- Mirsky, S. B., Curran, W. S., Mortensen, D. A., Ryan, M. R., and Shumway, D. L. (2009). Control of cereal rye with a roller/crimper as influenced by cover crop phenology. *Agronomy Journal*, 101(6), 1589-1596.
- Mirsky, S. B., Curran, W. S., Mortensen, D. M., Ryan, M. R., and Shumway, D. L. (2011). Timing of cover-crop management effects on weed suppression in no-till planted soybean using a roller-crimper. *Weed Science*, 59(3), 380-389.
- Mischler, R. A., Curran, W. S., Duiker, S. W., and Hyde, J. A. (2010). Use of a rolled-rye cover crop for weed suppression in no-till soybeans. *Weed Technology*, 24(3), 253-261.
- Mohler, C. L. and Teasdale, J. R. (1993). Response of weed emergence to rate of *Vicia villosa* Roth and *Secale cereale* L. residue. *Weed Research*, 33(6), 487-499.
- Morse, R. D. (1999). No-till vegetable production – its time is now. *Horttechnology*, 9(3), 373-379.
- Morse, R., and Creamer, N. (2006). Developing no-tillage systems without chemicals: the best of both worlds? In P. Kristiansen, A. Taji, & J. Reganold (Eds.), *Organic agriculture: A global perspective* (pp. 83-91). Ithaca, NY: Comstock Pub. Associates.
- Moyer, J. (2011). *Organic no-till farming*. Austin, TX: Acres U.S.A.
- National Organic Program. (2015). Final rule: Title 7: Agriculture. Part 205. USDA-AMS, Washington, D.C. Retrieved from: [http://www.ecfr.gov/cgi-bin/text-idx?tpl=/ecfrbrowse/Title07/7cfr205\\_main\\_02.tpl](http://www.ecfr.gov/cgi-bin/text-idx?tpl=/ecfrbrowse/Title07/7cfr205_main_02.tpl)
- Pekrun, C., Kaul, H., and Claupein, W. (2003). Soil tillage for sustainable nutrient management. In A. El Titi (Ed.), *Soil tillage in agroecosystems* (pp. 83-113). Boca Raton, FL: CRC Press.
- Putnam, A. R., Defrank, J. and Barnes, J. P. (1983). Exploitation of allelopathy for weed-control in annual and perennial cropping systems. *Journal of Chemical Ecology*, 9(8), 1001-1010.

- Reberg-Horton, S. C. et al. (2005). Changes over time in the allelochemical content of ten cultivars of rye (*Secale cereale* L.). *Journal of Chemical Ecology*, 31(1), 179-193.
- Reberg-Horton, S. C. et al. (2012). Utilizing cover crop mulches to reduce tillage in organic systems in the southeastern USA. *Renewable Agriculture and Food Systems*, 27(1), 41-48.
- Riemens, M., Groeneveld, R., Lotz, L., and Kropff, M. (2007). Effects of three management strategies on the seedbank, emergence and the need for hand weeding in an organic system. *Weed Research*, 47, 442-451.
- Schonbeck, M. W. (2015). What is "organic no-till," and is it practical? Extension Foundation, eOrganic Community of Practice. Retrieved from: [https://www.extension.org/pages/18526/what-is-organic-no-till-and-is-it-practical#.VAuqLhar\\_No](https://www.extension.org/pages/18526/what-is-organic-no-till-and-is-it-practical#.VAuqLhar_No)
- Schonbeck, M. W. and Morse, R. D. (2007). Reduced tillage and cover cropping systems for organic vegetable production. Virginia Association of Biological Farming Info. Sheet, 9-07. Retrieved from: <http://www.sare.org/Learning-Center/SARE-Project-Products/Southern-SARE-Project-Products/Reduced-Tillage-and-Cover-Cropping-Systems-for-Organic-Vegetable-Production>
- Seigler, D. S. (1996). Chemistry and mechanisms of allelopathic interactions. *Agronomy Journal*, 88(6), 876-885.
- Seiter, S. and Horwath, W. R. (2004). Strategies for managing soil organic matter to supply plant nutrients. In R. Weil and F. Magdoff (Eds.), *Soil organic matter in sustainable agriculture* (pp. 269-294). Boca Raton, FL: CRC Press.
- Shiple, P. R., Meisinger, J. J., and Decker, A. M. (1992). Conserving residual corn fertilizer nitrogen with winter cover crops. *Agronomy Journal*, 84(5), 869-876.
- Six, J. et al. (2004). The potential to mitigate global warming with no-tillage management is only realized when practised in the long term. *Global Change Biology*, 10(2), 155-160.
- Sooby, J., Landeck, J., and Lipson, M. (2007). 2007 National Organic Research Agenda. Santa Cruz, CA.: Organic Farming Research Foundation. Retrieved from: <http://ofrf.org/sites/ofrf.org/files/docs/pdf/nora2007.pdf>
- Swain, T. (1977). Secondary compounds as protective agents. *Annual Review of Plant Physiology*, 28, 479-501.

- Teasdale, J. R., & Abdul-Baki, A. A. (1998). Comparison of mixtures vs. monocultures of cover crops for fresh-market tomato production with and without herbicide. *Hortscience*, 33(7), 1163-1166.
- Teasdale, J. R., Coffman, C. B., and Mangum, R. W. (2007). Potential long-term benefits of no-tillage and organic cropping systems for grain production and soil improvement. *Agronomy Journal*, 99(5), 1297-1305.
- Teasdale, J.R. and Mohler, C. L. (1993). Light transmittance, soil temperature, and soil moisture under residue of hairy vetch and rye. *Agronomy Journal*, 85(3), 673-680.
- Teasdale, J.R. and Mohler, C. L. (2000). The quantitative relationship between weed emergence and the physical properties of mulches. *Weed Science*, 48(3), 385-392.
- Teasdale, J. R. (1996). Contribution of cover crops to weed management in sustainable agricultural systems. *Journal of Production Agriculture*, 9(4), 475-479.
- USDA-ERS (U.S. Department of Agriculture–Economic Research Service). (2014). Organic market overview. USDA-ERS, Washington, D.C. Retrieved from: <http://www.ers.usda.gov/topics/natural-resources-environment/organic-agriculture/organic-market-overview.aspx>
- USDA-ERS (U.S. Department of Agriculture–Economic Research Service). (2015). Organic agriculture overview. USDA-ERS, Washington, D.C. Retrieved from: <http://www.ers.usda.gov/topics/natural-resources-environment/organic-agriculture.aspx>
- Van Doren, D. M., Triplett, G. B., and Henry, J. E. (1976). Influence of long term tillage, crop rotation, and soil type combinations on corn yield. *Soil Science Society of America Journal*, 40(1), 100-105.
- Wagger, M. G. (1989). Time of desiccation effects on plant composition and subsequent nitrogen release from several winter annual cover crops. *Agronomy Journal*, 81(2), 236-241.
- West, T. O. and Post, W. M. (2002). Soil organic carbon sequestration rates by tillage and crop rotation: A global data analysis. *Soil Science Society of America Journal*, 66, 1930-1946.

## CHAPTER TWO: WEEDS, NITROGEN, AND YIELD: MEASURING THE EFFECTIVENESS OF AN ORGANIC NO-TILL SYSTEM

### Abstract

Previous research in the mid-Atlantic and midwestern USA has identified advantages and drawbacks of “organic no-till” vegetable production, but few studies have been conducted in the warmer southeastern region. The purpose of this study was to examine the effects of tillage [no-till (NT) vs. conventional tillage (CT) of a cereal rye/crimson clover cover crop] and three nitrogen fertilization rates on organic tomato (*Solanum lycopersicum* L.) and summer squash (*Cucurbita pepo* L.) yield, weed suppression, and soil N dynamics in two years in a soil series in Clemson, SC. Squash yields were similar between tillage treatments in both years. NT tomato yields were 43% greater than CT yields in 2014, whereas CT tomato yields were 46% greater than NT yields in 2015. Squash and tomato yields per unit of management labor (time) were significantly greater in NT compared to CT treatments for both years. There were no statistical differences in squash and tomato yields between N fertilization treatments in either year. Pre-season soil N levels were significantly higher in NT tomato plots in 2014 but similar between tillage treatments in tomato plots in 2015 and in squash plots both years. Post-season soil N levels in tomato plots were similar between tillage treatments both years. Post-season soil N levels were significantly higher in NT squash plots in 2014 and in CT squash plots in 2015. Roller-crimped NT mulches provided adequate early-season weed suppression in both years and saved considerable weed management and seedbed preparation labor. Overall, the results demonstrated that organic no-till is a

viable method for reduced tillage summer vegetable production in the South Carolina Piedmont region.

## **Introduction**

Weed management and the associated labor inputs are consistently some of the biggest challenges to organic crop production (Riemens et al., 2007; Sooby et al., 2007). Organic farmers, out of necessity, rely heavily on soil tillage and other forms of labor-intensive soil cultivation for weed management despite the well-known disadvantages to soil health associated with intensive soil disturbance (Schonbeck and Morse, 2007; Morse and Creamer, 2006).

A small but growing number of organic farmers have begun adopting reduce tillage techniques, which blend the soil-conserving and labor-saving methods of conventional no-till systems with traditional soil building practices (i.e. cover cropping) used in organic production (Leavitt et al., 2011; Mirsky et al., 2013). In organic no-till, an *in situ* mulch is created by mechanically terminating mature cover crops. Subsequent cash crops are direct seeded or transplanted into the mulch-covered soil. The cover crop mulch manages weeds in place of mechanical cultivation through physical impedance, light interception, and allelopathy (Mohler and Teasdale, 1993; Teasdale and Mohler, 1993; Teasdale and Mohler, 2000).

Weed suppression in no-till systems is achieved with high biomass ( $>3000 \text{ kg ha}^{-1}$ ) cover crops (Mohler and Teasdale, 1993). Organic no-till research has focused primarily

on fall-planted, winter annual cover crops that establish quickly, are competitive with weeds during the winter and spring, produce large amounts of biomass, and are terminated easily using mechanical methods (Delate et al., 2012). Good results have been found using monocultures or combinations of cereal rye (*Secale cereale* L.), hairy vetch (*Vicia villosa* Roth), and crimson clover (*Trifolium incarnatum* L.) (Delate et al., 2012; Mischler et al., 2010; Duzy et al., 2014; Abdul-Baki et al., 1996).

Adequate termination of cover crops is essential to organic no-till (Schonbeck and Morse, 2007; Creamer and Dabney, 2002). Optimal results have been documented with specialized roller-crimping devices that crush cover crop tissue and press the intact crop residue onto the soil surface (Creamer and Dabney, 2002; Mischler, et al., 2010, Ashford and Reeves, 2003; Morse, 1999; Moyer, 2011). Mirsky et al. (2009), Mischler et al. (2010), and Ashford and Reeves (2003) found good control of roller-crimped no-till cover crops with lasting weed suppression.

Despite the demonstrated weed suppression of no-till mulches, organic no-till vegetable production systems have produced mixed results (Delate et al., 2012). Yield reductions associated with no-till mulches were documented in squash (Leavitt et al., 2011), bell pepper (Diaz-Perez et al., 2008; Leavitt et al., 2011), and tomato production (Leavitt et al., 2011). On the contrary, comparable or positive yield responses in organic no-till systems compared to conventionally tilled systems were reported in tomatoes (Abdul-Baki et al., 1996; Madden et al., 2004; Delate et al., 2012).



Common problems researchers have identified regarding organic no-till production include: sub-optimal soil temperatures early in the season and shortened degree growing days caused by the cooling effect of cover crop mulches; loss of earliness due to a lack of synchrony between cover crop maturity and optimal cash crop planting dates; N immobilization when using high C/N cover crops (i.e. rye); increased weed pressure particularly when cover crop stands are inadequate; and reduced N mineralization and poor N synchrony due to a lack of cover crop incorporation (Leavitt et al., 2011; Schonbeck and Morse, 2007; Schonbeck, 2015; Moyer, 2011; Creamer et al., 1997; Morse, 1999; Mirsky et al., 2011; Parr, et al., 2014).

The purpose of this research is to compare an organic no-till vegetable production system to a conventionally tilled system on a Toccoa sandy loam soil in Clemson, SC along the following parameters: vegetable yield, soil N dynamics, and weed management inputs.

## **Materials and Methods**

### Field Experiment

The experiment was initiated in October 2013 at the Clemson University Student Organic Farm, a 5-acre USDA certified organic farm on the Calhoun Field Research Area on the Clemson University campus. The soil at the study site is a moderately well drained Toccoa sandy loam (Coarse-loamy, mixed, active, nonacid, thermic Typic Udifluvents) with an average organic matter content of 4.6%. Although the experiment began in 2013,

observations were taken only from 2014 to 2015. The experimental design for both years was a 2 by 3 factorial randomized complete block design replicated three times. The treatments consisted of two levels of tillage [no-till (NT) and conventional tillage (CT)] of a cereal rye/crimson clover cover crop biculture and three levels of N fertilization (0, 58, and 116 kg ha<sup>-1</sup> N) for tomato and summer squash production. Conventional tillage was accomplished with a disk harrow to a depth of 15 cm. The recommended N fertilizer rate (116 kg ha<sup>-1</sup>) was based on Clemson Agricultural Service Lab soil fertility recommendations from a standard soil analysis of composited 0-15 cm soil samples taken in March 2014 from the year one experiment site. In 2014, soil at the experiment plot was amended with P (from 0-10-0 bone meal) at a rate of 448 kg ha<sup>-1</sup> and K (from 0-0-50 potash) at a rate of 60 kg ha<sup>-1</sup> according to soil test recommendations. No P and K amendments were needed in 2015. The treatments were arranged in a split-split plot design. Tillage treatment split plots (6 m x 7.5 m) were established for each vegetable crop with 2 m alleys between each plot. Alleys were flail mowed, tilled, and planted to a buckwheat (*Fagopyrum esculentum*) cover crop in both years. Split-plots were divided into three split-split plots (vegetable rows) spaced 1.5 m apart for the N fertilization treatment. The treatments were replicated three times for each vegetable crop.

On 4 October, 2013 and 13 September, 2014 experiment plots were seeded with a mixture of cereal rye (VNS) and crimson clover (VNS in 2013 and 'Dixie' in 2014) using a tractor-mounted overseeder attachment with 10 cm row spacing. Two-year cropping history for the 2014 experiment plot included maize (*Zea mays*) and summer squash cash crops and Japanese millet (*Echinochloa esculenta*), sunn hemp (*Crotalaria juncea*),

cereal rye, crimson clover, and cowpea (*Vigna unguiculata*) cover crops. The 2015 plot cropping history included garlic (*Allium sativum*), carrot (*Daucus carota*), cole crops (*Brassica oleracea*) and beet (*Beta vulgaris*) cash crops and cereal rye, crimson clover, cowpea, and sudex (*Sorghum bicolor* × *S. bicolor* var. *sudanese*) cover crops. Prior to cover crop seeding, the plots had been disked to remove weeds and level the field. In 2013, a seeding rate of 112 kg ha<sup>-1</sup> rye and 39 kg ha<sup>-1</sup> clover was used. The high rate of clover was due to calibration problems with the overseeder's seed shoot. In 2014, the same rate of rye was used, but the clover rate was reduced to a more appropriate rate of 12 kg ha<sup>-1</sup>. Cover crops in CT plots were flail mowed on 5 May, 2014 and 6 May, 2015 and the plots were disked repeatedly the following day in both years to incorporate cover crop residue. NT termination was accomplished on 6 May, 2014 and 11 May, 2015 with a rear-mounted 2.4 m I & J (Gap, PA) roller-crimper that had been filled with 225 kg of water for a total weight of 860 kg. The drum could have accommodated more water, but the tractor's lifting capacity was a limiting factor. In both years crimping was done in one direction with roughly 0.3 m of roller-crimper overlap with each pass. In 2014, the initial round of crimping did not fully terminate the rye crop, which by 7 May, had begun to rebound. The NT plots were re-crimped on 8 May with a small-plot 0.7 m roller-crimper mounted to a two-wheel walk-behind tractor. (The farm's larger category 1 tractor was not operational at the time.) An additional 113 kg of weight was added to the small-plot crimper for a total weight of 230 kg. In 2015, two back-to-back passes with the 2.4 m roller-crimper were made over the cover crop (same direction) on 11 May to ensure adequate rye termination. At time of crimping, the rye had reached Zadoks stage 69

(anthesis complete) in 2014 and stage 75 (medium milk) in 2015. Crimson clover maturity was not noted in either year.

Five-week old ‘Celebrity’ tomato and two-week old ‘Success’ squash seedlings were transplanted by hand in the corresponding CT and NT tomato and squash plots on 09 May, 2014 (both crops) and on 14 May (tomatoes) and 18 May (squash) in 2015. Vegetable plots consisted of three rows 4.5 m in length with 0.3 m spacing between plants and 1.5 m spacing between row centers. Drip irrigation was installed on top of mulch in NT and on bare soil in CT prior to transplanting; plants were watered immediately after transplantation. In the NT plots, the cover crop mulch was spread 12-15 cm apart by hand creating a narrow planting slit in the row prior to transplanting. The mulch was then pushed back against the plants after transplantation to cover the soil surface. All plants had been started in the farm’s greenhouse using on-farm generated potting soil [50% compost (2, 0.02, and 0.02 kg t<sup>-1</sup> N, P, and K, respectively): 25% perlite: 25% peat moss] with 120 ml lime and 710 ml powderized 8-5-5 feather meal fertilizer added per 0.3 m<sup>3</sup> of potting soil mix. Tomatoes were seeded in 128-count seed trays and then transplanted after 3 weeks into grow out pots measuring 10 cm in diameter. Squash were seeded in 72-count seed trays. Squash seedlings had reached the second true-leaf stage prior to transplantation. All vegetable transplants were fertilized with a powderized 8-5-5 feather meal fertilizer while in the greenhouse and were hardened off prior to transplantation. Immediately after transplantation, plants were fertilized according to N fertilization treatment with a side-dressed split application of a slower release, pelletized 13-0-0 feather meal fertilizer. A second split application of 13-

0-0 was made at flowering stage for each crop. Tomatoes were trellised using the “Florida weave” technique.

#### Cover Crop Data

Aboveground cover crop biomass was sampled in three 0.5m<sup>2</sup> quadrants in 2014 and five 0.5 m<sup>2</sup> quadrants in 2015 from the alleys between vegetable plots immediately prior to CT plot flail mowing. The biomass samples were oven dried for 72 hrs at 55°C and then weighed. Additionally, subsamples from each biomass sample were sent to the Clemson University Agricultural Service Lab where they were dried at 70-80°C for 12-24 hrs, ground to pass through a 2 mm sieve, and analyzed for total N by combustion using a LECO® FP528 Nitrogen Combustion Analyzer. Cover crop N contribution to the following vegetable crops was estimated as  $0.40 \times \text{total cover crop biomass} \times \text{cover crop \%N}$  (Baldwin and Creamer, 2006). Four weeks after termination, NT plots were assessed visually for percentage cover crop regrowth (Leavitt et al., 2011).

#### Weeding and Labor Inputs

CT plots were weeded approximately every 1-2 weeks in 2014 and approximately every 2-3 weeks in 2015. Weeding in tilled plots consisted of rototilling, flame weeding, and hand hoeing. NT plots were weeded every 2-3 weeks in both years. Weeding in NT plots consisted of rotary and string mowing weeds that had emerged through the cover crop residue and hand-pulling of perennial weeds. Weed management labor (hours) was recorded for each vegetable crop by tillage treatment for both years. Additionally, a

visual assessment of all NT plots was made 6 weeks after crimping to estimate average percent ground coverage by weeds (Creamer et al., 1997). Labor (hours) spent preparing no-till (crimping) and tilled (mowing + disking) cover crop plots prior to transplanting was also recorded both years.

### Soil Analysis

Prior to transplantation of vegetable crops and N fertilization (9 May, 2014 and 13 May, 2015), six 0-15 cm soil samples were taken from each of the 36 split-split plots. Soil samples were composited by split-split plot and subsamples were sent to the USDA-ARS Grassland Soil and Water Research Laboratory, Temple, TX for soil health analysis using the Soil Health Tool (SHT) ver. 4.4. (Haney, n.d.). Another series of soil samples, same protocol, were taken at the end of the each growing season (30 July, 2014 and 31 July, 2015). The samples were dried at 50°C, ground to pass through a 2 mm sieve, extracted with DI water and H<sup>3</sup>A, and analyzed on a Seal Analytical rapid flow analyzer for NO<sub>3</sub>-N and NH<sub>4</sub>-N (Haney et al., 2008). The water extract was analyzed on a Teledyne-Tekmar Apollo 9000 C:N analyzer for water-extractable organic C and total N and 40 g of each dried soil sample was re-wetted with DI water and incubated with a Solvita<sup>®</sup> paddle in a 237 ml glass jar for 24 hours (Haney et al., 2008). At the end of 24-hour incubation, the paddle was removed and placed in the Solvita<sup>®</sup> digital reader for CO<sub>2</sub>-C analysis. The SHT couples inorganic N (NO<sub>3</sub>-N and NH<sub>4</sub>-N), water-extractable organic C and N, and CO<sub>2</sub>-C measurements to estimate plant available N in the soil.

## Vegetable Yield

Yield data (weight) were recorded for marketable tomatoes (USDA grades 1-3) and marketable squash (USDA grades 1 and 2) in each row for every harvest (USDA 1997a, 1997b). Squash were harvested 3-4 times per week and tomatoes 2-3 times per week in both years.

## Tissue Mineral Analysis

In 2015, leaf tissue samples (excluding petioles) were taken from the most recently mature leaf of each plant in every row for both crops at early the early flowering stage. Samples were composited by row and sent to the Clemson Agricultural Service Lab where they were dried at 70-80°C for 12-24 hrs, ground to pass through a 2 mm sieve, and analyzed for total N by combustion using a LECO® FP528 Nitrogen Combustion Analyzer.

## Statistical Analysis

Statistical analyses were performed with analysis of variance (ANOVA) using the Fit Model procedure of JMP® (version 11.0) to determine the effects of tillage and N fertilization on vegetable yield and soil N. Fisher's least significant difference tests ( $P \leq 0.05$ ) were used to separate means. To compare inputs between the two tillage treatments, an analysis of labor was compiled by recording the total labor hours required to prepare the seedbed for planting and manage weeds during the vegetable growing season for each crop (Leavitt et al, 2011). Additionally, an ANOVA was performed to

determine the effects of tillage and N fertilization on vegetable yield per unit of labor.

## **Results**

### **Cover crops**

Cover crop biomass averaged 8,400 kg ha<sup>-1</sup> in 2014 and 8,960 kg ha<sup>-1</sup> in 2015. Based on the average total N content of the cover crop samples, 1.74% (2014) and 1.72% (2015), total cover crop N content was approximately 146 kg ha<sup>-1</sup> in 2014 and 154 kg ha<sup>-1</sup> in 2015. Total N contribution to the vegetable crops ( $0.40 \times \text{total cover crop biomass} \times \text{cover crop \%N}$ ) was 58 kg ha<sup>-1</sup> (2014) and 61 kg ha<sup>-1</sup> (2015). Roller-crimping provided adequate control of cover crops. Cover crop regrowth at 4 weeks after termination was minimal (<1%) in both years.

### **Weeding and Labor Inputs**

Seedbed preparation labor was higher both years in CT plots (Table 1). Mowing + disking required 191% and 300% more labor in 2014 and 2015, respectively, compared to NT roller-crimping (Table 1). Managing weeds was also more labor-intensive in CT plots. Weed labor was 400% and 338% greater in CT tomato and squash plots, respectively, compared to NT plots in 2014 (Table 1). In 2015, weed labor was 45% greater in CT tomato plots compared to NT; squash plot weed management was comparable between tillage treatments. NT cover crop plots did become weedy later in the season particularly in 2014 when regrowth of the previous summer's cover crop (Japanese millet) required routine mowing to keep the millet from overwhelming the NT



plots. Average percent ground cover by weeds at 6 weeks after termination in NT plots was 35% (tomatoes) and 25% (squash) in 2014 and 10% (tomatoes) and 10% (squash) in 2015.

## Soil Analysis

### *Pre-season analysis*

Average plant available N and total N were significantly greater in NT tomato plots compared to CT in 2014 (Table 2). There were no significant differences in available and total N between tillage treatments in 2015. There were no significant differences in CO<sub>2</sub>-C levels between tillage treatments in tomato plots in either year. Average plant available N and total N in squash plots were not statistically different between tillage treatments for either year studied (Table 2). Average squash plot CO<sub>2</sub>-C was significantly higher with NT in 2014 but similar between tillage treatments in 2015.

### *Post-season analysis*

Tomato plant available N and total N levels were similar both years regardless of tillage treatment (Table 3a). Average CO<sub>2</sub>-C was significantly higher in 2014 with NT but similar between CT and NT in 2015. Average plant available N and total N were significantly greater in NT squash plots in 2014 and in CT plots in 2015 (Table 3b). Average CO<sub>2</sub>-C was significantly higher in CT squash plots in 2015. Based on post-season soil analysis, N fertilization treatments did not significantly affect plant available N, total N, or CO<sub>2</sub>-C in either year for either crop studied.

## Tissue Mineral Analysis

Based on mid-season analysis, there appeared to be no effect of either tillage or fertilization on leaf tissue N. Average leaf tissue total N values were similar between tillage and N fertilization treatments for both crops in 2015 (Table 4). Average %N for both CT and NT were within the sufficiency ranges for both tomato (3.5-5.0%) and squash (4.0-6.0%) crops (Campbell, 2000).

## Vegetable Yield

Average NT tomato yields were significantly greater than CT yields in 2014; in 2015 CT yields were significantly greater than NT (Table 5). Squash yields were similar between tillage treatments for both years studied (Table 5). Nitrogen fertilization treatments did not significantly affect vegetable yield in either year for either crop studied nor were there significant tillage x fertilization interactions. Vegetable yields per unit of labor (seedbed preparation + weeding) were significantly greater in both NT crops in both years studied (Table 6).

## Discussion

The significant reduction in CT tomato yield in 2014 was likely due to disease. CT tomatoes were severely damaged by a combination of Southern blight (*Sclerotium rolfsii*) and Pythium root rot (*Pythium spp.*). Plant pathogen diagnosis was confirmed by the Clemson University Plant Problem Clinic. Both pathogens thrive in moist conditions found in poorly drained sites (Kluepfel et al., 2014). Because of the no-till component of

the study, we did not create raised beds in either tillage treatment. The roller-crimper, we found, provides optimal cover crop termination on level terrain, although there are roller-crimping devices designed for use in raised-bed systems (Reberg-Horton et al., 2012; Moyer, 2011). Normal farm CT practices include post-tillage raised-bed making to improve field drainage for cultural management of soil-borne diseases. In all, roughly 14% of the CT tomato plants in 2014 were lost to disease, which impacted average row yields. However, when row yield data were transformed from yield per row to yield per plant (by dividing row yields by number of plants per row) and analyzed using the same statistical model, there were no significant differences between tillage treatments (data not shown). NT tomatoes remained disease free in 2014, which was notable because: 1) CT and NT crops were grown in spatially similar parts of the field with identical cropping histories; 2) soil conditions with high levels of available carbon (i.e. poorly decomposed plant tissue) such as those found in reduced tillage systems are conducive to Southern blight (diagnostician's notes; Averre, 2009). A different field at the farm was used in the second year of the study and both CT and NT crops remained disease free.

Seedbed prep and weed management labor were much higher in CT plots in 2014 compared to 2015. Mowing and incorporating the cover crop residue took longer in 2014. More in-field tractor turns and repositioning were required in 2014 during mowing and subsequent tillage because of plot proximity to adjacent crop fields. Regarding 2015, roughly 75% of the cover crop at the study site (visual estimation) was lodged by severe rain and wind events three weeks prior to cover crop termination. At termination, approximately 20% of the cover crop stand in CT plots remained lodged – only 20% of

the crop was left standing at its original height. The lack of a fully erect cover crop made mowing less time intensive in 2015. Regarding in-season weeding, field conditions in 2014 were generally wetter in the first several weeks of the vegetable-growing season compared to 2015. In 2015, there were no rain events for the first 2.5 weeks after transplanting, which decreased the amount of early-season weeding that had to be done in CT plots after transplantation.

Overall, we found that the labor required to establish and manage weeds in CT plots was considerably greater when compared to NT. When yields were similar (squash, in both years) or even greater using CT practices (i.e. tomatoes in 2015), the management savings associated with NT translated to significantly greater yields per unit of labor input. Further, having a 5-7 week relatively weed-free window early in the growing season where little weeding was needed in newly transplanted crops – as was our experience with no-till – was particularly advantageous on a small, diversified farm such as ours where labor demands are high in the early summer growing season with 20+ crops growing in the fields at any given time.

N levels were high across all treatments each year. N fertility treatments had no significant effect on vegetable yield. Thus, soil N was likely not a limiting factor in either year probably as a result of high soil organic matter content and residual N from previous cover crops. Plant available N, total N, and CO<sub>2</sub>-C were generally higher in soil from the 2015 site.

Because of the short duration of the study, we were unfortunately unable to identify any discernible trends regarding differences in soil health across tillage treatments. Longer-term, mostly conventional agronomic-crop no-till studies, though, have demonstrated significant advantages to soil health (increased SOM) after multiple (>5) years of conventional no-tillage management (Arshad et al., 1990; Dick, 1983; Blevins et al., 1977; Bauer and Black, 1981; Edwards et al., 1992).

Regarding greater farmer adoption of organic no-till, one potential drawback we recognized is weediness later in the growing season, which we experienced particularly in year one of the study. Repeated hand weeding or “rouging” of no-till plots can be especially labor intensive and could negate the early season weed management savings realized with no-till mulches. We found adequate weed management later in the growing season using string and rotary mowing to keep emerged weeds in check but not necessarily controlled. Unrelated to this study, the Student Organic Farm has also experimented with adding off-farm mulches (e.g. leaf litter) to no-till mulches for longer season summer vegetable (e.g. pepper and eggplant) production in an attempt to compensate for decomposition of no-till mulches over the course of the growing season. Further research into cost-effective weed management strategies in organic no-till systems is warranted.

One main disadvantage we found using no-till was loss of earliness, which is an identified problem with spring/early-summer organic no-till vegetable cropping systems (Schonbeck and Morse, 2007). By prolonging the cover crop growing season until the rye

had matured to a stage where it could be managed by either tillage treatment, vegetables were not available for market until late June (squash) and mid-July (tomatoes). In order to produce tomatoes and squash for the early summer market period, the cover crops would had to have been terminated in early April, which would not have been compatible with no-till practices – the cover crop is too immature at this stage to be crimped and has not produced nearly enough biomass for effective no-till mulch. Thus, we found no-till vegetable production is perhaps best used for mid-to-late season summer crops. In future work, we would like to explore earlier fall planting/spring termination dates with faster maturing cover crop varieties to improve reduced tillage vegetable cropping earlier in the season.

### **Acknowledgements**

Special thanks to Dr. Rick Haney and the staff of the USDA-ARS Grassland, Soil and Water Research Laboratory, Temple, TX, for their assistance with the soil analyses. Also, thanks to farm manager Shawn Jadrnicek, Charles Pellett, and other Student Organic Farm staff who assisted with this project. This research was supported by the USDA SARE, Southern region, under award number GS13-126.

### **Literature Cited**

- Abdul-Baki, A., Stommel, J., Watada, A., Teasdale, J., and Morse, R. (1996). Hairy vetch mulch favorably impacts yield of processing tomato. *HortScience*, 31(338), 340.
- Arshad, M. A., Schnitzer, M., Angers, D. A., and Ripmeester, J. A. (1990). Effects of till vs no-till on the quality of soil organic matter. *Soil Biology and Biochemistry*, 22(5), 595-599.

- Ashford, D. L. and Reeves, D. W. (2003). Use of a mechanical roller-crimper alternative kill method for cover crops. *American Journal of Alternative Agriculture*, 18, 37-45.
- Averre, C. W. 2009. Southern blight of vegetable crops. *Vegetable Disease Information Note 9 (VDIN-009)*. North Carolina State University. Retrieved from: <http://www.ces.ncsu.edu/depts/pp/notes/oldnotes/vg9.htm>
- Baldwin, K. and Creamer, N. (2006). Cover crops for organic farms. North Carolina Cooperative Extension Service # E06-45788. Retrieved from: <http://content.ces.ncsu.edu/cover-crops-for-organic-farms.pdf>
- Bauer, A., and Black, A. L. (1981). Soil carbon, nitrogen, and bulk density comparisons in two cropland tillage systems after 25 years and in virgin grassland. *Soil Science Society of America Journal*, 45, 1166-1170.
- Blevins, R. L., Thomas, G. W., and Cornelius, P. L. (1977). Influence of no-tillage and nitrogen-fertilization on certain soil properties after 5 years of continuous corn. *Agronomy Journal*, 69(3), 383-386.
- Campbell, R. C. (2000). Sufficiency ranges for plant analysis in the southern region of the United States. *Southern Cooperative Series Bulletin #394*. Retrieved from: <http://www.ncagr.gov/agronomi/saaesd/scsb394.pdf>
- Clark, A. J., Decker, A. M., and Meisinger, J. J. (1994). Seeding rate and kill date effects on hairy vetch cereal rye cover crop mixtures for corn production. *Agronomy Journal*, 86(6), 1065-1070.
- Creamer, N. G, and Dabney, S. M. (2002). Killing cover crops mechanically: review of recent literature and assessment of new research results. *American Journal of Alternative Agriculture*, 17(01), 32-40.
- Creamer, N. G., Bennett, M. A., and Stinner, B. R. (1997). Evaluation of cover crop mixtures for use in vegetable production systems. *HortScience*, 32(5), 866-870.
- Delate, K., Cwach, D., and Chase, C. (2012). Organic no-tillage system effects on soybean, corn and irrigated tomato production and economic performance in Iowa, USA. *Renewable Agriculture and Food Systems*, 27, 49-59.
- Diaz-Perez, J., Silvoy, J., Phatak, S., Ruberson, J., and Morse, R. (2008). Effect of winter cover crops and no-till on the yield of organically grown bell pepper (*Capsicum annuum* L.). In R. Prange and S. Bishop (eds.). (Ed.), *Proc. XXVII IHC-S11 Sustainability through integrated and organic horticulture*. pp. 767.

- Dick, W. A. (1983). Organic-carbon, nitrogen, and phosphorus concentrations and pH in soil profiles as affected by tillage intensity. *Soil Science Society of America Journal*, 47(1), 102-107.
- Duzy, L. M., Kornecki, T. S., Balkcom, K. S., and Arriaga, F. J. (2014). Net returns and risks for cover crop use in Alabama tomato production. *Renewable Agriculture and Food Systems*, 29(4), 334-344.
- Edwards, J. H., Wood, C. W., Thurlow, D.L., and Ruf, M. E. (1992). Tillage and crop-rotation effects on fertility status of a Hapludult soil. *Soil Science Society of America Journal*, 56(5), 1577-1582.
- Haney, R. L., n.d. Soil health tool (SHT) ver 4.4: an integrated approach to soil testing. Retrieved from:  
[http://www.nrcs.usda.gov/wps/PA\\_NRCSCConsumption/download/?cid=nrcseprd333873&ext=pdf](http://www.nrcs.usda.gov/wps/PA_NRCSCConsumption/download/?cid=nrcseprd333873&ext=pdf)
- Haney, R. L., Brinton, W. H., and Evans, E. (2008). Estimating soil carbon, nitrogen, and phosphorus mineralization from short-term carbon dioxide respiration. *Communications in Soil Science and Plant Analysis*, 39(17-18), 2706-2720.
- Kluepfel, M., Blake, J. H., Keinath, A. P., and Williamson, J. (2014). Tomato diseases and disorders. HGIC #2217. Clemson University. Retrieved from:  
[http://www.clemson.edu/extension/hgic/pests/plant\\_pests/veg\\_fruit/hgic2217.html](http://www.clemson.edu/extension/hgic/pests/plant_pests/veg_fruit/hgic2217.html)
- Leavitt, M. J., Sheaffer, C. C., Wyse, D. L., and Allan, D. L. (2011). Rolled winter rye and hairy vetch cover crops lower weed density but reduce vegetable yields in no-tillage organic production. *HortScience*, 46(3), 387-395.
- Madden, N. M. et al. (2004). Evaluation of conservation tillage and cover crop systems for organic processing tomato production. *Horttechnology*, 14(2), 243-250.
- Mirsky, S. B. et al. (2013). Overcoming weed management challenges in cover crop-based organic rotational no-till soybean production in the eastern United States. *Weed Technology*, 27(1), 193-203.
- Mirsky, S. B., Curran, W. S., Mortensen, D. A., Ryan, M. R., and Shumway, D. L. (2009). Control of cereal rye with a roller/crimper as influenced by cover crop phenology. *Agronomy Journal*, 101(6), 1589-1596.
- Mirsky, S. B., Curran, W. S., Mortensen, D. M., Ryan, M. R., and Shumway, D. L. (2011). Timing of cover-crop management effects on weed suppression in no-till planted soybean using a roller-crimper. *Weed Science*, 59(3), 380-389.



- Mischler, R. A., Curran, W. S., Duiker, S. W., and Hyde, J. A. (2010). Use of a rolled-rye cover crop for weed suppression in no-till soybeans. *Weed Technology*, 24(3), 253-261.
- Morse, R. D. (1999). No-till vegetable production – its time is now. *Horttechnology*, 9(3), 373-379.
- Morse, R., and Creamer, N. (2006). Developing no-tillage systems without chemicals: The best of both worlds? In P. Kristiansen, A. Taji, & J. Reganold (Eds.), *Organic agriculture: a global perspective* (pp. 83-91). Ithaca, NY: Comstock Pub. Associates.
- Moyer, J. (2011). *Organic no-till farming*. Austin, TX: Acres U.S.A.
- Parr, M., Grossman, J. M., Reberg-Horton, S. C., Brinton, C., and Crozier, C. (2014). Roller-crimper termination for legume cover crops in North Carolina: impacts on nutrient availability to a succeeding corn crop. *Communications in Soil Science and Plant Analysis*, 45(8), 1106-1119.
- Reberg-Horton, S. C. et al. (2012). Utilizing cover crop mulches to reduce tillage in organic systems in the southeastern USA. *Renewable Agriculture and Food Systems*, 27(1), 41-48.
- Riemens, M., Groeneveld, R., Lotz, L., and Kropff, M. (2007). Effects of three management strategies on the seedbank, emergence and the need for hand weeding in an organic system. *Weed Research*, 47, 442-451.
- Schonbeck, M. W. (2015). What is "organic no-till," and is it practical? Extension Foundation, eOrganic Community of Practice. Retrieved from [https://www.extension.org/pages/18526/what-is-organic-no-till-and-is-it-practical#.VAuqLhar\\_No](https://www.extension.org/pages/18526/what-is-organic-no-till-and-is-it-practical#.VAuqLhar_No)
- Schonbeck, M. W. and Morse, R. D. (2007). Reduced tillage and cover cropping systems for organic vegetable production. Virginia Association of Biological Farming Info Sheet, 9-07. Retrieved from: <http://www.sare.org/Learning-Center/SARE-Project-Products/Southern-SARE-Project-Products/Reduced-Tillage-and-Cover-Cropping-Systems-for-Organic-Vegetable-Production>
- Sooby, J., Landeck, J., and Lipson, M. (2007). 2007 National Organic Research Agenda. Santa Cruz, CA.: Organic Farming Research Foundation. Retrieved from: <http://ofrf.org/sites/ofrf.org/files/docs/pdf/nora2007.pdf>

Teasdale, J.R. and Mohler, C. L. (1993). Light transmittance, soil temperature, and soil moisture under residue of hairy vetch and rye. *Agronomy Journal*, 85(3), 673-680.

Teasdale, J.R. and Mohler, C. L. (2000). The quantitative relationship between weed emergence and the physical properties of mulches. *Weed Science*, 48(3), 385-392.

USDA. 1997a. United States standards for grades of tomatoes. Effective 1 Oct. 1991. USDA-AMS, Washington, D.C.

USDA. 1997b. United States standards for grades of summer squash. Effective 6 Jan. 1984. USDA-AMS, Washington, D.C.

**Table 1. Total labor inputs by tillage treatment.**

<u>Tomato</u>	<u>Treatment</u>	<u>Seedbed preparation (hrs)</u>		<u>Weeding (hrs)</u>	
		<u>2014</u>	<u>2015</u>	<u>2014</u>	<u>2015</u>
	NT	0.9	0.3	1.3	1.1
	CT	2.75	1.2	6.5	1.6
<u>Squash</u>	<u>Treatment</u>	<u>2014</u>	<u>2015</u>	<u>2014</u>	<u>2015</u>
	NT	0.9	0.3	1.3	0.9
	CT	2.75	1.2	5.7	0.9

**Table 2. Effects of tillage (NT and CT) on average plant available N, total N, and CO<sub>2</sub>-C, pre-season<sup>x</sup> analysis.**

		Plant Available N (kg ha <sup>-1</sup> )		Total N (kg ha <sup>-1</sup> )		CO <sub>2</sub> -C (ppm)	
<b>Tomato</b>	<u>Tillage</u>	<u>0 DBT<sup>y</sup> 2014</u>	<u>1 DBT 2015</u>	<u>0 DBT 2014</u>	<u>1 DBT 2015</u>	<u>0 DBT 2014</u>	<u>1 DBT 2015</u>
	NT	61.36a <sup>z</sup>	94.98a	74.12a	102.57a	132.34a	242.57a
	CT	51.89b	100.07a	61.34b	113.44a	128.61a	189.50a
		<i>P</i> = 0.0222	<i>P</i> = 0.5483	<i>P</i> = 0.0197	<i>P</i> = 0.2731	<i>P</i> = 0.7757	<i>P</i> = 0.2335
<b>Squash</b>	<u>Tillage</u>	<u>0 DBT 2014</u>	<u>4 DBT 2015</u>	<u>0 DBT 2014</u>	<u>4 DBT 2015</u>	<u>0 DBT 2014</u>	<u>4 DBT 2015</u>
	NT	65.21a	90.60a	78.56a	99.99a	65.01a	217.76a
	CT	54.97a	91.78a	65.92a	103.84a	44.21b	172.32a
		<i>P</i> = 0.0699	<i>P</i> = 0.8377	<i>P</i> = 0.0846	<i>P</i> = 0.5761	<i>P</i> = 0.0493	<i>P</i> = 0.3342

x

After cover crop termination but before vegetable transplantation.

<sup>y</sup> DAT=Days after transplantation.

<sup>z</sup> Mean separation within tillage and year by Fisher's least significant difference test,  $P \leq 0.05$ .

**Table 3a. Effects of tillage (NT and CT) and N fertilization (0, 58, and 116 kg ha<sup>-1</sup>) on average plant available N, total N, and CO<sub>2</sub>-C, post-season<sup>x</sup> analysis.**

<b>Tomato</b>	<b>Tillage</b>	<b>Plant Available N (kg ha<sup>-1</sup>)</b>		<b>Total N (kg ha<sup>-1</sup>)</b>		<b>CO<sub>2</sub>-C (ppm)</b>	
		<b>82 DAT<sup>y</sup> 2014</b>	<b>78 DAT 2015</b>	<b>82 DAT 2014</b>	<b>78 DAT 2015</b>	<b>82 DAT 2014</b>	<b>78 DAT 2015</b>
	NT	68.80a	89.48a	78.14a	104.03a	161.56a <sup>z</sup>	315.98a
	CT	59.99a	89.41a	67.22a	104.76a	97.29b	250.40a
	<b>Fertilization</b>						
	0	69.45a	84.62a	78.11a	98.45a	94.83b	272.02a
	58	61.26a	88.83a	69.23a	103.97a	162.67a	278.43a
	116	62.57a	94.88a	70.69a	110.77a	130.77ab	299.12a
	Tillage	<i>P</i> = 0.1383	<i>P</i> = 0.9925	<i>P</i> = 0.1441	<i>P</i> = 0.9358	<i>P</i> = 0.0260	<i>P</i> = 0.0823
	Fert	<i>P</i> = 0.4601	<i>P</i> = 0.5047	<i>P</i> = 0.5495	<i>P</i> = 0.5366	<i>P</i> = 0.1288	<i>P</i> = 0.7972
	Tillage x Fert	<i>P</i> = 0.7432	<i>P</i> = 0.4997	<i>P</i> = 0.7638	<i>P</i> = 0.5544	<i>P</i> = 0.4815	<i>P</i> = 0.4051

<sup>x</sup> After all harvesting completed for the season.

<sup>y</sup> DAT=Days after transplantation.

<sup>z</sup>Mean separation within tillage or fertilization and within year by Fisher's least significant difference test, *P* ≤ 0.05.

**Table 3b. Effects of tillage (NT and CT) and N fertilization (0, 58, and 116 kg ha<sup>-1</sup>) on average plant available N, total N, and CO<sub>2</sub>-C, post-season<sup>x</sup> analysis.**

<b>Squash</b>	<b>Tillage</b>	<b>Plant Available N (kg ha<sup>-1</sup>)</b>		<b>Total N (kg ha<sup>-1</sup>)</b>		<b>CO<sub>2</sub>-C (ppm)</b>	
		<b>82 DAT 2014</b>	<b>74 DAT 2015</b>	<b>82 DAT 2014</b>	<b>74 DAT 2015</b>	<b>82 DAT 2014</b>	<b>74 DAT 2015</b>
	NT	70.92a	80.56b	82.31a	93.89b	180.94a	202.80b
	CT	59.88b	94.26a	67.89b	112.06a	130.67a	305.19a
	<b>Fertilization</b>						
	0	65.84a	91.47a	75.39a	106.78a	154.97a	253.68a
	58	64.38a	86.85a	73.79a	103.53a	171.05a	245.22a
	116	65.98a	83.90a	76.12a	98.60a	141.39a	263.08a
	Tillage	<i>P</i> = 0.0239	<i>P</i> = 0.0099	<i>P</i> = 0.0232	<i>P</i> = 0.0138	<i>P</i> = 0.0843	<i>P</i> = 0.0105
	Fert	<i>P</i> = 0.9413	<i>P</i> = 0.3880	<i>P</i> = 0.9372	<i>P</i> = 0.5629	<i>P</i> = 0.6638	<i>P</i> = 0.9055
	Tillage x Fert	<i>P</i> = 0.8259	<i>P</i> = 0.3235	<i>P</i> = 0.8481	<i>P</i> = 0.4375	<i>P</i> = 0.5631	<i>P</i> = 0.9274

<sup>x</sup> After all harvesting completed for the season.

<sup>y</sup> DAT=Days after transplantation.

<sup>z</sup>Mean separation within tillage or fertilization and within year by Fisher's least significant difference test, *P* ≤ 0.05.

**Table 4. Effects of tillage (NT and CT) and N fertilization (0, 58, 116 kg ha<sup>-1</sup>) on average leaf tissue N (2015).**

		<b>Total N (kg ha<sup>-1</sup>)</b>
<b><u>Tomato</u></b>	<u>Tillage</u>	
	NT	4.18a <sup>z</sup>
	CT	4.01a
	<u>Fertilization</u>	
	0	4.09a
	58	4.14a
	116	4.05a
	Tillage	<i>P</i> = 0.2710
	Fert	<i>P</i> = 0.8676
	Tillage x Fert	<i>P</i> = 0.3072
<b><u>Squash</u></b>	<u>Tillage</u>	
	NT	4.95a
	CT	4.56a
	<u>Fertilization</u>	
	0	4.76a
	58	4.82a
	116	4.67a
	Tillage	<i>P</i> = 0.0853
	Fert	<i>P</i> = 0.8200
	Tillage x Fert	<i>P</i> = 0.7217

<sup>z</sup>Mean separation within tillage or fertilization by Fisher's least significant difference test, *P* ≤ 0.05.

**Table 5. Effects of tillage (NT and CT) and N fertilization (0, 58, 116 kg ha<sup>-1</sup>) on average marketable vegetable yields per row.**

		<b>Yield (kg)</b>	
<b><u>Tomato</u></b>	<u>Tillage</u>	<u>2014</u>	<u>2015</u>
	NT	13.74a <sup>z</sup>	14.79b
	CT	9.27b	21.55a
	<u>Fertilization</u>		
	0	10.91a	15.89a
	58	10.60a	18.85a
	116	13.02a	19.80a
	Tillage	<i>P</i> = 0.0440	<i>P</i> = 0.0029
	Fert	<i>P</i> = 0.5609	<i>P</i> = 0.2019
	Tillage x Fert	<i>P</i> = 0.8515	<i>P</i> = 0.5212
<b><u>Squash</u></b>	<u>Tillage</u>	<u>2014</u>	<u>2015</u>
	NT	30.37a	23.32a
	CT	32.21a	25.14a
	<u>Fertilization</u>		
	0	27.56a	22.46a
	58	32.79a	26.45a
	116	33.52a	23.78a
	Tillage	<i>P</i> = 0.5459	<i>P</i> = 0.4830
	Fert	<i>P</i> = 0.2459	<i>P</i> = 0.4404
	Tillage x Fert	<i>P</i> = 0.9747	<i>P</i> = 0.7050

<sup>z</sup>Mean separation within tillage or fertilization and within year by Fisher's least significant difference test, *P* ≤ 0.05.



**Table 6. Effects of tillage (NT and CT) on average marketable vegetable yield per unit of effort.**

		<b>Yield per Unit of Effort<sup>y</sup> (kg/hr)</b>	
<b><u>Tomato</u></b>	<b><u>Tillage</u></b>	<b><u>2014</u></b>	<b><u>2015</u></b>
	NT	57.26a <sup>z</sup>	92.46a
	CT	9.00b	69.52b
		P < 0.0001	P = 0.0371
<b><u>Squash</u></b>	<b><u>Tillage</u></b>	<b><u>2014</u></b>	<b><u>2015</u></b>
	NT	126.54a	179.40a
	CT	34.27b	109.29b
		P < 0.0001	P = 0.0003

<sup>y</sup> Rate determined by dividing labor input totals for each tillage treatment by number of rows (9) to determine hours of labor input per row of crop by tillage treatment. Crop row yields were then divided by hours of labor per row to determine yield per unit of effort rate for each row.

<sup>z</sup>Mean separation within tillage and year by Fisher's least significant difference test,  $P \leq 0.05$ .