



University of Tennessee, Knoxville
Trace: Tennessee Research and Creative Exchange

Faculty Publications and Other Works -- Large
Animal Clinical Sciences

Veterinary Medicine -- Faculty Publications and
Other Works

4-2011

Updates on Crias

Christine Cocquyt
University of Tennessee - Knoxville

Follow this and additional works at: http://trace.tennessee.edu/utk_largpubs

Recommended Citation


Cocquyt, Christine, "Updates on Crias" (2011). *Faculty Publications and Other Works -- Large Animal Clinical Sciences*.
http://trace.tennessee.edu/utk_largpubs/21

This Conference Proceeding is brought to you for free and open access by the Veterinary Medicine -- Faculty Publications and Other Works at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Faculty Publications and Other Works -- Large Animal Clinical Sciences by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Updates on Crias

Dr. Christine Cocquyt
Large Animal Medicine Resident
University of Tennessee

The Newborn Cria



The Basics

- ⦿ Normal gestation length
- ⦿ Normal birth weights

Alpacas

>5.5 kg
Ave 7 kg

Llamas

>7 kg
Ave 9 kg


- ⦿ May lose up to 0.5 pounds in first day
- ⦿ Should gain 0.5-1 pound per day after

Cria Birth Weights

- Study in South American alpacas
- Trend of increasing birth weight as dam's age increased from 3 to 9 years of age
- Declined after 12 years of age
- May not apply to more intensively managed alpacas in US

Signs of Prematurity

- ◉ Weak, unable to hold head up, lateral
- ◉ Low birth weight
- ◉ Weak/absent suckle
- ◉ Floppy ears
- ◉ Silky coat
- ◉ Tendon laxity
- ◉ Incisors not erupted
- ◉ Soft hooves




Prematurity-related Concerns

- ◎ Failure of passive transfer
 - > Delayed or absent colostrum intake
 - > Decreased absorption of Ig
- ◎ Abnormal lung function
 - > Low oxygen to tissues
- ◎ Low blood sugar
- ◎ Dehydration
- ◎ Poor thermoregulation


Colostrum Antibodies

- Hypogammaglobulinemic
- Epitheliochorio placenta prevents Ig transfer to fetus → require colostrum intake
- Starts to close after first feeding; complete by 24 hours
- Mechanism of transfer may be impaired in premature crias



Fowler, ME. Medicine and Surgery of Camelids. 2010

IgG Concentration




Fowler, ME. Medicine and Surgery of Camelids. 2010

- Healthy crias may be okay with low [IgG]
- Peak [IgG] between 1-2 days of age
- Studies in Peru
 - Lower [IgG] in crias that died
 - Highest [IgG] when first suckle 2-4 hours after birth
 - Marginal [IgG] when first suckle 6-8 hours after birth

Failure of Passive Transfer

- Gold standard = radial immunodiffusion test (RID)
 - Measures [IgG]
 - 36 hours post birth
 - 24 hours to get result
 - Ideal >1000 mg/dl
- TP < 4.5 g/dl indicative
> 5.5 g/dl adequate
- Sodium sulfite precipitation test indicative
- Variable findings on other tests




<http://sciencera.com/biology/immunology-laboratory-procedures/>

Plasma Transfusion

- Oral – only effective soon after birth
 - Decreased efficacy in premies
- Intraperitoneal
 - Field veterinarians, breeders
 - Rapid infusion (5-10 min)
 - Risk of pain from distention, organ laceration, peritonitis
- Intravenous
 - 20-30 ml/kg over 1.5-3 hours
 - Variable increase in IgG levels
 - Recommend 2 units (approx 700 ml high concentration llama plasma) (unpublished, UT, 2009)



Supportive Care



- ⦿ IV catheter
- ⦿ Plasma!!!
- ⦿ IV fluid support – add dextrose
- ⦿ Antibiotics
- ⦿ Heat lamp
- ⦿ Intranasal oxygen
- ⦿ Feeding tube

Flexural Deformities

- ⦿ Premature or dysmature crias
- ⦿ Laxity in tendons
- ⦿ Fetlocks drop
 - Controlled exercise
- ⦿ May splay
 - Use hobbles
- ⦿ Good prognosis with time



Umbilicus

- ◎ 2-3 inches long is ideal
- ◎ Dip with dilute chlorhexidene or betadine solution every 12 hours until dried up
- ◎ Best to leave unclamped
 - > If clamp, 10-15 minutes recommended
 - > Increased risk of abscess with clamp
- ◎ Heat, pain, swelling, or discharge may indicate infection
- ◎ Route of infection to whole body!!!

Topics in the News

- ◉ Choanal Atresia
- ◉ Vitamin D toxicosis
- ◉ Sodium phosphate enemas
- ◉ Bovine viral diarrhea virus
- ◉ Coronavirus


Choanal Atresia

Choanal Atresia

- Congenital defect
- One or both sides
- Complete or partial

● Signs

- > Difficulty breathing
- > Weak
- > Aspiration possible
- > Other defects?



What can be done?

- ◎ Diagnosis
 - > Contrast radiographs
 - > Computed tomography
- ◎ Surgical repair under general anesthesia only option to correct
- ◎ Prognosis guarded to poor

Choanal Atresia Gene

- Genetic component suspected
- Similar to condition in humans (CHD7 gene)
- Researchers at University of Minnesota found gene CHD7 in affected alpaca cria
- Working on sequencing and then evaluating for mutations
- Eventually may have genetic test for breeding stock

Vitamin D Intoxication

Acute Renal Failure and Anuria Associated with Vitamin D Intoxication in Two Alpaca (*Vicugna pacos*) Cria

C. Gerspach et al. 2010

Journal of Veterinary Internal Medicine. 24(2), 443–449.

Vitamin D



- ◉ Converted in skin by UV radiation
- ◉ SA mountain habitat → high UV exposure
 - › Reduced converting ability
- ◉ Less UV exposure in North America
 - › Important to supplement
- ◉ Recommended
 - › SAC: 30 IU/kg BW (Van Saun)
 - › single dose 1000-2000 IU/kg
 - › Other species: 6.6 IU/kg

Vitamin D Functions

- ◎ Calcium and phosphorus metabolism
 - › Skeletal development
 - › Bone mineralization

- ◎ Deficiency → rickets




Too much of a good thing...

- ◎ Two alpaca crias
 - > 18-day-old male: 3,750 IU/kg/d for 7 days
 - Powdered goat colostrum supplement
 - > 8-day-old male: 12,987 IU/kg/d for 5 days
 - ADE paste → 100,000 IU vitamin D/day
- ◎ Signs
 - > Anorexia
 - > Weakness
 - > Depression

Vitamin D Intoxication

- ⦿ High calcium
- ⦿ High phosphorus
- ⦿ Kidney failure
- ⦿ Mineralization of organs



Vitamin D in supplements

Supplement	Vitamin D
Manna Pro Colostrum Powder	5000 IU/lb
MannaPro Kid Colostrum	5000 IU/lb
Kaeco Colostrum Powder	2750 IU/tsp
LaBelle Alpaca/Llama Powder	3200 IU/lb
ADE paste	varies

Recommended amount of 30 IU/kg:

7 kg alpaca cria = 210 IU/day

9kg llama cria = 270 IU/day

Enemas

Inadvertent transvaginal administration of sodium
phosphate enemas in 2 alpaca crias

Bragg et al. 2010.
JVECC 20(6), 623-627

Sodium Phosphate Enemas

- ⦿ 2 female alpaca crias
- ⦿ Given sodium phosphate enemas for suspected meconium impactions
- ⦿ Inadvertently given in reproductive tract
→ vaginal tear → enema fluid in abdomen
- ⦿ Electrolyte levels altered
- ⦿ One euthanized, one recovered

Enemas

- Meconium should pass w/in 18-20 hours of birth
- Sodium phosphate
 - Dogs < 25 lbs 2 fluid ounces toxic
- Warm soapy water



- Use only if indicated
- Do not repeat
- Ensure proper restraint
- Gentle applicator and lubrication



Bovine Viral Diarrhea Virus

Disseminated Bovine viral diarrhea virus in a persistently
infected alpaca cria

Byers et al. 2009

J Vet Diag Invest 21, 145-148.

Bovine Viral Diarrhea Virus

- ◎ Emerging disease in North American camelids?
- ◎ Study of 63 herds (voluntary)
 - > 25% of herds had positive animals
 - > 6% of herds had PI crias
- ◎ Cattle
 - > Acute → fever, low WBC, anorexic
 - > Persistent infection

BVDV-1b in Camelids

Natural transient infection

Decreased feed intake, mild lethargy

>60 days

Acute BVD

Mild fever, anorexia, lethargy

Intro of PI
→ 9/52 sick

Embryonic loss

Pregnancy loss

Abortion

Premature births

PI cria

Acute

chronic

What is a PI cria?

- ◎ **P**ersistantly **I**nfected
- ◎ Dam infected during pregnancy
 - › Virus crosses placenta
- ◎ Fetus infected **before** immune system develops
 - › Probably first trimester (cattle < 145 days)
 - Immune system accepts virus as “self”
 - Virus multiplies
- ◎ Often low birth weights, poor-doers, chronic respiratory and GI infections

Disseminated Bovine viral diarrhoea virus in a persistently infected alpaca (*Vicugna pacos*) cria

- ◎ Case report from Washington State
- ◎ 4-month-old male alpaca cria
 - › Negative antibody test
 - › Positive PCR and virus isolation tests
- ◎ Decreased weight gain, poor-doer, anorexia
- ◎ BVDV-1b found in many tissues
 - › Salivary glands → transmission through spit?
 - › Testes, prostate → reproductive transfer?
 - › Kidneys → urine?

Recommendations

- ◎ Avoid comingling especially pregnant females
 - > Isolate new animals until tested
- ◎ Test all crias
 - > PCR
- ◎ Ab test (ELISA)
 - > Bovine colostrum may give false (+)
 - > Vaccination
 - > 0.9-25.4 % positive
- ◎ PCR screen
 - > 10% of herd or 15 animals
 - > Retest positive in 3-4 weeks (transient up to 60 d)
 - > Pooled sample

Coronavirus

Identification of a novel coronavirus possibly associated with acute respiratory syndrome in alpacas in California

Crossley et al. 2010.
J Vet Diag Invest 22, 94-97.

Coronavirus

- ◎ First identified in NWC in 1998
 - > Herd outbreaks
 - > Severe diarrhea
- ◎ Diarrhea outbreaks – all ages
 - > Diagnosis with electron microscopy of feces
- ◎ “Snots” – outbreaks in 2007



Acute Respiratory Syndrome

- ◎ California 2007
- ◎ Fever, Mild flu signs to fatalities
 - > Severe fluid in lungs and chest cavity
 - > Fibrin in alveoli
- ◎ Isolated coronavirus from 1 lung
 - > Different than diarrheal virus
 - > More work needed to identify in respiratory cases

Questions?



Sources

- Bravo et al. (2009). Cria alpaca body weight and perinatal survival in relation to age of the dam. *Animal Reproduction Scienc* 111, 214-219.
- Davis et al. (1997) Seasonal effects on gestation length and birth weight in alpacas. *Animal Reproduction Scienc* 46, 297-303.
- Whitehead, Claire. (2009). Neonatal diseases in llamas and alpacas. *Vet Clin Food Anim* 25, 367-384.
- Whitehead, Claire. (2009) Management of neonatal llamas and alpacas. *Vet Clin Food Anim* 25, 353-366.
- C. Gerspach et al. (2010) Acute Renal Failure and Anuria Associated with Vitamin D Intoxication in Two Alpaca (*Vicugna pacos*) Cria. *Journal of Veterinary Internal Medicine*. 24(2), 443-449.

Sources (cont.)

- Bragg et al (2010) Inadvertent transvaginal administration of sodium phosphate enemas in 2 alpaca crias. JVECC 20(6), 623-627.
- Byers et al. (2009) Disseminated Bovine viral diarrhea virus in a persistently infected alpaca cria. J Vet Diag Invest 21, 145-148.
- Van Amstel, S and Kennedy, M. (2010) Bovine viral diarrhea infections in new world camelids-A review. Small Ruminant Research 91, 121-126.
- Topliff et al (2009) Prevalence of bovine viral diarrhea virus infections in alpacas in the United States. JAVMA 234(4), 519-529.
- Crossley et al (2010). Identification of a novel coronavirus possibly associated with acute respiratory syndrome in alpacas in California. J Vet Diag Invest 22, 94-97.