



August 2002

Molecular Cloning and Characterization of a Novel Mouse Macrophage C-type Lectin, mMGL2, Which Has a Distinct Carbohydrate Specificity from mMGL1

Thandi M. Onami

University of Tennessee - Knoxville, tonami@utk.edu

M. Tsuiji

University of Tokyo

M. Fujimori

University of Tokyo

Y. Ohashi

University of Tokyo

N. Higashi

University of Tokyo

Follow this and additional works at: https://trace.tennessee.edu/utk_micrpubs



Part of the [Immune System Diseases Commons](#), [Immunity Commons](#), [Immunology of Infectious Disease Commons](#), [Immunopathology Commons](#), [Microbiology Commons](#), and the [Virus Diseases Commons](#)

[Commons](#)

Recommended Citation

Onami, Thandi M.; Tsuiji, M.; Fujimori, M.; Ohashi, Y.; Higashi, N.; Hendrick, S. M.; and Irimura, T., "Molecular Cloning and Characterization of a Novel Mouse Macrophage C-type Lectin, mMGL2, Which Has a Distinct Carbohydrate Specificity from mMGL1" (2002). *Microbiology Publications and Other Works*. https://trace.tennessee.edu/utk_micrpubs/28

This Article is brought to you for free and open access by the Microbiology at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Microbiology Publications and Other Works by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Authors

Thandi M. Onami, M. Tsuiji, M. Fujimori, Y. Ohashi, N. Higashi, S. M. Hendrick, and T. Irimura

Molecular Cloning and Characterization of a Novel Mouse Macrophage C-type Lectin, mMGL2, Which Has a Distinct Carbohydrate Specificity from mMGL1*

Received for publication, April 18, 2002, and in revised form, May 16, 2002
Published, JBC Papers in Press, May 16, 2002, DOI 10.1074/jbc.M203774200

Makoto Tsuiji, Mayuko Fujimori‡, Yoshimi Ohashi, Nobuaki Higashi, Thandi M. Onami§, Stephen M. Hedrick§, and Tatsuro Irimura¶

From the Laboratory of Cancer Biology and Molecular Immunology, Graduate School of Pharmaceutical Sciences, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan and the §Cancer Center and the Division of Biology, University of California, San Diego, La Jolla, California 92093-0687

A novel mouse macrophage galactose-type C-type lectin 2 (mMGL2) was identified by BLAST analysis of expressed sequence tags. The sequence of mMGL2 is highly homologous to the mMGL, which should now be called mMGL1. The open reading frame of mMGL2 contains a sequence corresponding to a type II transmembrane protein with 332 amino acids having a single extracellular C-type lectin domain. The 3'-untranslated region included long terminal repeats of mouse early transposon. The *Mgl2* gene was cloned from a 129/SvJ mouse genomic library and sequenced. The gene spans 7,136 base pairs and consists of 10 exons, which is similar to the genomic organization of mMGL1. The reverse transcriptase-PCR analysis indicates that mMGL2 is expressed in cell lines and normal mouse tissues in a macrophage-restricted manner, also very similar to that of mMGL1. The mMGL2 mRNA was also detected in mMGL1-positive cells, which were sorted from thioglycollate-induced peritoneal cells with a mMGL1-specific monoclonal antibody, LOM-8.7. The soluble recombinant proteins of mMGL2 exhibited carbohydrate specificity for α - and β -GalNAc-conjugated soluble polyacrylamides, whereas mMGL1 preferentially bound Lewis X-conjugated soluble polyacrylamides in solid phase assays. These two lectins may function cooperatively as recognition and endocytic molecules on macrophages and related cells.

Macrophages (MØs)¹ and related cells are widely distributed throughout the body, displaying a morphological and functional diversity. They are found in the lymphoid organs, liver,

lungs, gastrointestinal tract, central nervous system, serous cavities, bones, synovia, and skin. Resident MØs mediate clearance of senescent or apoptotic cells, produce and secrete cytokines, are involved in hemopoiesis and bone resorption, transport and present antigens, and regulate neuroendocrine processes. Activated MØs are recruited to sites of infection, tissue injury, inflammation, and neoplasia and play crucial roles in tissue repair and pathogenesis (1).

The distribution and functional heterogeneity of MØs derive in part from their specialized plasma membrane receptors (2). Cell surface markers such as F4/80, sialoadhesin, MØ mannose receptor and scavenger receptor type A have significantly contributed to the current understanding of MØ ontogeny and function (3). However, in comparison to other immune cells such as B and T lymphocytes, relatively few MØ-restricted cell surface molecules have been identified. The physiological and pathological roles of these putative markers remain unknown.

Protein-carbohydrate interactions serve a variety of functions in the immune system. A number of lectins (carbohydrate-binding proteins) mediate both pathogen recognition and cell-to-cell interactions using structurally related carbohydrate recognition domains (CRDs). One of the most diversified families of these CRDs is Ca²⁺-dependent and termed C-type CRDs (4, 5). MØs and related cells such as dendritic cells are known to express several subfamilies of C-type lectins: type 1 multi-lectins such as MØ mannose receptor and lectins having type II transmembrane configurations such as MØ galactose-type (known previously as galactose/*N*-acetylgalactosamine-specific) C-type lectin (MGL). These lectins seem to mediate carbohydrate-specific endocytosis (6–11). Little is known regarding the expression of selectins, another subfamily of C-type lectins, in MØs and related cells.

Previously, MGL was cloned from rats, mice, and humans and characterized. The lectin from mice was shown to have specific affinity for highly branched *N*-linked carbohydrate chains with terminal β -galactosyl groups and glycopeptides carrying three consecutive α -GalNAc-Ser/Thr (Tn-antigens) (12). A human MGL was also shown to have affinity for glycopeptides carrying three consecutive Tn-antigens (13). Surface

* This work was supported by grants-in-aid from the Ministry of Education, Science, Sports and Culture of Japan (09254101, 11557180, 11672162, and 12307054), from the Research Association for Biotechnology, Special Coordination Funds for Promoting Science and Technology of the Ministry of Education, Culture, Sports, Science and Technology, and from the Program for Promotion of Basic Research Activities for Innovative Biosciences. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

The nucleotide sequence(s) reported in this paper has been submitted to the DDBJ/GenBank™/EBI Data Bank with accession number(s) AY103461, AY103462.

‡ Present address: Dept. of Applied Biological Chemistry, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, 113-8657, Japan.

¶ To whom correspondence should be addressed. Tel.: 813-5841-4870; Fax: 813-5841-4879; E-mail: irimura@mol.f.u-tokyo.ac.jp.

¹ The abbreviations used are: MØ, macrophage; ABTS, 2,2'-amino-bis(3-ethylbenzthiazoline-6-sulfonic acid); ASGR, hepatic asialoglyco-

protein receptor; BAC, bacterial artificial chromosome; BSA, bovine serum albumin; CRD, carbohydrate-recognition domain; DPBS, Dulbecco's phosphate-buffered saline; HRP, horseradish peroxidase; MGL, macrophage galactose-type C-type lectin; RHL, rat hepatic lectin; RT, reverse transcriptase; TG-PEC, thioglycollate-induced peritoneal cells; RACE, rapid amplification of cDNA ends; STAT, signal transducers and activators of transcription; mAb, monoclonal antibody; MES, 4-morpholineethanesulfonic acid; MOPS, 4-morpholinepropanesulfonic acid; h, human; m, mouse; r, rat.

plasmon resonance revealed that affinity of recombinant hMGL for immobilized glycopeptides increased in parallel with the number of GalNAc residues (14). Tn-antigen was known as a marker of malignant cells, and MGL was shown to play a role as a recognition molecule on MØs for tumor cells (15–17). Immunohistochemical localization of mouse MGL (mMGL) with specific monoclonal antibodies revealed that this lectin has a strong association with MØs residing in connective tissue and those infiltrated into tumor tissues (18–20). Recent studies demonstrated that this lectin is also expressed on the surface of immature dendritic cells and is involved in the uptake of glycosylated antigens in mice and humans (21, 22).

A C-type lectin highly homologous to MGL and expressed mainly on hepatocytes, the hepatic asialoglycoprotein receptor (ASGR), has two isomers in mice, rats, and humans, apparently due to recent gene duplication. Previously, MGL was believed to have a single gene. Southern blotting analysis supported this notion. However, in this report, we describe the properties of another novel MØ galactose-type C-type lectin, mMGL2. As an obvious consequence, the previous mMGL must now be called mMGL1. These lectins are highly homologous to each other except in their cytoplasmic domains and CRDs. We found that mMGL2 has a distinct carbohydrate specificity from mMGL1. These lectins seem to be expressed on the same cells and therefore seem to function cooperatively.

EXPERIMENTAL PROCEDURES

Cells—The following cell lines were provided by Cell Resource Center for Biomedical Research, Institute of Development, Aging and Cancer Tohoku University, Sendai, Japan: L929, JLS-V9, EL4, RL δ 1, YAC-1, BCL1-B20, P815, P388, and M1 cells. RAW264.7 cells were purchased from ATCC. All cells were cultured in RPMI 1640 media with 10% fetal calf serum at 37 °C with 5% CO₂.

Thioglycollate-induced peritoneal cells (TG-PEC) were obtained from 6–8-week old female C57BL/6J mice from Clea Japan, Inc (Kawasaki, Japan). These were maintained under pathogen-free conditions. These mice received 1 ml of 4% thioglycollate broth (Difco, Detroit, MI) by intraperitoneal injection. Four days later, mice were sacrificed by neck dislocation, and peritoneal exudate cells were harvested by lavage with 5 ml of RPMI 1640 media on ice.

TG-PEC were suspended in chilled 0.1% BSA/Dulbecco's modified phosphate-buffered saline (DPBS: 137 mM NaCl, 13.4 mM KCl, 40.5 mM Na₂HPO₄, 7.35 mM KH₂PO₄, 0.49 mM MgCl₂, 0.905 mM CaCl₂) containing phycoerythrin-labeled mAb LOM-14 (reactive with mMGL1 and mMGL2 as described below) and biotin-labeled mAb LOM-8.7 (reactive with mMGL1) and then incubated with streptavidin-fluorescein isothiocyanate (Zymed Laboratories Inc., South San Francisco, CA) on ice. The cells were sorted for the expression of mMGL1 (mAb LOM-8.7 staining) by Epics Elite (Beckman Coulter).

Mice—*Mgl1*^{-/-} mice were prepared in Hedrick's laboratory at the University of California, San Diego, CA. The details will be published in a separate study (23).

RNA Preparations and RT-PCR Analysis—Total RNAs were extracted by using Ultraspec RNA zol (Biotecx, Houston, TX) according to the manufacturer's instructions. First-strand cDNA synthesis was carried out using oligo(dT)_{12–18} and Superscript II (Invitrogen). The cDNA was used as the template in PCR reactions using Ampli Taq Gold polymerase (Applied Biosystems). PCR was performed with specific primers for *mMgl1* (5'-TCTCTGAAAGTGGATGTGGAGG-3', 5'-CAC-TACCCAGCTCAAACAATCC-3'), *mMgl2* (5'-TCTCTGAAAGTG-GATGTGGAGG-3', 5'-GCTATAAGTTGTGGGGAGTGGGC-3'), and *G3pdh* (5'-TGAAGTCGGTGTGAACGGATTGCG-3', 5'-CATGTAG-GCCATGAGGTCCACCAC-3'). The specific primers were designed with Genetyx-Mac (Software Development Co. Ltd., Tokyo, Japan). The conditions were 94 °C for 30 s, 65 °C for 30 s, and 72 °C for 2 min for 40 cycles. Amplifications of *G3pdh* were used as a control for cDNA quantity. The conditions were 94 °C for 30 s, 61 °C for 30 s, and 72 °C for 1 min for 25 cycles. The PCR products were then separated on 1% agarose gels, stained with ethidium bromide, and visualized with the Fluor-S image analyzer (Bio-Rad).

Rapid Amplification of cDNA Ends (RACE)—The poly(A)⁺ RNA from RAW264.7 cells were isolated using a μ MACS mRNA isolation kit (Miltenyi Biotec). Adapter-ligated cDNA were synthesized using the

Marathon cDNA amplification kit (CLONTECH). To obtain a full-length cDNA of mMGL2, an antisense primer, AS3 (5'-TCCTCCACATCCACTTTCAGAG-3'), and a sense primer, 3S1 (5'-TTGGAGCGG-GAAGAGAAAAACCAG-3'), were designed from the expressed sequence tag clones for 5'-RACE and 3'-RACE reactions, respectively. Reactions were incubated at 95 °C for 10 min followed by 43 cycles at 94 °C for 45 s, 60 °C for 45 s, and 72 °C for 2 min. The resulting 1.5-kbp and 700-bp products were subcloned into the pGEM-T easy vector (Promega UK, Southampton, UK) and sequenced with the dye primer method.

Screening of the Genomic Clone Coded the mMgl2 Gene—A 129/SvJ mouse genomic library (Stratagene; vector γ FIXII) was screened with the complete mMGL1 cDNA coding sequence labeled by random priming with ³²P by plaque hybridization. The fragments of positive clones were subcloned into pBluescript SK(+) (Stratagene). These clones were sequenced with 24 specific primers with the dye-terminator method. These sequences were assembled and aligned using Genetyx-Mac.

Production and Isolation of Soluble mMGL2—cDNA encoding mMGL1 and mMGL2 was cloned into pGEM-T easy vector (Promega). The vector was digested with *Bam*HI and *Sac*I (for mMGL1) or *Bam*HI and *Not*I (for mMGL2). The fragments encoding these neck and CRD domains were separated with agarose gel electrophoresis and inserted into each site of expression plasmid vector pET-21a (Novagen). The BL21(DE3) cells containing the plasmid were grown to mid-log phase at 37 °C in 2 \times TY medium (1.2 liters) and then treated with isopropyl β -D-thiogalactoside at a concentration of 1 mM. After isopropyl β -D-thiogalactopyranoside induction, the cultured cells were washed with 50 mM Tris-HCl, pH 8.0, containing 0.15 M NaCl (TBS) and suspended in TBS containing phenylmethanesulfonyl fluoride. The cell lysates were prepared by freezing and thawing, and after the addition of DNase I (final 50 units/ml) and lysozyme (final 0.2 mg/ml), they were incubated 1 h at 37 °C and then centrifuged at 15,000 \times g for 10 min at 4 °C. Expressed proteins formed inclusion bodies. The pellets were washed with TBS containing 0.5% Triton X-100 and 10 mM EDTA and then with H₂O. The washed pellets were solubilized with 2 M NH₄OH (20 ml) and then added to 25 mM MOPS buffer, pH 7.0, containing 2 mM glutathione, reduced form, 0.2 mM glutathione, oxidized form, 20 mM CaCl₂, 0.5 M NaCl, and 0.02% NaN₃ (20 ml). These solutions were then dialyzed against 25 mM MOPS buffer, pH 7.0, containing 20 mM CaCl₂, 0.5 M NaCl, and 0.02% NaN₃. Soluble recombinant mMGL1 and mMGL2 were purified by affinity chromatography on a column of galactose-Sepharose 4B, as described previously (24).

Solid Phase Binding Assays—Absorption of the purified soluble recombinant mMGL1 or mMGL2 onto enzyme-linked immunosorbent assay plates (655061, Greiner) was carried out by adding 100 μ l of solution (3 μ g/ml in DPBS) to each well and incubating the plates for 18 h at 4 °C. After blocking of the wells using 3% bovine serum albumin (BSA) in DPBS for 2 h at room temperature, 150 μ l solution of biotinylated soluble polyacrylamide with attached mono- or oligosaccharides (GlycoTech, Rockville, MD) or hybridoma culture supernatant, diluted into varying concentrations with DPBS containing 3% BSA, were added to each well. After incubation for 2 h at 4 °C (at room temperature for mAb reactions), the wells were washed three times with DPBS to remove unbound materials, and then 100 μ l of HRP-conjugated streptavidin solution (1.25 μ g/ml in DPBS) or HRP-conjugated goat anti-rat IgG (H + L) solution (0.375 μ g/ml in DPBS) for rat mAbs was added to detect bound materials. After incubation for 1 h at room temperature, the wells were washed three times with DPBS. Subsequently, 100 μ l of 1 mM 2, 2'-amino-bis(3-ethylbenzthiazoline-6-sulfonic acid) ammonium (ABTS) solution containing 0.34% H₂O₂ in 0.1 M sodium citrate buffer (pH 4.3) was added, and the absorbance was measured at 405 nm on a microplate reader (25).

To determine pH dependence of the binding of mMGL1 and mMGL2, the incubation conditions were modified as follows. After the blocking buffer was removed, the wells were washed three times with DPBS. Aliquots (50 μ l) of biotinylated soluble polyacrylamide in 2 \times incubation buffer (274 mM NaCl, 26.8 mM KCl, 0.98 mM MgCl₂, 1.810 mM CaCl₂, 2% BSA) were mixed with equal volumes of 2 \times pH buffer (50 mM sodium acetate buffer at pH 4.5–6.0, 50 mM MES buffer at pH 6.0–7.0, 50 mM Tris-HCl buffer at pH 7.0–8.0) and transferred to each well. Incubation was performed at 4 °C for 1.5 h. The wells were emptied and washed three times with DPBST (DPBS containing 0.1% Tween 20) and 100 μ l of HRP-conjugated streptavidin solution (Zymed Laboratories Inc., 1000 \times in DPBS containing 1% BSA) was added and incubated for 1 h at 4 °C. The wells were emptied and washed three times with DPBST, and then 100 μ l of ABTS solution containing 0.034% H₂O₂ (1 mM ABTS dissolved in 0.1 M sodium citrate buffer, pH 4.2) was added, and absorbance at 405 nm was determined.

TABLE I
Splice junction sequences, exon sizes, and estimated intron sizes of the mouse *Mgl2* gene

Exon sequences are in capital letters; intron sequences are in lowercase letters.

Exon no.	Position ^a	Size	Exon/Intron Junction			
			Acceptor sequence	Donor sequence	Intron size	Intron phase
		bp			bp	
1	1–23	23		CACAGgtaac	3684	
2	24–168	145	ttcagTTCTG	TCTAAgtatg	787	1
3	169–279	111	agcagTTCCT	CCAAAgtggg	428	1
4	280–375	96	atcagATTCC	CAGGGgtgag	83	1
5	376–447	72	tcagCTGAC	GGCAGgtgag	79	1
6	448–519	72	ttcagGCCGA	AACAGgtgag	258	1
7	520–606	87	cccagATCTG	CAATGgtgag	103	1
8	607–758	152	cgtagGCTCG	AGCAGgtgat	77	0
9	759–865	107	tttagAATTT	TTCAAgtacg	144	2
10	866–1530	629	ttcagGAATT			

^a Exon positions in coding sequence.

included in the *Mgl* (*Mgl1*) gene (GenBankTM accession number AF132744) (29), but the other clone (termed 41a) was included in the *Mgl2* gene, confirmed by PCR using primers of mMGL2. The latter gene spans 7136 bp, consists of 10 exons, and is similar to *Mgl* (*Mgl1*) in genomic organization (Fig. 2A). The GenBankTM accession number of the *Mgl2* gene is AY103462. The intron/exon boundaries were defined using DNA sequencing. All splice sites conform to the AG/GT rule (Table I). This clone contains 5'-upstream sequences. To identify the promoter sequences of the *Mgl2* gene, transcription factor binding site consensus sequences were searched using the transcription factor data base (www.cbrc.jp/research/db/TFSEARCH.html). The promoter lacks a classical TATA box but contains several binding sites for other transcription factors, C/EBP β , CF1, AML-1, c-Ets, PU.1, c-Rel, Oct-1, Lyf-1, AP-1, GATAs, and STATs. These features are almost identical to that of the 5'-flanking motifs of the *Mgl1* gene. These are the sites found upstream of genes expressed preferentially by cells of monocyte/M ϕ lineages (Fig. 2C).

We identified the BAC clone RPCI-23-172M21, which coded the *Mgl2*, *Asgr1*, and *Asgr2* genes. The other clone RPCI-23-198E14 encodes the *Mgl2* and *Mgl1* genes. These BAC clones have been constructed from the genomic DNA of female C57BL/6J mice. Therefore, these four C-type lectin genes should be located within about 150 kb on mouse chromosome 11 (Fig. 2B).

Reactivity with Monoclonal Antibodies—Previously, we have obtained mAbs specific for MGL1 using purified mMGL (likely to be a mixture of mMGL1 and mMGL2) and recombinant MGL1 as immunogens. Screening to obtain specific monoclonal antibodies was performed previously with recombinant mMGL1 (25). The antibodies are mAb LOM-4.7, mAb LOM-8.2, mAb LOM-8.7, all shown to have blocking activity, mAb LOM-11, shown to recognize the ligand-induced binding site, and a non-blocking mAb LOM-14 (25, 30). As shown in Fig. 3, mAb LOM-4.7, mAb LOM-8.2, mAb LOM-8.7, and mAb LOM-11 were shown to be specific for MGL1. LOM-14 bound to both MGL1 and MGL2 (Fig. 3). These results suggested that the most diverse region of these lectins was the ligand binding site.

Distribution of mMGL2 mRNA in Tissue and Cells—As a means of testing the expression profiles of this putative gene and *mMgl1*, RT-PCR analysis was performed on 10 mouse cell lines including L929 (fibroblast line), JLS-V9 (fibroblast-like bone marrow-derived cell line), EL4 (thymoma cell line), RL δ 1 (lymphoma cell line), YAC-1 (lymphoma cell line), BCL1-B20 (malignant B cells), P815 (mastocytoma cells), P388 (M ϕ -like lymphoid cell line), M1 (myeloblastic leukemia cells), and RAW264.7 (M ϕ -like cell line). The 644-bp bands indicating *mMgl2* were found only in the cell lines




FIG. 3. Binding of mAbs to the recombinant mMGL1 and mMGL2. The binding of mAbs LOM-4.7, LOM-8.2, LOM-8.7, LOM-11, and LOM-14 to the immobilized recombinant mMGL1 (A) and mMGL2 (B) was measured according to mAb concentration. The binding of mAbs was detected using HRP-conjugated goat mAbs specific for rat IgG (H + L). Absorbance at 405 nm was measured on a microplate reader. The values represent means of triplicate determinations, and the error bars indicate S.D.

P388 and RAW264.7 (Fig. 4A). These cell lines were also high expressers of *mMgl1*. To assess the expression patterns of *mMgl1* and *mMgl2* *in vivo*, RT-PCR analysis was conducted




FIG. 4. RT-PCR analysis of mMGL1 and mMGL2 mRNA expression in cells and tissue. *A*, the cDNA was prepared from mouse cell lines (L929, JLS-V9, EL4, RL σ 1, YAC-1, BCL1-B20, P815, P388, RAW264.7, and M1). *B*, tissue cDNA (CLONTECH) was used after normalizing the relative amounts according to the levels of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) cDNA. *C*, TG-PEC were analyzed for the binding of mAbs LOM-14 and LOM-8.7. The cells highly or poorly reactive with mAb LOM-8.7 were separated and tested for mAb LOM-14 binding by flow cytometry. cDNA was prepared from the sorted cells, and RT-PCR analysis was performed. The PCR products were then separated on 1% agarose gels, stained with ethidium bromide, and visualized with the image analyzer. The ratios of band intensity of PCR products were measured. The panels represent a typical result of two separate experiments including cell fractionations. The results of these two separate experiments were almost identical.




FIG. 5. Distribution of cells expressing mMGL1 and mMGL2 in skins. Frozen sections of skins from *Mg11*^{-/-} and *Mg11*^{+/-} mice were stained with mAb LOM-8.7 (specific for mMGL1) and mAb LOM-14 (cross-reactive between mMGL1 and mMGL2).

on RNA isolated from 13 different normal mouse tissues and embryos (Fig. 4B). These genes were expressed almost throughout the body, and the apparent relative intensities among different organs were similar between *mMgl1* and *mMgl2*. To assess whether the expression levels of these genes correspond at single cell levels, TG-PEC were sorted for the binding of mAb LOM-8.7, an mAb specific for mMGL1, and then reacted with mAb LOM-14 reactive with both mMGL1 and mMGL2. As shown in Fig. 4C, cells strongly reactive with mAb LOM-8.7 were also reactive with mAb LOM-14. The ratios of expressed mRNAs corresponding to *mMgl1* and *mMgl2* shown by RT-PCR analysis were almost identical when unsorted mAb LOM-8.7-positive and mAb LOM-8.7-negative cells were compared. These results indicate that the *mMgl1*-positive cells also express the *mMgl2* mRNA.

mMGL2-positive Cells in Tissue Sections from *Mg11*^{-/-} Mice—Frozen sections of skins from *Mg11*^{-/-} mice were stained with mAb LOM-8.7 and mAb LOM-14 (Fig. 5). The results revealed that *Mg11*^{-/-} mice expressed epitopes reactive with mAb LOM-14 at lower levels than *Mg11*^{+/-} mice but not with mAb LOM-8.7. These epitopes are likely to represent mMGL2. The subcellular localization of the mAb LOM-14 staining in *Mg11*^{-/-} mice did not appear to be distinct from that of mAb LOM-8.7 staining in *Mg11*^{+/-} mice.

Carbohydrate Specificity of mMGL2—A variety of biotin-labeled soluble polyacrylamides with mono- or oligosaccharides were applied to determine the carbohydrate specificity of mMGL1 and mMGL2 (Fig. 6). The carbohydrates tested are listed in the figure. The recombinant form of mMGL1 had the highest affinity with Le^x residues among all carbohydrate-modified polyacrylamides tested. mMGL2 showed a very low affinity with Le^x but showed the highest affinity with β -linked GalNAc residues (Fig. 6). The bindings were inhibited with 5 mM EDTA (data not shown).

pH-dependent Binding of Ligands to mMGL—Enzyme-linked immunosorbent assays were performed under various pH conditions to determine the binding of soluble polyacrylamides with Le^x residues or α -GalNAc residues to immobilized recombinant mMGL1 or mMGL2, respectively. Under a Ca²⁺ concentration similar to that found in extracellular fluids (0.905 mM), the binding decreased when pH was lowered from extracellular (pH 7.3) to endosomal (pH 5.4). The profiles of the pH-dependent binding were similar to that observed with intact hepatic asialoglycoprotein receptors


(31). The pH value of a half-maximal ligand binding, designated as pH_B, was 5.5 for MGL1 and 5.7 for MGL2 (Fig. 7).

DISCUSSION

In the present study, we found a novel MØ C-type lectin, mMGL2, and characterized its genomic structure, expression patterns, and carbohydrate specificity. The *Mgl2* gene spans 7136 bp and consists of 10 exons, which is a similar genomic organization to that of the *Mgl* gene now renamed the *Mgl1* gene. We reported that the *Mgl1* gene linked to *Trp53* and is located 1.8 ± 1.2 cM distal to *D11Mit5* and 1.8 ± 1.2 cM proximal to *Htt* on mouse chromosome 11. Using a panel of DNA samples from two parental mice, C3H/HeJ-*gld* and (C3H/HeJ-*gld* × *Mus spretus*) F1 were digested with various restriction endonucleases and hybridized with a mMGL1 cDNA probe to determine the restriction fragment length polymorphism and to allow haplotype analyses (29). The hepatic asialoglycoprotein receptor genes (*Asgr1* and *Asgr2*) were also known to be linked to *Trp53* (32). We found BAC clones, which included these two C-type lectins. These results indicate that the highly homologous *Mgl2* gene is linked within about 50 kb. Their homology and close genomic localization indicate that the original gene was duplicated recently. The human *MGL* (*HML-2*) gene was mapped in chromosome 17p and linked to *ASGR1* and *ASGR2*. These results show that there is a cluster of type 2 C-type lectin genes on mouse chromosome 11 and on human 17p12–13. Likewise, the natural killer gene complex resided on mouse chromosome 6 and on human 12p (33, 34).

Although there was a high degree of sequence homology between mMGL1 and mMGL2, the sequences corresponding to cytoplasmic and CRD showed differences, suggesting that cytoplasmic tail associations and the carbohydrate recognitions are unique between mMGL1 and mMGL2. We showed that recombinant mMGL1 bound soluble polyacrylamides containing Le^x oligosaccharides. mMGL2 had affinity with polyacrylamides containing α - or β -GalNAc. It has been believed that C-type lectins contain a peptide segment that corresponds to the carbohydrate binding specificity within the CRD and that the sequence in Gal and GalNAc-binding lectins, such as ASGRs, is QPD. In contrast, the sequence in mannose-, fucose-, and GlcNAc specific lectins, such as serum mannose-binding proteins, was EPN. The sequence in both mMGL1 and -2 was QPD. However, it was obvious from our results that their fine specificities depended on other amino acids in the CRD (Figs. 6 and 8). Some of the amino acid residues important in the

FIG. 6. Carbohydrate specificity of recombinant mMGL1 and mMGL2. Binding of biotin-labeled soluble polyacrylamides containing mono- or oligosaccharides to immobilized recombinant mMGL1 (A and C) and mMGL2 (B and D). Amounts of bound polymers incubated at various concentrations were quantified using HRP-conjugated streptavidin. ABTS was used as a substrate, and absorbance at 405 nm was measured on a microplate reader. The panels represent a typical result of three separate determinations, which showed almost identical results. Each assay was performed in duplicate, and the mean value is shown.



carbohydrate recognitions were conserved in mMGL1, mMGL2, hMGL, RHL-1 and -2, mASGR-1, and hASGR-1. Crystallographic determination of the structure of human ASGR-1 revealed amino acid residues important in sugar binding (35). Furthermore, rat hepatic asialoglycoprotein receptor was subjected to site-directed mutation. As a result, the pH dependence of ligand binding was shown to be mediated by His-256, Asp-266, and Arg-270 (36). Because these residues are also conserved in mMGL1 and -2, they should also be responsible for their pH dependence (Fig. 8). Amino acid residues responsible for the differential carbohydrate specificity between mMGL1 and -2 and other members of the C-type lectin family and the phylogeny of these lectins (Fig. 8) are yet to be elucidated.

There is a possibility that mMGL1 and -2 form heterooligomers. These lectins are known to form trimeric structures through the interactions of the neck domain in a similar manner to that of ASGRs. ASGRs are abundantly expressed on the sinusoidal surface of hepatic parenchymal cells (37, 38). Its primary role is the removal and degradation of desialylated glycoproteins from circulation. High affinity binding requires the receptor to be assembled as a heterooligomer consisting of two highly homologous subunits, termed hepatic lectin 1 and 2 (39). Experiments with recombinant rat hepatic lectins suggested that the binding properties of the major subunit (RHL-1) and the minor subunit (RHL-2/-3) were optimized for different ligands (40). In the case of mMGL1 and -2, the recombinant form corresponding to each one showed affinity with different carbohydrates, indicating that heterooligomer formation was not required for their carbohydrate recognition. However, these lectins were likely to be expressed on the same cells at the single cell level and cooperatively functioned to recognize and uptake extracellular molecules.

mMGL1-positive cells are abundant in connective tissues throughout the body (19). mMGL1 was shown to be expressed on the surfaces of bone marrow-derived immature dendritic cells (21). hMGL is also shown to be expressed on monocyte-derived immature dendritic cells and monocyte-derived immature MØs in humans (41). We have shown that mMGL1-positive cells migrate from dermis to regional lymph nodes during the sensitization phase of contact hypersensitivity (42). The




FIG. 7. Effects of pH on binding of recombinant mMGL1 and mMGL2. The binding of biotin-labeled soluble polyacrylamides (0.22 µg/ml Le^x polyacrylamide or 0.67 µg/ml α-GalNAc polyacrylamide) to the immobilized recombinant mMGL1 (A) or mMGL2 (B) was determined under varied pH conditions. The assays were performed in the presence of 0.905 mM CaCl₂. The values obtained in different buffers (25 mM sodium acetate buffer for pH 4.5–6.0, 25 mM MES buffer for pH 6.0–7.0, 25 mM Tris-HCl buffer for pH 7.0–8.0). The panels represent a typical result of two separate determinations, which showed almost identical results. The values represent means of duplicate determinations.

A

mMGL2	1:S-EVACCPHWHTEHEGSCYWFSESEKSWPEADKYCRLENSHLVVVNSLEEQNF--LQNR	57
mMGL1	1:S-EVACCPHWHTEHEGSCYWFSESEKSWPEADKYCRLENSHLVVVNSLEEQNF--LQNR	57
rMGL	1:S-AVACCPHWHMEHEGSCYWFSSQSGKPWPEADKYCQLENSNLVVVNSLAEQNF--LQTH	57
hMGL	1:--EGTCCPVNWEHQDSCYWFSSGMSWAEAEKYCQLENSHLVVVNSLAEQNF--VQKY	56
hASGR1	1:S-ERTCCPVNWEHERSICYWFSSRSGKAWADADNYCRLEDAHLVVVTSWEEQKF--VQHH	57
RHL-1	1:S-ERICCPINWVEYEGSCYWFSSSVKPWTEADKYCQLENSHLVVVTSWEEQRF--VQHM	57
MHL1	1:S-ERTCCPVNWEYEGSCYWFSSSVRPWTEADKYCQLENSHLVVVTSRDEQNF--LQRH	57
hASGR2	1:--SQRTCCPVNWEHQGSCYWFSSGKAWAEAEKYCQLENSHLVVVNSWEEQKF--IVQHT	57
RHL-2/3	1:--SNGTECCPVNWEHFQGGSCYWFSSRDLTWAEDQYQCMENAHLLVINSREEQEF--VVKH	58
MHL2	1:--TECCPVNWEHFQGGSCYWFSSRDLTWAEDQYQCMENAHLLVINSREEQDF--VVKH	55
rKCR	1:-----AENFCVSQGAHLASVTSQEEQAF--LVQIT	28
mKCR	1:-----AEKFCVSQGAHLASVTSQEEQAF--LVQIT	28
hCD23	1:-----TCPEKWINFQKCYFYFGKTKQWVHARYACDDMEGQLVSIHSPREEQDF--LTKH	53
mCD23	1:-----ICPKNWLHFQKCYFYFGKSKQWIQARFACSDLQGLVSIHSPREEQDF--LQHI	53
hDC-SIGN	1:-----PCFEWTFQGNICYFMSNSQRNWHDSITACKEVGAQLVVIKSAAEQNFLLQSSR	55
hL-SIGN	1:-----HCPKDWTFQGNICYFMSNSQRNWHDSVTACQEVRAQLVVIKTAEEQNFLQLQTSR	55
	* * *	
mMGL2	58:ANVLSWMGL--TDQNGPWRWVDG--TDFDKGFKNWRP	113
mMGL1	58:ANVVSWIGL--TDQNGPWRWVDG--TDFEKGFKNWAP	113
rMGL	58:GSVVTWIGL--TDQNGPWRWVDG--TDYEKGFTHWAPK	113
hMGL	57:GSAYTWIGL--SDPEGAWKWDG--TDYATGFQNWKPG	112
hASGR1	58:GPVNTWIGL--HDQNGPWKWDG--TDYETGFKNWRPE	113
RHL-1	58:GPLNTWIGL--TDQNGPWKWDG--TDYETGFKNWRPE	113
MHL1	58:GPLNTWIGL--TDQNGPWKWDG--TDYETGFQNRPE	113
hASGR2	58:NPFNTWIGL--TDSGWSWKWDG--TDYRHNKKNWAVT	113
RHL-2/3	59:GAFHIWIGL--TDKGSWKWDG--TEYRSNFKNWAVT	114
MHL2	56:SQFHIWIGL--TDRDGSWKWDG--TDYRSNYRNWAVT	111
rKCR	29:NAVDHWIGLTDQGTGEGNRWRWVDGTFPDYVQSRRFWRK	85
mKCR	29:SSGDHWIGLTDQGTGEGIRWRWVDGTFPNNAQSKGFWGKN	85
hCD23	54:SHTGSWIGLRNLDLKGFEIWDG--S--HVDYSNWAPGPTS-RSQ--G-EDCVMRGS	104
mCD23	54:NKKDSWIGLQDLNMEGEFVWSDG--S--PVGYSNWNPC	104
hDC-SIGN	56:SNRFTWIGLSDLNQEGTWQWVDG--SPLLPSFKQYWRGEPN	108
hL-SIGN	56:SNRFSWMGLSDLNQEGTWQWVDG--SPLSPSFQRYWNSGEPN	108
	* * * * *	
mMGL2	114:GRWNDDVC-QRHYHWICETELGKASSAHSQPQIASVP-----	149
mMGL1	114:GPWNDDVC-QRTFRWICEMKLAKES-----	137
rMGL	114:GRWNDDVC-QRPYRWVCEMKLAKDS-----	137
hMGL	113:GRWNDDVC-QRPYHWVCEAGLGQTSQESH-----	140
hASGR1	114:GRWNDDVC-QRPYRWVCETELDKASQEPPLL-----	143
RHL-1	114:GHWNDVC-RRPYRWVCETELKGAN-----	137
MHL1	114:GRWNDDVC-RRPYRWVCETELKDKAN-----	137
hASGR2	114:GRWNDDVC-LQVYRWVCE-KRRNATGEVA-----	140
RHL-2/3	115:GLWNDNFC-QQVNRWACE-RKRDRITY-----	138
MHL2	112:GHWNDNFC-QQVNRWACE-KRRNITH-----	135
rKCR	86:RMWNDMAC-GTAYNWVCKKSTDWSVARTDQS-----	115
mKCR	86:QQWNDMAC-GSSYPWVCKKSTGWSAARVG-----	113
hCD23	105:GRWDAFCRDKLGAWVCDRLATCTPPASEGSAESMGFDSRDPDGRLETPSAPLHS	160
mCD23	105:GQWDAFCRSYLDWVCEQLATCEISAPLASVTPTRETPKSEP-----	147
hDC-SIGN	109:G-WNDKCN-LAKFWIC-KKSAASCRDEEQFLSPAPATPNPPPA-----	150
hL-SIGN	109:G-WNDNRC-DVDNYWIC-KKPAA-CFRDE-----	133
	*** * * *	


B

Fig. 8. Comparison of the amino acid sequences of the CRD of C-type lectins. A, multialignment of the CRD sequences of C-type lectins. B, evolutionary tree of sequence of CRD of C-type lectins. Abbreviations used in this figure are: *MHL*, mouse hepatic lectin; *rKCR*, rat Kupffer cell receptor; *mKCR*, mouse Kupffer cell receptor.

migration was initiated by cytokine-mediated release of mMGL1-positive cells from dermis (42–44). The cells homed to the boundary of the T-cell area in the regional lymph nodes, and the prevention of migration seemed to interfere with sensitization. Involvement of coordinated functions of mMGL1 and -2 in such pathogenic processes should be the most important subject of future investigations.

REFERENCES

- Lewis, C., and McGee, J. O. D. (1992) *The Macrophage: The Natural Immune System*, pp. 1–413, IRL Press at Oxford University Press, Oxford
- Feizi, T. (2000) *Immunol. Rev.* **173**, 79–88
- McKnight, A. J., and Gordon, S. (1998) *Adv. Immunol.* **68**, 271–314
- Linehan, S. A., Martinez-Pomares, L., and Gordon, S. (2000) *Microbes Infect.* **2**, 279–288
- Weis, W. I., Taylor, M. E., and Drickamer, K. (1998) *Immunol. Rev.* **163**, 19–34
- Stahl, P. D., and Ezekowitz, R. A. (1998) *Curr. Opin. Immunol.* **10**, 50–55
- Liu, Y., Chirino, A. J., Misulovin, Z., Leteux, C., Feizi, T., Nussenzweig, M. C., and Bjorkman, P. J. (2000) *J. Exp. Med.* **191**, 1105–1116
- Leteux, C., Chai, W., Loveless, R. W., Yuen, C. T., Uhlin-Hansen, L., Combarous, Y., Jankovic, M., Maric, S. C., Misulovin, Z., Nussenzweig, M. C., and Ten, F. (2000) *J. Exp. Med.* **191**, 1117–1126
- Ozaki, K., Ii, M., Itoh, N., and Kawasaki, T. (1992) *J. Biol. Chem.* **267**, 9229–9235
- Kawakami, K., Yamamoto, K., Toyoshima, S., Osawa, T., and Irimura, T. (1994) *Jpn. J. Cancer Res.* **85**, 744–749
- Valladeau, J., Duvert-Frances, V., Pin, J. J., Kleijmeer, M. J., Ait-Yahia, S., Ravel, O., Vincent, C., Vega, F., Jr., Helms, A., Gorman, D., Zurawski, S. M., Zurawski, G., Ford, J., and Saeland, S. (2001) *J. Immunol.* **167**, 5767–5774
- Yamamoto, K., Ishida, C., Shinohara, Y., Hasegawa, Y., Konami, Y., Osawa, T., and Irimura, T. (1994) *Biochemistry* **33**, 8159–8166
- Suzuki, N., Yamamoto, K., Toyoshima, S., Osawa, T., and Irimura, T. (1996) *J. Immunol.* **156**, 128–135
- Iida, S., Yamamoto, K., and Irimura, T. (1999) *J. Biol. Chem.* **274**, 10697–10705
- Sakamaki, T., Imai, Y., and Irimura, T. (1995) *J. Leukocyte Biol.* **57**, 407–414
- Ichii, S., Imai, Y., and Irimura, T. (1997) *J. Leukocyte Biol.* **62**, 761–770
- Ichii, S., Imai, Y., and Irimura, T. (2000) *Cancer Immunol. Immunother.* **49**, 1–9
- Imai, Y., Akimoto, Y., Mizuochi, S., Kimura, T., Hirano, H., and Irimura, T. (1995) *Immunology* **86**, 591–598
- Mizuochi, S., Akimoto, Y., Imai, Y., Hirano, H., and Irimura, T. (1997) *Glycobiology* **7**, 137–146
- Mizuochi, S., Akimoto, Y., Imai, Y., Hirano, H., and Irimura, T. (1998) *Glycoconj. J.* **15**, 397–404
- Denda-Nagai, K., Kubota, N., Tsuiji, M., Kamata, M., and Irimura, T. (2002) *Glycobiology*, **12**, 443–450
- Higashi, N., Fujioka, K., Denda-Nagai, K., Hashimoto, S., Nagai, S., Sato, T., Fujita, Y., Morikawa, A., Tsuiji, M., Miyata-Takeuchi, M., Sano, Y., Suzuki, N., Yamamoto, K., Matsushima, K., and Irimura, T. (2002) *J. Biol. Chem.* **277**, 20686–20693
- Onami, T. M., Lin M.-Y., Page, D. M., Reynolds, S. A., Katayama, C. D., Marth, J. D., Irimura, T., Varki, A., Varki, N., and Hedrick, S. M. (2002) *Mol. Cell. Biol.* **22**, 5173–5181
- Sato, M., Kawakami, K., Osawa, T., and Toyoshima, S. (1992) *J. Biochem. (Tokyo)* **111**, 331–336
- Kimura, T., Imai, Y., and Irimura, T. (1995) *J. Biol. Chem.* **270**, 16056–16062
- Tanaka, I., and Ishihara, H. (2001) *Virology* **280**, 107–114
- Mager, D. L., and Freeman, J. D. (2000) *J. Virol.* **74**, 7221–7229
- Trowbridge, I. S. (1991) *Curr. Opin. Cell Biol.* **3**, 634–641
- Tsuiji, M., Fujimori, M., Seldin, M. F., Taketo, M. M., and Irimura, T. (1999) *Immunogenetics* **50**, 67–70
- Kimura, T., Hosoi, T., Yamamoto, K., Suzuki, N., Imai, Y., and Irimura, T. (2000) *Mol. Immunol.* **37**, 151–160
- Hudgin, R. L., Pricer, W. E., Jr., Ashwell, G., Stockert, R. J., and Morell, A. G. (1974) *J. Biol. Chem.* **249**, 5536–5543
- Blackburn, C. C., Griffith, J., and Morahan, G. (1995) *Genomics* **26**, 308–317
- Yokoyama, W. M., and Seaman, W. E. (1993) *Annu. Rev. Immunol.* **11**, 613–635
- Brown, M. G., Fulmek, S., Matsumoto, K., Cho, R., Lyons, P. A., Levy, E. R., Scalzo, A. A., and Yokoyama, W. M. (1997) *Genomics* **42**, 16–25
- Meier, M., Bider, M. D., Malashkevich, V. N., Spiess, M., and Burkhard, P. (2000) *J. Mol. Biol.* **300**, 857–865
- Wragg, S., and Drickamer, K. (1999) *J. Biol. Chem.* **274**, 35400–35406
- Collins, J. C., Stockert, R. J., and Morell, A. G. (1984) *Hepatology* **4**, 80–83
- Braun, J. R., Willnow, T. E., Ishibashi, S., Ashwell, G., and Herz, J. (1996) *J. Biol. Chem.* **271**, 21160–21166
- Bider, M. D., and Spiess, M. (1998) *FEBS Lett.* **434**, 37–41
- Ruiz, N. I., and Drickamer, K. (1996) *Glycobiology* **6**, 551–559
- Higashi, N., Morikawa, A., Fujioka, K., Fujita, Y., Sano, Y., Miyata-Takeuchi, M., Suzuki, N., Irimura, T. (2002) *Int. Immunol.* **14**, 545–554
- Sato, K., Imai, Y., and Irimura, T. (1998) *J. Immunol.*, **161**, 6835–6844
- Chun, K. H., Imai, Y., Higashi, N., Irimura, T. (2000) *J. Leukocyte Biol.* **68**, 471–478
- Chun, K. H., Imai, Y., Higashi, N., Irimura, T. (2000) *Int. Immunol.* **12**, 1695–1703