



University of Tennessee, Knoxville  
**Trace: Tennessee Research and Creative Exchange**

---

Insects, Pests, Plant Diseases and Weeds

UT Extension Publications

---

5-14-2013

# W 289-J IPM QuickFacts Series: Potato Leafhopper

Amy Fulcher

Heather Bowers

Follow this and additional works at: [http://trace.tennessee.edu/utk\\_agexdise](http://trace.tennessee.edu/utk_agexdise)

 Part of the [Agricultural Education Commons](#), and the [Plant Sciences Commons](#)

---

## Recommended Citation

"W 289-J IPM QuickFacts Series: Potato Leafhopper," Amy Fulcher and Heather Bowers, W289-J, [http://trace.tennessee.edu/utk\\_agexdise/73](http://trace.tennessee.edu/utk_agexdise/73)

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Ornamental insects, Pests & Diseases is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Insects, Pests, Plant Diseases and Weeds by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

# Potato Leafhopper

## *Empoasca fabae*

### Description

Potato leafhoppers can be identified visually by their wedge shape and pale green color. Adult potato leafhoppers are about 1/8 inch long and hold their wings “roof-like” over the abdomen. Potato leafhopper nymphs look like miniature adults and have visible wing “buds” instead of fully developed wings.

### Life Cycle

An adult female potato leafhopper mates and deposits 60 to 100 eggs into leaf veins and leaf petioles of host plants. Nymphs hatch in six to nine days. Development from egg to adult takes between 20 and 30 days, with the duration increasing as temperatures decrease. Adults typically survive about 30 days. Several overlapping generations may occur during a typical growing season. In Tennessee, potato leafhopper cannot overwinter due to the cold winter temperatures. They overwinter south of Tennessee and are blown north by prevailing winds, typically arriving in the state between April and early May.

### Monitoring

Potato leafhoppers can be hard to spot on plants; however, when disturbed, adults will either jump or make short-distance flights. Adults and nymphs will crab-walk sideways and hide on the opposite side of leaves when disturbed. Look for potato leafhoppers feeding on the underside of leaves and look for stippling on leaf surfaces and curled or stunted shoot tips. Another good diagnostic cue may be the cast skins left by nymphs that have molted and may be attached to lower leaf surfaces. Potato leafhopper is often found on alfalfa until it is cut and then migrates to ornamentals. Activity can be monitored by using yellow sticky traps. Populations often decline by mid-July.

### Host Plants

Red maples are the most frequently affected ornamental plant in nurseries.

Hosts for other leafhopper species include the following:

- Apple
- Ash
- Birch
- Crabapple
- Dogwood
- Elm
- Hickory
- Oak
- Redbud

# Damage Symptoms



Potato leafhoppers use their piercing-sucking mouthparts to feed on plant vascular tissues. Feeding removes the chlorophyll, creating small, light, angular stippled spots on leaves. During feeding, potato leafhoppers inject a salivary toxin that disrupts sap flow and causes leaf tips and edges to turn brown, curl and die. This characteristic symptom is known as “hopperburn.” When the vascular tissue of shoots is damaged, the internode length of the new growth is greatly reduced.

Hopperburn is primarily aesthetic and seldom results in tree death, although the damage can reduce plant cold hardiness. Mature landscape trees typically experience little potato leafhopper damage and can generally tolerate feeding injury. In production, extra labor will be needed to prune out multiple terminal leaders. Poor aesthetic appearance of damaged trees may also reduce the market value of affected trees.

## Integrated Pest Management

### BIOLOGICAL CONTROL

Ladybird beetles, green lacewings, parasitic wasps, damsel bugs, bigeyed bugs and assassin bugs are all effective predators of potato leafhopper adults and nymphs.

### CULTURAL CONTROL

Resistant cultivars include *Acer rubrum* ‘Brandywine’, *Acer rubrum* ‘Somerset’ and *Acer rubrum* ‘Sun Valley’.

### CHEMICAL CONTROL

Soil drenches or foliar spray applications are common control options. However, pesticides applied to control leafhoppers may trigger outbreaks of maple spider mite (see Frank and Sadof below).

Please refer to [http://eppserver.ag.utk.edu/redbook/sections/trees\\_flowers.htm](http://eppserver.ag.utk.edu/redbook/sections/trees_flowers.htm) for current recommendations.

## Resources

- Photo credits: Amy Fulcher, University of Tennessee; Ric Bessin, University of Kentucky; Steve L. Brown, University of Georgia; Bugwood.org
- Cloyd, R. 2001. Potato leafhopper. University of Illinois Extension. Home, Yard & Garden Pest Newsletter (4). <http://hyg.ipm.illinois.edu/pastpest/200104c.html>
- Frank, S. and C. Sadof. 2011. Reducing insecticide volume and nontarget effects of ambrosia beetle management in nurseries. J. Econ. Ent. 104(6):1960-1968.
- Gill, S. 2010. Potato leafhopper (*Empoasca fabae*). University of Maryland Extension fact sheet. <http://ipmnet.umd.edu/factsheets/PotatoLeafhopper-UMD.pdf>
- Oliver, J.B., D.C. Fare, N. Youssef, M.A. Halcomb, M.E. Reding, and C.M. Ranger. 2009. Evaluation of systemic insecticides for potato leafhopper control in field-grown red maple. Journal of Environmental Horticulture 27(1):17-23. [http://www.ars.usda.gov/research/publications/publications.htm?seq\\_no\\_115=225970](http://www.ars.usda.gov/research/publications/publications.htm?seq_no_115=225970)
- Potter, D. and P. Spicer. 1993. Seasonal phenology, management, and host preferences of potato leafhopper on nursery-grown maples. Journal of Environmental Horticulture 11(3):101-106. [http://www.hriresearch.org/docs/publications/JEH/JEH\\_1993/JEH\\_1993\\_11\\_3/JEH%2011-3-101-106.pdf](http://www.hriresearch.org/docs/publications/JEH/JEH_1993/JEH_1993_11_3/JEH%2011-3-101-106.pdf)
- Riemann, A.B., W.E. Klingeman, and F.A. Hale. 2006. Spurn “Hopper Burn” by understanding and managing leafhoppers. Tennessee Greentimes 7(1):24-27.

THE UNIVERSITY of TENNESSEE   
INSTITUTE of AGRICULTURE

Prepared by Heather Bowers, Intern, Department of Plant Sciences  
and Dr. Amy Fulcher, Assistant Professor, Department of Plant Sciences  
Publication funded by USDA Extension IPM Grant in partnership with University of Kentucky Integrated Pest Management Program.  
The authors thank M. Halcomb, B. Klingeman, W. Russell and F. Hale for their careful review.

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.  
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.  
UT Extension provides equal opportunities in programs and employment.