



5-14-2013

## W289-L IPM QuickFacts Series: Black Root Rot

Amy Fulcher

Heather Bowers

Follow this and additional works at: [https://trace.tennessee.edu/utk\\_agexdise](https://trace.tennessee.edu/utk_agexdise)



Part of the [Agricultural Education Commons](#), and the [Plant Sciences Commons](#)

---

### Recommended Citation


"W289-L IPM QuickFacts Series: Black Root Rot," Amy Fulcher and Heather Bowers,  
W 289-L  
, [https://trace.tennessee.edu/utk\\_agexdise/74](https://trace.tennessee.edu/utk_agexdise/74)

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Ornamental insects, Pests & Diseases is brought to you for free and open access by the UT Extension Publications at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Insects, Pests, Plant Diseases and Weeds by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

# Black Root Rot

## *Thielaviopsis basicola*




### Description

Black root rot is caused by the fungus *Thielaviopsis basicola*. Although it attacks mainly herbaceous plants, it is a very serious disease of Japanese, inkberry and the blue Meserve hybrid hollies.

### Host Plants

- Begonia
- Cyclamen
- Geranium
- Gerbera
- Gloxinia
- Holly (Japanese, Blue, Inkberry)
- Pansy
- Petunia
- Phlox
- Poinsettia
- Sweet pea
- Verbena
- Violet



### Disease Cycle

The black root rot pathogen prefers soil temperatures between 55 and 65 degrees F. The disease can spread in irrigation water and substrate by insects such as shore flies and fungus gnats and also can spread long distance via infected plants. The fungus can survive in the soil or it can survive as a saprophyte on plant debris. It overwinters as chlamydospores.

### Monitoring

Monitor environmental conditions that stress the host plant, such as high temperatures for pansy. When trying to identify black root rot, look for black lesions on the roots. The contrast in color makes it easy to spot on otherwise healthy white roots. Check roots of otherwise healthy-appearing hollies, as ideal nursery conditions can mask infection.



# Symptoms



Some early symptoms of black root rot include yellowing of foliage, marginal leaf scorch and overall plant decline. Black root rot may be mistaken for a nutrient deficiency. Infected herbaceous plants wilt rapidly during the hottest part of the day and recover during the evening hours. Branches may die back, and the entire plant may eventually die. Root systems are reduced to roots that are short, stubby, black and decayed.



## Integrated Pest Management

### BIOLOGICAL CONTROL

*Pseudomonas fluorescens* and mycorrhizal fungi have been used to suppress *Thielaviopsis*.

### CULTURAL CONTROL

Maintain the soil pH below 6.5. Avoid excessive irrigation, which could attract fungus gnats and shore flies. Plant resistant species. Examine root systems prior to planting container plants. Avoid planting susceptible species in soils previously infested with the fungus. Sanitize recycled containers before using. Avoid using contaminated substrate.

### CHEMICAL CONTROL

Please refer to [http://eppserver.ag.utk.edu/redbook/sections/trees\\_flowers.htm](http://eppserver.ag.utk.edu/redbook/sections/trees_flowers.htm) for the most up-to-date recommendations.

# Resources

Photo credits: Alan Windham, University of Tennessee

Adkins, C., G. Armel, M. Chappell, J.C. Chong, S. Frank, A. Fulcher, F. Hale, K. Ivors, W. Klingeman III, A. LeBude, J. Neal, A. Senesac, S. White, and A. Windham. 2010. Pest management strategic plan for container and field-produced nursery crops in GA, KY, NC, SC, TN. A. Fulcher, ed. Southern Region IPM Center.

Bost, S. and D. Hensely. 2011. Black root rot of tobacco. University of Tennessee Extension publication W274.

<https://utextension.tennessee.edu/publications/Documents/W274.pdf>

Moorman, G.W. Black root rot (Thielaviopsis). Penn State University.

<http://extension.psu.edu/plant-disease-factsheets/all-fact-sheets/thielaviopsis-or-black-root-rot>

THE UNIVERSITY of TENNESSEE   
INSTITUTE of AGRICULTURE

Prepared by Heather Bowers, Intern, Department of Plant Sciences  
and Dr. Amy Fulcher, Assistant Professor, Department of Plant Sciences  
Publication funded by USDA Extension IPM Grant in partnership with University of Kentucky Integrated Pest Management Program.  
The authors thank M. Halcomb, W. Russell and A. Windham for their careful review.

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.  
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.  
UT Extension provides equal opportunities in programs and employment.