



7-23-2010

## Comparative functional genomic study of substrate specificity evolution of the SABATH family of methyltransferases in plants

Nan Zhao  
*University of Tennessee - Knoxville*

Jean-Luc Ferrer  
*Universite Joseph Fourier (Grenoble I)*

Xiaofeng Zhuang  
*University of Tennessee - Knoxville*

Feng Chen  
*University of Tennessee - Knoxville, fengc@utk.edu*

Follow this and additional works at: [https://trace.tennessee.edu/utk\\_planpubs](https://trace.tennessee.edu/utk_planpubs)



Part of the [Bioinformatics Commons](#), and the [Plant Sciences Commons](#)

---

### Recommended Citation

BMC Bioinformatics 2010, 11(Suppl 4):P7 doi:10.1186/1471-2105-11-S4-P7

This Article is brought to you for free and open access by the Plant Sciences at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Plant Sciences Publications and Other Works by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

POSTER PRESENTATION

Open Access

# Comparative functional genomic study of substrate specificity evolution of the SABATH family of methyltransferases in plants

Nan Zhao<sup>1</sup>, Jean-Luc Ferrer<sup>2</sup>, Xiaofeng Zhuang<sup>1</sup>, Feng Chen<sup>1\*</sup>

From UT-ORNL-KBRIN Bioinformatics Summit 2010  
Cadiz, KY, USA. 19-21 March 2010

## Background

The plant SABATH protein family is composed of a group of related small molecule methyltransferases (MTs) that catalyze the S-adenosyl-L-methionine dependent methylation of a variety of plant small molecular weight metabolites encompassing widely divergent structures. Some of these substrates are important plant hormones and signaling molecules, such as indole-3-acetic acid (IAA), jasmonic acid (JA) and salicylic acid (SA). Methylating these compounds may have important impacts on plant growth and development. In the previous paper, we presented Indole-3-acetic acid (IAA) methyltransferase (IAMT) as an evolutionarily ancient member of the SABATH family in higher plants. Whether the IAMT exists in less evolutionarily advanced plants is still unknown.

## Materials and methods

To further understand the evolution of the SABATH family in land plants, we undertook an integrated functional genomic approach to identify and characterize SABATH genes in a lower plant moss (*Physcomitrella patens*). Four putative moss SABATH genes were identified using bioinformatics tools.

## Results and conclusion

Enzymatic assay displayed none of them had IAMT activity, suggesting that IAMT might be evolved after the divergence of lower and higher plants. However, one of them, *PpSABATH1*, showed methyltransferase activity with a number of compounds containing sulhydryl or selenohydryl groups. Tobacco plants overexpressing the

*PpSABATH1* gene under the control of CaMV35S promoter exhibited an enhanced tolerance to thiobenzoic acid. Together with these results, we hypothesize that IAMT, the evolutionarily ancient member of the SABATH family in higher plants, evolved from a sulfur methyltransferase.

## Author details

<sup>1</sup>Department of Plant Sciences, University of Tennessee, Knoxville, TN 37996, USA. <sup>2</sup>Institut de Biologie Structurale, Centre National de la Recherche Scientifique, Université Joseph Fourier, 38027 Grenoble cedex 1, France.

Published: 23 July 2010

doi:10.1186/1471-2105-11-S4-P7

Cite this article as: Zhao et al.: Comparative functional genomic study of substrate specificity evolution of the SABATH family of methyltransferases in plants. *BMC Bioinformatics* 2010 **11**(Suppl 4):P7.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit



\* Correspondence: fengc@utk.edu

<sup>1</sup>Department of Plant Sciences, University of Tennessee, Knoxville, TN 37996, USA