



11-12-2013

Acyl-homoserine lactone-based quorum sensing in the Roseobacter clade: complex cell-to-cell communication controls multiple physiologies

W Nathan Cude

Alison Buchan

University of Tennessee - Knoxville

Follow this and additional works at: https://trace.tennessee.edu/utk_micrpubs

 Part of the [Microbiology Commons](#)

Recommended Citation

Cude WN and Buchan A (2013) Acyl-homoserine lactone-based quorum sensing in the Roseobacter clade: complex cell-to-cell communication controls multiple physiologies. *Front. Microbiol.* 4:336. doi: 10.3389/fmicb.2013.00336

This Article is brought to you for free and open access by the Microbiology at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Microbiology Publications and Other Works by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.



Acyl-homoserine lactone-based quorum sensing in the *Roseobacter* clade: complex cell-to-cell communication controls multiple physiologies

W. Nathan Cude and Alison Buchan *

Department of Microbiology, University of Tennessee, Knoxville, TN, USA

Edited by:

Benjamin Van Mooy, Woods Hole Oceanographic Institution, USA

Reviewed by:

Shawn R. Campagna, University of Tennessee, Knoxville, USA

Laura Hmelo, University of Washington, USA

Shady A. Amin, University of Washington, USA

*Correspondence:

Alison Buchan, Department of Microbiology, University of Tennessee, M409 Walters Life Sciences, Knoxville, TN 37996-0845, USA

e-mail: abuchan@utk.edu

Bacteria have been widely reported to use quorum sensing (QS) systems, which employ small diffusible metabolites to coordinate gene expression in a population density dependent manner. In Proteobacteria, the most commonly described QS signaling molecules are *N*-acyl-homoserine lactones (AHLs). Recent studies suggest that members of the abundant marine *Roseobacter* lineage possess AHL-based QS systems and are environmentally relevant models for relating QS to ecological success. As reviewed here, these studies suggest that the roles of QS in roseobacters are varied and complex. An analysis of the 43 publically available *Roseobacter* genomes shows conservation of QS protein sequences and overall gene topologies, providing support for the hypothesis that QS is a conserved and widespread trait in the clade.

Keywords: quorum sensing, *Roseobacter*, marine bacteria, biogeochemical cycles, acyl-homoserine lactones

INTRODUCTION

When acting as coordinated communities, bacterial populations are able to influence their local environment in manners that are unachievable by individual cells. It has been widely reported that phylogenetically diverse bacteria use genetic regulatory systems, known as quorum sensing (QS) systems, to coordinate gene expression in a population density dependent manner (e.g., Fuqua et al., 2001; Pappas et al., 2004; Case et al., 2008; Ng and Bassler, 2009). Among other things, QS is hypothesized to facilitate maximal access to available nutrients through the use of exoenzymes (Vetter et al., 1998; Schimel and Weintraub, 2003), the colonization of desirable niches (Nadell et al., 2008, 2009), and competitive advantages against other organisms (Folcher et al., 2001; Chin-a-Woeng et al., 2003; Barnard et al., 2007). The chemical mediators of QS are often small molecular weight diffusible molecules (Fuqua et al., 2001; Churchill and Chen, 2011). A well-characterized type of QS uses *N*-acyl-homoserine lactones (AHLs) and appears exclusive to Proteobacteria (Case et al., 2008). Canonical AHL-QS systems produce and respond to AHLs using two proteins that mediate signal production and response, LuxI and LuxR-like proteins, respectively (Nealson et al., 1970; Ruby, 1996). The genes encoding these two proteins are often located adjacent to one another on the chromosome (Fuqua et al., 1996; Churchill and Chen, 2011; Gelencsér et al., 2012). LuxI-like proteins synthesize AHLs by cyclizing *S*-adenosyl methionine into a lactone ring and the addition of an acylated carbon chain from fatty acid biosynthesis pathways (Schaefer et al., 1996). Chain length and modification at the third carbon (either -H, -OH, or -O) allow for species or group specificity (Schaefer et al., 1996; Fuqua et al., 2001). LuxR-like proteins are response regulators that mediate the expression of genes required for communal

behavior in response to intracellular concentrations of cognate AHLs (Fuqua and Winans, 1994; Fuqua et al., 1996). Activated LuxR proteins often upregulate *luxI* transcription to enhance the rate of AHL synthesis, increasing AHL concentrations, and also modulate the expression of other genes (Fuqua et al., 1996, 2001; Case et al., 2008).

AHL-based QS is common in Proteobacteria, which are abundant in coastal marine systems (Dang and Lovell, 2002; Waters and Bassler, 2005; Ng and Bassler, 2009). One of the most abundant and biogeochemically active groups of marine α -proteobacteria is the *Roseobacter* clade (Gonzalez and Moran, 1997; Buchan et al., 2005). Roseobacters can comprise up to 30% of the total 16S rRNA genes in coastal environments and up to 15% in the open ocean (Buchan et al., 2005; Wagner-Dobler and Bibel, 2006). In coastal salt marshes, roseobacters are the primary colonizers of surfaces and mediate a wide range of biogeochemically relevant processes, including mineralization of plant-derived compounds and transformations of reduced inorganic and organic sulfur compounds (Gonzalez and Moran, 1997; Dang and Lovell, 2000; Buchan et al., 2005; Dang et al., 2008). Here, we describe some of the most compelling recent research that focuses on QS in the *Roseobacter* clade, provide a genomic perspective of QS systems in roseobacters, and highlight areas for further investigation.

ROSEOBACTERS AND QUORUM SENSING

QS was first reported in roseobacters associated with marine snow and hypothesized to contribute to the ability of group members to colonize particulate matter in the ocean (Gram et al., 2002). Subsequent studies further demonstrated that roseobacters are prolific colonizers of a variety of marine surfaces, both

inert and living, and the contribution of QS to this ability and other physiologies is of growing interest (Dang and Lovell, 2002; Berger et al., 2011; Zan et al., 2012). Characterized *Roseobacter* isolates produce diverse AHL structures with acyl chains ranging from eight to eighteen carbons in length that display varying degrees of saturation as well as all three possible oxidation states (-H, -OH, or -O) at the third carbon (for structures see Gram et al., 2002; Wagner-Dobler et al., 2005; Cicirelli et al., 2008; Mohamed et al., 2008; Thiel et al., 2009; Berger et al., 2011; Zan et al., 2012). The production of AHLs has been detected by LuxR-LacZ fusion bioreporters and mass spectrometry for several isolates (Gram et al., 2002; Wagner-Dobler et al., 2005; Martens et al., 2007; Thiel et al., 2009; Berger et al., 2011; Zan et al., 2012). Of the 43 publicly available *Roseobacter* genomes, only five lack annotated *luxI* homologs: *Oceanicola batsensis* HTCC2597, *Oceanicola* sp. S124, *Pelagibaca bermudensis* HTCC2601, Rhodobacterales bacterium HTCC2255, and *Ruegeria* sp. TM1040. All except HTCC2255, however, have *luxR* homologs (Table A2). Thus far, experimental studies of QS have primarily focused on isolated representatives of the *Ruegeria-Phaeobacter* branch of the *Roseobacter* clade, with the exception of the description of a diunsaturated long chain AHL produced by *Jannaschia helgolandensis* (Thiel et al., 2009), a survey of 31 AHL producing isolates (Wagner-Dobler et al., 2005), and a recent analysis of QS in *Dinoroseobacter shibae*, where QS was shown to control motility, expression of a type IV secretion system, and whether the cells divided by binary fission or budding (Patzelt et al., 2013).

Culture-based studies of bacterial symbionts of marine sponges suggest that roseobacters are the primary producers of AHLs in these systems (Taylor et al., 2004). A model for sponge-associated roseobacters has been established using *Ruegeria* sp. KLH11 (Zan et al., 2011). Studies with this strain have been informative in providing insight into the contributions of QS to host-bacterial interactions. KLH11 contains two sets of *luxRI* homologs, designated *ssaRI* (RKLH11_1559 and RKLH11_2275) and *ssbRI* (RKLH11_1933 and RKLH11_260), and a recently discovered orphan *luxI*, designated *ssCI*, that is not annotated in the publicly available KLH11 genome. While orphan *luxI* have not been widely described in the literature, they are best described as *luxI* homologs that are not immediately adjacent to a corresponding *luxR* homolog on the chromosome. It has been proposed that *ssCI* is a recent duplication of *ssBI* (Zan et al., 2012). Heterologous expression of *SsaI*, *SsbI*, and *SscI* in *Escherichia coli* showed that they predominantly produce long chain saturated and unsaturated AHLs (C12-16). More specifically, *SsaI* produces 3O-AHL variants whereas *SsbI* and *SscI* produce 3OH-AHLs (Zan et al., 2012). The modification at the third carbon has been shown to affect the binding affinity of signaling molecules to LuxR homologs, and may allow KLH11 to finely tune its metabolism to cellular density and AHL diversity (Koch et al., 2005). KLH11 mutants deficient in QS display impaired motility, which corresponds to decreased transcription of genes encoding flagella biosynthesis machinery. The QS and motility impaired mutants form drastically thicker biofilms, suggesting when motility or QS is retarded, biofilm formation is increased (Zan et al., 2012). This

may also suggest that biofilm formation may not be directly controlled by QS, but that when quorum is achieved, motility and biofilm dispersion are induced. Recent work has shown a phosphorelay system that controls motility in KLH11 is induced by QS (Zan et al., 2013). A similar phenotype has been observed in other roseobacters, and this trend may extend across the *Ruegeria-Phaeobacter* subgroup (Bruhn et al., 2006; Dobretsov et al., 2007).

QS-mediated physiologies have been implicated in one of the few examples of roseobacters demonstrating antagonistic behavior toward a eukaryotic host. *Nautella* (formerly *Ruegeria*) sp. R11 readily colonizes the macroalga *Delisea pulchra* resulting in bleaching and subsequent death (Case et al., 2011; Fernandes et al., 2011). To combat infection, *D. pulchra* produces halogenated furanones, which have been shown to block AHL-based QS systems in many bacterial species. Active synthesis of furanones prevents macroalgal colonization by epiphytic bacteria, including *Nautella* sp. R11. However, in the absence of halogen substrates required for furanone biosynthesis, colonization occurs rapidly (Manefield et al., 1999; Hentzer et al., 2002; Defoirdt et al., 2007). Further, it appears furanones may be effective against other potentially pathogenic *Ruegeria* spp. (Zhong et al., 2003).

QS is closely connected to antimicrobial production in several roseobacters. In *Phaeobacter* sp. strain Y4I, the regulatory controls dictating the production of the antimicrobial compound indigoidine are complex and include QS. Indigoidine production confers a competitive advantage to Y4I when grown in co-culture with *Vibrio fischeri*. Transposon insertions in either of two separate *luxRI*-like systems leads to an inability of Y4I mutants to produce wildtype levels of indigoidine and an inability to inhibit the growth of *V. fischeri*. This indicates a role for both QS systems in the synthesis of indigoidine (Cude et al., 2012). The presence of multiple QS systems in the genomes of many roseobacters suggests multi-layered control is a common feature to regulate energy intensive processes, including secondary metabolite production.

Tropodithietic acid (TDA) is a broad spectrum antimicrobial produced by multiple roseobacters in response to QS (Bruhn et al., 2005; Porsby et al., 2008; Berger et al., 2011). Genome analyses of *Phaeobacter gallaeciensis* strains isolated from geographically distant locations suggest they are capable of producing both AHLs and TDA (Thole et al., 2012). *P. gallaeciensis* 2.10 has been suggested to produce TDA in response to AHLs while colonizing the marine alga *Ulva australis*, thus protecting the alga from bacterial, fungal, and larval pathogens (Rao et al., 2007). A closely related strain, *P. gallaeciensis* DSM17395, which has also been shown to colonize *U. australis* (Thole et al., 2012), produces *N*-3-hydroxydecanoyl-homoserine lactone (3OHC10-HSL) using the LuxI homolog PgaI. 3OHC10-HSL activates the adjacent regulator, PgaR, in a concentration dependent manner, which leads to the upregulation of a TDA biosynthetic operon (Berger et al., 2011). Interestingly, in a Δ *pgal* strain of DSM17395, addition of exogenous TDA is sufficient to upregulate TDA biosynthesis machinery, suggesting that regulation of TDA biosynthesis may involve multiple signals in some strains (Berger et al., 2011). The dual role of TDA as an autoinducer and an antimicrobial has also




FIGURE 1 | Continued

FIGURE 1 | Maximum likelihood phylogenetic trees of *Roseobacter* LuxR- (A) and LuxI-like (B) deduced amino acid sequences (see Appendix for details). Strain designations are shown and gene locus tags of the corresponding gene sequences can be found in **Table A1**. The scale bar represents the substitutions per sequence position. The *Roseobacter* clade number is represented in parentheses after the organism name and follows the classification system identified in Newton et al., 2010. Proposed designations

of LuxR and LuxI subgroups in roseobacters are indicated by Greek character subscript and color. Bootstrap values <50% (from 1000 iterations) are shown at branch nodes. Sequences designated with a closed pentagon indicate organisms that have been shown experimentally, by either bioreporters or mass spectrometry, to produce AHLs (Wagner-Dobler et al., 2005; Rao et al., 2006; Bruhn et al., 2007; Berger et al., 2011; Case et al., 2011; Zan et al., 2012). Sequences designated with a circle are non-roseobacters.

been demonstrated in *Ruegeria* sp. TM1040, which lacks AHL-based QS (Geng and Belas, 2010). Collectively, these data show that in addition to AHLs, roseobacters use novel autoinducers. In fact, recent investigations into novel non-fatty acyl-HSLs have shown that at least one *Roseobacter*, *Ruegeria pomeroyi* DSS-3, is capable of producing *p*-coumaroyl-homoserine lactone when grown in the presence of the aromatic lignin breakdown product *p*-coumaric acid (Schaefer et al., 2008). This discovery raises the possibility that many novel signaling molecules could be produced by roseobacters in response to available local substrates, specifically plant-derived aromatics which are primary growth substrates for roseobacters (Buchan et al., 2000; Gulvik and Buchan, 2013). The production of specific signaling molecules in response to exogenously supplied substrates suggest a single signal may convey information about both population density and environmental conditions (i.e., availability of a substrate that serves as both a source of organic nutrients and a colonizable surface), which would dictate a specific set of behaviors.

QUORUM SENSING GENE HOMOLOGY AND TOPOLOGY

To understand the relatedness of AHL-based QS systems in roseobacters, we performed a phylogenetic reconstruction of the LuxI- and their neighboring LuxR-like sequences in 38 *Roseobacter* genomes. As solo LuxR homologs have been found to bind a variety of ligands, including non-AHL molecules from eukaryotic organisms (Pappas et al., 2004; Subramoni and Venturi, 2009), it is difficult to infer their contribution in AHL-based QS. Thus, *luxR* genes that are not adjacent to *luxI* genes were not included in this analysis, but they are listed in **Table A2**. Likely a result of the close relatedness of clade members and instances of horizontal gene transfer (HGT), many of the LuxR- and LuxI-like proteins analyzed show high sequence similarity and can be grouped together (**Figures 1A,B**). Our phylogenetic trees suggests there are four LuxR-like (designated R_{α} , R_{β} , R_{γ} , and R_{δ}) and four LuxI-like protein types (designated I_{α} , I_{β} , I_{γ} , and I_{δ}) found in most sequenced roseobacters, though more sequence variants may be discovered as more genome sequences become available.

Genome analysis demonstrates that multiple conserved QS gene topologies are present within sequenced roseobacters, allowing for classification by sequence similarity and gene orientation (**Figure 2** and **Table A1**). The most conserved gene topologies are the A and B groups, of which 28 different *Roseobacter* genomes contain one of the orientations, and three *Ruegeria* genomes contain both. Genomes that contain the A topology have highly similar LuxI and LuxR sequences (>63 and >70% similarity, respectively) and its presence in three different roseobacter subclades (defined in Newton et al., 2010) may be suggestive of

HGT (**Figures 1A,B**). Genomes with topology A share a Trigger Factor (TF) encoding gene downstream from *luxRI* (**Figure 2**). The location of this TF is conserved in seven genomes. Though the function has not been examined in roseobacters, in *Vibrio cholera*, TFs play a role in the folding and secretion of proteins (Ludlam et al., 2004). The LuxI and LuxR of the A topology have been designated I_{α} and R_{α} , respectively (**Figure 2**).

The B topology is the most prevalent among the sequenced roseobacters and is found in four variations in 24 genomes (**Table A1**). Like the A topology, the LuxI and LuxR protein sequences are highly similar (>73%) between the organisms that contain the B topology. This topology is found in members of all five *Roseobacter* subclades identified by Newton et al. (2010) (**Figures 1A,B**). The LuxI and LuxR of the B topology have been labeled I_{β} and R_{β} , respectively (**Figure 2**). The conserved regions of the B topology include genes encoding a crotonyl-CoA reductase preceding *luxRI* and a putative ATP-dependent helicase following *luxRI*. In some organisms, crotonyl-CoA reductase interconverts unsaturated crotonyl-CoA to saturated butyryl-CoA as a precursor to fatty acid biosynthesis (Wallace et al., 1995). The helicase may be involved in DNA repair, protein degradation, or gene regulation (Snider et al., 2008). The B1 subgroup is the most abundant orientation within the B group, and contains a short-chain dehydrogenase following the helicase. This gene orientation is conserved in 14 *Roseobacter* genomes. Short-chain dehydrogenases are a large family of proteins that modify carbon chains of many substrates (Joernvall et al., 1995). The protein encoded by this gene may function to modify AHL biosynthesis substrates before or after AHL production.

Variations of the D topology are found in six *Roseobacter* genomes, all belonging to members of the *Roseobacter* subclade 4 (**Figures 1A,B**). These LuxI and LuxR proteins share >52 and >64% sequence similarity, respectively. The LuxI and LuxR of the D topology have been designated I_{γ} and R_{δ} (**Figure 2**). This topology shares two genes in common between the variations, *fliG* in the opposite orientation upstream of *luxRI* and an adenylosuccinate lyase encoding gene downstream. In *E. coli*, FliG is the flagellar motor switch that controls the spin direction of flagella (Roman et al., 1993). The characterized role of QS and motility in roseobacters was addressed previously (Zan et al., 2012), but none of the organisms containing the D topology have been investigated with respect to QS. The direct connection between QS and flagellar machinery may be an interesting avenue for future investigation. The other gene in this orientation putatively encodes an adenylosuccinate lyase, which is important in the *de novo* purine biosynthetic pathway and in controlling the levels of AMP and fumarate inside the cell (Tsai et al., 2007), suggesting purine biosynthesis may respond to QS.




FIGURE 2 | The gene orientation of all putative *luxRI* operons in available *Roseobacter* genomes. Green arrows represent AHL synthase encoding genes (*luxI*), orange arrows represent AHL binding response regulators (*luxR*), and black arrows represent adjacent genes. Abbreviations used: Crotonyl CoA, Crotonyl CoA reductase; HK, histidine kinase; HK/RR, hybrid histidine kinase/response regulator; HK/REC, histidine kinase with REC domain; CHP, conserved hypothetical protein; RND, RND multidrug efflux pump; Sig B, sigma B factor; SCD, short chain dehydrogenase; Trans., transposase; L17, L17 component of the 50S ribosomal protein; 2CRS, two-component regulatory system; TctA, TctA family transmembrane transporter; *mdlB*, mandelate dehydrogenase; *mdlB*; *spaβ*, α and β subunits of sulfopyruvate decarboxylase. R_x and I_x designations above the response regulators and AHL synthases indicate their corresponding phylogenetic subgroupings in **Figures 1A,B**, respectively. Those without R_x and I_x designations indicate unique sequences not found in the conserved groupings. The corresponding genomes that contain these topologies can be found in **Table A1**.

The presence of orphan *luxI* genes appears common, especially in the *Sulfitobacter*, *Ruegeria*, and *Phaeobacter* genera (Table A1). The synteny of these *luxI* and their adjacent genes is conserved in the H, I, and J topologies. In organisms that have these three orientations, there is a *luxI*-like gene of the I_8 . The *LuxI* of these topologies share >52% sequence similarity. Shared among the H, I, and J topologies are different types of putative histidine kinase (HK) encoding genes upstream of the orphan *luxI*, suggesting the protein is part of a two-component phosphorelay (Dutta et al., 1999; Stock et al., 2000). These genes are in the same direction as the *luxI* in H and I and in the opposite in J (Figure 2). In *Vibrio harveyi*, the hybrid two-component HK *LuxN* has been shown to activate gene circuits that lead to coordinated behaviors, such as bioluminescence, in response to AHLs (Freeman and Bassler, 1999; Laub and Goulian, 2007). The HKs found these topologies share modest identity with the *Vibrio harveyi* *LuxN* ($\leq 26\%$) suggesting similar regulatory systems may be present in roseobacters. While the similarity of gene sequence does not directly predict regulatory cascades or phenotypes, the development of model systems for each of these topologies will prove valuable for comparative studies across lineage members.

FUTURE DIRECTIONS

The repertoire of chemical signals in roseobacters is anticipated to be large and result in complex chemical signaling pathways in lineage members, some of which may contribute to interspecies interactions and should be investigated further. For example, uncharacterized roseobacters have been shown to be epibionts of the abundant cyanobacterial lineage *Trichodesmium*. While AHL-based interactions between *Trichodesmium* and select epibionts have been shown to stimulate mechanisms for phosphorus acquisition in this host (Hmelo et al., 2012; Van Mooy et al., 2012), a definitive role for roseobacters in this symbiosis has not yet been demonstrated. Similarly, it has been hypothesized that QS plays a role in the switch from mutualistic to antagonistic behavior proposed for *P. gallaeciensis* in its interactions with the phytoplankter *Emiliana huxleyi* (Seyedsayamdost et al., 2011). Finally, the relationships roseobacters have with vascular plants as they colonize plant material and transform plant-derived compounds

(Buchan et al., 2000; Dang and Lovell, 2000; Buchan et al., 2001) is suggestive of inter-kingdom communication, such as that found in other α -proteobacteria [e.g., *Agrobacterium tumefaciens* and *Sinorhizobium meliloti* (Hughes and Sperandio, 2008)]. Research in these areas would help elucidate the role of QS in the ability of roseobacters to colonize and interact with a diverse group of organisms.

The presence of orphan *luxR*-like genes in Proteobacterial genomes has been widely described, and their gene products have been shown to respond to AHLs and other molecules produced by other QS systems in the same organism or by other organisms (Malott et al., 2009; Patankar and González, 2009; Sabag-Daigle et al., 2012). Furthermore, it is possible that these LuxR family proteins bind structurally similar molecules that are not related to QS. In fact, it has been shown that cross-domain signaling can be mediated through LuxR homologs that bind non-AHL eukaryotic molecules (Subramoni and Venturi, 2009). In contrast, detailed studies of orphan *luxI*-like gene products are rare and are an area ripe for study. Perhaps either novel non-LuxR-like proteins or proteins encoded by genes located in distal regions of the genome (Table A2) respond to the orphan LuxI-derived AHLs. Undoubtedly, more detailed characterization of such systems will lead to a better understanding of their biological roles in roseobacters as well as other lineages.

To date, experimental studies of QS in relatively few select roseobacters have revealed complex and multi-layered control mechanisms as well as novel signaling molecules. In addition to expanding our knowledge of these characterized systems, it is our hope that future studies also broaden our understanding of currently under investigated systems within the clade and their contribution to complex multi-species interactions.

REFERENCES

- Barnard, A. M. L., Bowden, S. D., Burr, T., Coulthurst, S. J., Monson, R. E., and Salmond, G. P. C. (2007). Quorum sensing, virulence and secondary metabolite production in plant soft-rotting bacteria. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 362, 1165–1183. doi: 10.1098/rstb.2007.2042
- Berger, M., Neumann, A., Schulz, S., Simon, M., and Brinkhoff, T. (2011). Tropodithietic acid production in *Phaeobacter gallaeciensis* is regulated by N-acyl homoserine lactone-mediated quorum sensing. *J. Bacteriol.* 193, 6576–6585. doi: 10.1128/JB.05818-11
- Bruhn, J. B., Gram, L., and Belas, R. (2007). Production of antibacterial compounds and biofilm formation by *Roseobacter* species are influenced by culture conditions. *Appl. Environ. Microbiol.* 73, 442–450. doi: 10.1128/AEM.02238-06
- Bruhn, J. B., Haagensen, J. A., Bagge-Ravn, D., and Gram, L. (2006). Culture conditions of *Roseobacter* strain 27-4 affect its attachment and biofilm formation as quantified by real-time PCR. *Appl. Environ. Microbiol.* 72, 3011–3015. doi: 10.1128/AEM.72.4.3011-3015.2006
- Bruhn, J. B., Nielsen, K. F., Hjelm, M., Hansen, M., Bresciani, J., Schulz, S., and Gram, L. (2005). Ecology, inhibitory activity, and morphogenesis of a marine antagonistic bacterium belonging to the *Roseobacter* clade. *Appl. Environ. Microbiol.* 71, 7263–7270. doi: 10.1128/AEM.71.11.7263-7270.2005
- Buchan, A., Collier, L. S., Neidle, E. L., and Moran, M. A. (2000). Key aromatic-ring-cleaving enzyme, protocatechuate 3, 4-dioxygenase, in the ecologically important marine *Roseobacter* lineage. *Appl. Environ. Microbiol.* 66, 4662–4672. doi: 10.1128/AEM.66.11.4662-4672.2000
- Buchan, A., Gonzalez, J. M., and Moran, M. A. (2005). Overview of the marine *Roseobacter* lineage. *Appl. Environ. Microbiol.* 71, 5665–5677. doi: 10.1128/AEM.71.10.5665-5677.2005
- Buchan, A., Neidle, E. L., and Moran, M. A. (2001). Diversity of the ring-cleaving dioxygenase gene *pcaH* in a salt marsh bacterial community. *Appl. Environ. Microbiol.* 67, 5801–5809. doi: 10.1128/AEM.67.12.5801-5809.2001
- Case, R. J., Labbate, M., and Kjelleberg, S. (2008). AHL-driven quorum-sensing circuits: their frequency and function among the Proteobacteria. *ISME J.* 2, 345–349. doi: 10.1038/ismej.2008.13
- Case, R. J., Longford, S. R., Campbell, A. H., Low, A., Tujula, N., Steinberg, P. D., et al. (2011). Temperature induced bacterial virulence and bleaching disease in a chemically defended marine macroalga. *Environ. Microbiol.* 13, 529–537. doi: 10.1111/j.1462-2920.2010.02356.x
- Chin-a-Woeng, T. F. C., Bloemberg, G. V., and Lugtenberg, B. J. J. (2003). Phenazines and their role in biocontrol by *Pseudomonas* bacteria. *New Phytol.* 157, 503–523. doi: 10.1046/j.1469-8137.2003.00686.x
- Churchill, M. E. A., and Chen, L. L. (2011). Structural basis of acyl-homoserine lactone-dependent signaling. *Chem. Rev.* 111, 68–85. doi: 10.1021/cr1000817
- Cicirelli, E. M., Williamson, H., Tait, K., and Fuqua, C. (2008). “Acylated homoserine lactone signaling in marine bacterial systems,” in *Chemical Communication Among Bacteria*. eds S. C. Winans and B. L. Bassler (Washington, DC: ASM Press), 251–272.
- Cude, W. N., Mooney, J., Tavanaei, A. A., Hadden, M. K., Frank, A. M., Gulvik, C. A., et al. (2012). Production of the antimicrobial secondary metabolite indigoidine contributes to competitive surface colonization by the marine roseobacter *Phaeobacter* sp. strain Y4I. *Appl. Environ. Microbiol.* 78, 4771–4780. doi: 10.1128/AEM.00297-12
- Dang, H. Y., Li, T. G., Chen, M. N., and Huang, G. Q. (2008). Cross-ocean distribution of Rhodobacterales bacteria as primary surface colonizers in temperate coastal marine waters. *Appl. Environ. Microbiol.* 74, 52–60. doi: 10.1128/AEM.01400-07
- Dang, H. Y., and Lovell, C. R. (2000). Bacterial primary colonization and early succession on surfaces in marine waters as determined by amplified rRNA gene restriction analysis and sequence analysis of 16S rRNA genes. *Appl. Environ. Microbiol.* 66, 467–475. doi: 10.1128/AEM.66.2.467-475.2000
- Dang, H. Y., and Lovell, C. R. (2002). Seasonal dynamics of particle-associated and free-living marine Proteobacteria in a salt marsh tidal creek as determined using fluorescence *in situ* hybridization. *Environ. Microbiol.* 4, 287–295. doi: 10.1046/j.1462-2920.2002.00295.x
- Defoirdt, T., Miyamoto, C. M., Wood, T. K., Meighen, E. A., Sorgeloos, P., Verstraete, W., et al. (2007). The natural furanone (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone disrupts quorum sensing-regulated gene expression in *Vibrio harveyi* by decreasing the DNA-binding activity of the transcriptional regulator protein LuxR. *Environ. Microbiol.* 9, 2486–2495. doi: 10.1111/j.1462-2920.2007.01367.x
- Dobretsov, S., Dahms, H.-U., Yili, H., Wahl, M., and Qian, P.-Y. (2007). The effect of quorum-sensing blockers on the formation of marine microbial communities and larval attachment. *FEMS Microbiol. Ecol.* 60, 177–188. doi: 10.1111/j.1574-6941.2007.00285.x
- Dutta, R., Qin, L., and Inouye, M. (1999). Histidine kinases: diversity of domain organization. *Mol. Microbiol.* 34, 633–640. doi: 10.1046/j.1365-2958.1999.01646.x
- Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32, 1792–1797. doi: 10.1093/nar/gkh340
- Fernandes, N., Case, R. J., Longford, S. R., Seyedsayamdost, M. R., Steinberg, P. D., Kjelleberg, S., et al. (2011). Genomes and virulence factors of novel bacterial pathogens causing bleaching disease in the marine red alga *Delisea pulchra*. *PLoS ONE* 6:e27387. doi: 10.1371/journal.pone.0027387
- Folcher, M., Gaillard, H., Nguyen, L. T., Nguyen, K. T., Lacroix, P., Bamas-Jacques, N., et al. (2001). Pleiotropic functions of a *Streptomyces pristinaespiralis* autoregulator receptor in development, antibiotic biosynthesis, and expression of a superoxide dismutase. *J. Biol. Chem.* 276, 44297–44306. doi: 10.1074/jbc.M101109200
- Freeman, J. A., and Bassler, B. L. (1999). Sequence and function of LuxU: a two-component phosphorelay protein that regulates quorum sensing in *Vibrio harveyi*. *J. Bacteriol.* 181, 899–906.
- Fuqua, C., Parsek, M. R., and Greenberg, E. P. (2001). Regulation of gene expression by cell-to-cell communication: acyl-homoserine lactone quorum sensing. *Annu. Rev. Genet.* 35, 439–468. doi: 10.1146/annurev.genet.35.102401.090913
- Fuqua, C., Winans, S. C., and Greenberg, E. P. (1996). Census and consensus in bacterial ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators. *Annu. Rev. Microbiol.* 50, 727–751. doi: 10.1146/annurev.micro.50.1.727

- Fuqua, W. C., and Winans, S. C. (1994). A LuxR-LuxI type regulatory system activates *Agrobacterium* Ti plasmid conjugal transfer in the presence of a plant tumor metabolite. *J. Bacteriol.* 176, 2796–2806.
- Gelencsér, Z., Choudhary, K. S., Coutinho, B. G., Hudaiberdiev, S., Galbáts, B., Venturi, V., et al. (2012). Classifying the topology of AHL-driven quorum sensing circuits in Proteobacterial genomes. *Sensors* 12, 5432–5444. doi: 10.3390/s120505432
- Geng, H., and Belas, R. (2010). Expression of tropodithetic acid biosynthesis is controlled by a novel autoinducer. *J. Bacteriol.* 192, 4377–4387. doi: 10.1128/JB.00410-10
- Gonzalez, J. M., and Moran, M. A. (1997). Numerical dominance of a group of marine bacteria in the alpha-subclass of the class Proteobacteria in coastal seawater. *Appl. Environ. Microbiol.* 63, 4237–4242.
- Gram, L., Grossart, H. P., Schlingloff, A., and Kiorboe, T. (2002). Possible quorum sensing in marine snow bacteria: production of acylated homoserine lactones by *Roseobacter* strains isolated from marine snow. *Appl. Environ. Microbiol.* 68, 4111–4116. doi: 10.1128/AEM.68.8.4111-4116.2002
- Gulvik, C. A., and Buchan, A. (2013). Simultaneous catabolism of plant-derived aromatic compounds results in enhanced growth for members of the *Roseobacter* lineage. *Appl. Environ. Microbiol.* 79, 3716–3723. doi: 10.1128/AEM.00405-13
- Hentzer, M., Riedel, K., Rasmussen, T. B., Heydorn, A., Andersen, J. B., Parsek, M. R., et al. (2002). Inhibition of quorum sensing in *Pseudomonas aeruginosa* biofilm bacteria by a halogenated furanone compound. *Microbiology* 148, 87–102. Available online at: <http://mic.sgmjournals.org/content/148/1/87.long>
- Hmelo, L. R., Van Mooy, B. A. S., and Mincer, T. J. (2012). Characterization of bacterial epibionts on the cyanobacterium *Trichodesmium*. *Aquat. Microb. Ecol.* 67, 488–496. doi: 10.3354/ame01571
- Hughes, D. T., and Sperandio, V. (2008). Inter-kingdom signalling: communication between bacteria and their hosts. *Nat. Rev. Microbiol.* 6, 111–120. doi: 10.1038/nrmicro1836
- Joernvall, H., Persson, B., Krook, M., Atrian, S., Gonzalez-Duarte, R., Jeffery, J., et al. (1995). Short-chain dehydrogenases/reductases (SDR). *Biochemistry* 34, 6003–6013. doi: 10.1021/bi00018a001
- Koch, B., Liljefors, T., Persson, T., Nielsen, J., Kjelleberg, S., and Givskov, M. (2005). The LuxR receptor: the sites of interaction with quorum-sensing signals and inhibitors. *Microbiology* 151, 3589–3602. doi: 10.1099/mic.0.27954-0
- Laub, M. T., and Goulian, M. (2007). Specificity in two-component signal transduction pathways. *Annu. Rev. Genet.* 41, 121–145. doi: 10.1146/annurev.genet.41.042007.170548
- Ludlam, A. V., Moore, B. A., and Xu, Z. (2004). The crystal structure of ribosomal chaperone trigger factor from *Vibrio cholerae*. *Proc. Natl. Acad. Sci. U.S.A.* 101, 13436–13441. doi: 10.1073/pnas.0405868101
- Malott, R. J., O'grady, E. P., Toller, J., Inhülsen, S., Eberl, L., and Sokol, P. A. (2009). A *Burkholderia cenocepacia* orphan luxR homolog is involved in quorum-sensing regulation. *J. Bacteriol.* 191, 2447–2460. doi: 10.1128/JB.01746-08
- Manefield, M., De Nys, R., Naresh, K., Roger, R., Givskov, M., Peter, S., et al. (1999). Evidence that halogenated furanones from *Delisea pulchra* inhibit acylated homoserine lactone (AHL)-mediated gene expression by displacing the AHL signal from its receptor protein. *Microbiology* 145, 283–291. doi: 10.1099/13500872-145-2-283
- Martens, T., Gram, L., Grossart, H.-P., Kessler, D., Muller, R., Simon, M., et al. (2007). Bacteria of the *Roseobacter* clade show potential for secondary metabolite production. *Microb. Ecol.* 54, 31–42. doi: 10.1007/s00248-006-9165-2
- Mohamed, N. M., Cicirelli, E. M., Kan, J., Chen, F., Fuqua, C., and Hill, R. T. (2008). Diversity and quorum-sensing signal production of Proteobacteria associated with marine sponges. *Environ. Microbiol.* 10, 75–86. doi: 10.1111/j.1462-2920.2007.01431.x
- Nadell, C. D., Xavier, J. B., and Foster, K. R. (2009). The sociobiology of biofilms. *FEMS Microbiol. Rev.* 33, 206–224. doi: 10.1111/j.1574-6976.2008.00150.x
- Nadell, C. D., Xavier, J. B., Levin, S. A., and Foster, K. R. (2008). The evolution of quorum sensing in bacterial biofilms. *PLoS Biol.* 6:e14. doi: 10.1371/journal.pbio.0060014
- Nealson, K. H., Platt, T., and Hastings, J. W. (1970). Cellular control of the synthesis and activity of the bacterial luminescent system. *J. Bacteriol.* 104, 313–322.
- Newton, R. J., Griffin, L. E., Bowles, K. M., Meile, C., Gifford, S., Givens, C. E., et al. (2010). Genome characteristics of a generalist marine bacterial lineage. *ISME J.* 4, 784–798. doi: 10.1038/ismej.2009.150
- Ng, W.-L., and Bassler, B. L. (2009). Bacterial quorum-sensing network architectures. *Annu. Rev. Genet.* 43, 197–222. doi: 10.1146/annurev-genet-102108-134304
- Pappas, K. M., Weingart, C. L., and Winans, S. C. (2004). Chemical communication in proteobacteria: biochemical and structural studies of signal synthases and receptors required for intercellular signalling. *Mol. Microbiol.* 53, 755–769. doi: 10.1111/j.1365-2958.2004.04212.x
- Patankar, A. V., and González, J. E. (2009). An orphan luxR homolog of *Sinorhizobium meliloti* affects stress adaptation and competition for nodulation. *Appl. Environ. Microbiol.* 75, 946–955. doi: 10.1128/AEM.01692-08
- Patzelt, D., Wang, H., Buchholz, I., Rohde, M., Grobe, L., Pradella, S., et al. (2013). You are what you talk: quorum sensing induces individual morphologies and cell division modes in *Dinoroseobacter shibae*. *ISME J.* doi: 10.1038/ismej.2013.107. Available online at: <http://dx.doi.org/10.1038/ismej.2013.107>
- Porsby, C. H., Nielsen, K. F., and Gram, L. (2008). *Phaeobacter* and *ruegeria* species of the *Roseobacter* clade colonize separate niches in a danish turbot (*Scophthalmus maximus*)-rearing farm and antagonize *Vibrio anguillarum* under different growth conditions. *Appl. Environ. Microbiol.* 74, 7356–7364. doi: 10.1128/AEM.01738-08
- Rao, D., Webb, J. S., Holmstrom, C., Case, R., Low, A., Steinberg, P., et al. (2007). Low densities of epiphytic bacteria from the marine alga *Ulva australis* inhibit settlement of fouling organisms. *Appl. Environ. Microbiol.* 73, 7844–7852. doi: 10.1128/AEM.01543-07
- Rao, D., Webb, J. S., and Kjelleberg, S. (2006). Microbial colonization and competition on the marine alga *Ulva australis*. *Appl. Environ. Microbiol.* 72, 5547–5555. doi: 10.1128/AEM.00449-06
- Roman, S. J., Frantz, B. B., and Matsumura, P. (1993). Gene sequence, overproduction, purification and determination of the wild-type level of the *Escherichia coli* flagellar switch protein FlhG. *Gene* 133, 103–108. doi: 10.1016/0378-1119(93)90232-R
- Ruby, E. G. (1996). Lessons from a cooperative, bacterial-animal association: the *Vibrio fischeri*–*Euprymna scolopes* light organ symbiosis. *Annu. Rev. Microbiol.* 50, 591–624. doi: 10.1146/annurev.micro.50.1.591
- Sabag-Daigle, A., Soares, J. A., Smith, J. N., Elmasry, M. E., and Ahmer, B. M. (2012). The acyl homoserine lactone (AHL) receptor, SdiA, of *E. coli* and *Salmonella* does not respond to indole, but at high concentrations indole can interfere with AHL detection. *Appl. Environ. Microbiol.* 78, 5424–5431. doi: 10.1128/AEM.00046-00012
- Schaefer, A. L., Greenberg, E. P., Oliver, C. M., Oda, Y., Huang, J. J., Bittan-Banin, G., et al. (2008). A new class of homoserine lactone quorum-sensing signals. *Nature* 454, U595–U596. doi: 10.1038/nature07088
- Schaefer, A. L., Val, D. L., Hanzelka, B. L., Cronan, J. E., and Greenberg, E. P. (1996). Generation of cell-to-cell signals in quorum sensing: acyl homoserine lactone synthase activity of a purified *Vibrio fischeri* LuxI protein. *Proc. Natl. Acad. Sci. U.S.A.* 93, 9505–9509. doi: 10.1073/pnas.93.18.9505
- Schimel, J. P., and Weintraub, M. N. (2003). The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biol. Biochem.* 35, 549–563. doi: 10.1016/S0038-0717(03)00015-4
- Seyedsayamdost, M. R., Case, R. J., Kolter, R., and Clardy, J. (2011). The Jekyll-and-Hyde chemistry of *Phaeobacter gallaeciensis*. *Nat. Chem.* 3, 331–335. doi: 10.1038/nchem.1002
- Snider, J., Thibault, G., and Houry, W. A. (2008). The AAA+ superfamily of functionally diverse proteins. *Genome Biol.* 9, 216. doi: 10.1186/gb-2008-9-4-216
- Stock, A. M., Robinson, V. L., and Goudreau, P. N. (2000). Two-component signal transduction. *Annu. Rev. Biochem.* 69, 183–215. doi: 10.1146/annurev.biochem.69.1.183
- Subramoni, S., and Venturi, V. (2009). LuxR-family 'solos': bachelor sensors/regulators of signalling molecules. *Microbiology* 155, 1377–1385. doi: 10.1099/mic.0.026849-0
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., and Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739. doi: 10.1093/molbev/msr121
- Taylor, M. W., Schupp, P. J., Baillie, H. J., Charlton, T. S., De Nys, R., Kjelleberg, S., et al. (2004). Evidence for acyl homoserine lactone signal production in bacteria associated with marine sponges. *Appl. Environ. Microbiol.* 70, 4387–4389. doi: 10.1128/AEM.70.7.4387-4389.2004

- Thiel, V., Kunze, B., Verma, P., Wagner-Dobler, I., and Schulz, S. (2009). New structural variants of homoserine lactones in bacteria. *Chembiochem* 10, 1861–1868. doi: 10.1002/cbic.200900126
- Thole, S., Kalhoefer, D., Voget, S., Berger, M., Engelhardt, T., Liesegang, H., et al. (2012). *Phaeobacter gallaeciensis* genomes from globally opposite locations reveal high similarity of adaptation to surface life. *ISME J.* 6, 2229–2244. doi: 10.1038/ismej.2012.62
- Tsai, M., Koo, J., Yip, P., Colman, R. F., Segall, M. L., and Howell, P. L. (2007). Substrate and product complexes of *Escherichia coli* adenylosuccinate lyase provide new insights into the enzymatic mechanism. *J. Mol. Biol.* 370, 541–554. doi: 10.1016/j.jmb.2007.04.052
- Van Mooy, B. A. S., Hmelo, L. R., Sofen, L. E., Campagna, S. R., May, A. L., Dyhrman, S. T., et al. (2012). Quorum sensing control of phosphorus acquisition in *Trichodesmium* consortia. *ISME J.* 6, 422–429. doi: 10.1038/ismej.2011.115
- Vetter, Y. A., Deming, J. W., Jumars, P. A., and Krieger-Brockett, B. B. (1998). A predictive model of bacterial foraging by means of freely released extracellular enzymes. *Microb. Ecol.* 36, 75–92. doi: 10.1007/s002489900095
- Wagner-Dobler, I., and Bibel, H. (2006). Environmental biology of the marine *Roseobacter* lineage. *Annu. Rev. Microbiol.* 60, 255–280. doi: 10.1146/annurev.micro.60.080805.142115
- Wagner-Dobler, I., Thiel, V., Eberl, L., Allgaier, M., Bodor, A., Meyer, S., et al. (2005). Discovery of complex mixtures of novel long-chain quorum sensing signals in free-living and host-associated marine alphaproteobacteria. *Chembiochem* 6, 2195–2206. doi: 10.1002/cbic.200500189
- Wallace, K. K., Bao, Z.-Y., Dai, H., Digate, R., Schuler, G., Speedie, M. K., et al. (1995). Purification of crotonyl-CoA reductase from *Streptomyces collinus* and cloning, sequencing and expression of the corresponding gene in *Escherichia coli*. *Eur. J. Biochem.* 233, 954–962. doi: 10.1111/j.1432-1033.1995.954_3.x
- Waters, C. M., and Bassler, B. L. (2005). Quorum sensing: cell-to-cell communication in bacteria. *Annu. Rev. Cell Dev. Biol.* 21, 319–346. doi: 10.1146/annurev.cellbio.21.012704.131001
- Zan, J., Fricke, W. F., Fuqua, C., Ravel, J., and Hill, R. T. (2011). Genome sequence of *Ruegeria* sp. strain KLH11, an N-acylhomoserine lactone-producing bacterium isolated from the marine sponge *Mycale laxissima*. *J. Bacteriol.* 193, 5011–5012. doi: 10.1128/JB.05556-11
- Zan, J., Heindl, J. E., Liu, Y., Fuqua, C., and Hill, R. T. (2013). The CckA-ChpT-CtrA phosphorelay system is regulated by quorum sensing and controls flagellar motility in the marine sponge symbiont *Ruegeria* sp. KLH11. *PLoS ONE* 8:e66346. doi: 10.1371/journal.pone.0066346
- Zan, J. D., Cicirelli, E. M., Mohamed, N. M., Sibhatu, H., Kroll, S., Choi, O., et al. (2012). A complex LuxR-LuxI type quorum sensing network in a roseobacterial marine sponge symbiont activates flagellar motility and inhibits biofilm formation. *Mol. Microbiol.* 85, 916–933. doi: 10.1111/j.1365-2958.2012.08149.x
- Zhong, Z. P., Caspi, R., Helinski, D., Knauf, V., Sykes, S., O'byrne, C., et al. (2003). Nucleotide sequence based characterizations of two cryptic plasmids from the marine bacterium *Ruegeria* isolate PR1b. *Plasmid* 49, 233–252. doi: 10.1016/S0147-619X(03)00014-3

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 12 June 2013; accepted: 24 October 2013; published online: 12 November 2013.

Citation: Cude WN and Buchan A (2013) Acyl-homoserine lactone-based quorum sensing in the *Roseobacter* clade: complex cell-to-cell communication controls multiple physiologies. *Front. Microbiol.* 4:336. doi: 10.3389/fmicb.2013.00336

This article was submitted to *Aquatic Microbiology*, a section of the journal *Frontiers in Microbiology*.

Copyright © 2013 Cude and Buchan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

APPENDIX

Maximum likelihood phylogenetic trees of LuxI-like and genetically linked LuxR-like sequences from 38 published roseobacter genomes were constructed. Protein alignments of the LuxI and LuxR homologs were done using the MUSCLE algorithm with default parameters (Edgar, 2004), and manually curated. The phylogenetic trees were generated using MEGA 5.2 following published methods (Hall, 2013). The Maximum Likelihood statistical method was used with the WAG model of amino acid substitution and gamma distribution with invariant sites (G+I) selected. Gaps were handled with a 95% partial deletion data treatment, and the phylogeny was tested with 1000 bootstrap replications (Tamura et al., 2011). Bootstrap values are reported in percentages and shown at nodes where values are >50%. Groups were divided and defined by natural divisions in the trees and gene topology in the genome (Figure 2). The LuxRI protein sequences of *Vibrio fischeri* (Accession: AAQ90231.1 and AAP22376.1) were used to root the trees. LuxR and LuxI homologs of six proteobacterial species with sequence similarity to at least one roseobacter sequence in each

subgroup (>30% identity) were included in the alignments to assess the validity of the groupings. The non-roseobacter LuxRI included were: *Sinorhizobium meliloti* (Accession: ABC88593.1 and CAC46417.1), *Bradyrhizobium elkanii* (Accession: WP_018273827.1 and WP_018272735), *Rhizobium leguminosarum* (Accession: YP_002281222.1 and CAD20929.1), *Agrobacterium tumefaciens* (Accession: WP_003501811.1 and AAZ50597.1), *Pseudomonas putida* (Accession: CAO85746.1 and CAO85747), *Pseudomonas aeruginosa* (CAO85753.1 and CAO85754.1).

REFERENCES

- Hall, B. G. (2013). Building phylogenetic trees from molecular data with MEGA. *Mol. Biol. Evol.* 30, 1229–1235. doi: 10.1093/molbev/mst012
- Newton, R. J., Griffin, L. E., Bowles, K. M., Meile, C., Gifford, S., Givens, C. E., et al. (2010). Genome characteristics of a generalist marine bacterial lineage. *ISME J.* 4, 784–798. doi: 10.1038/ismej.2009.150
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., and Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739. doi: 10.1093/molbev/msr121

Table A1 | Paired LuxRI and orphan LuxI^a homologs identified in the 38 sequenced roseobacters.

Strains	Gene orientation	<i>luxR</i> gene locus	<i>luxI</i> gene locus
Rhodobacterales bacterium HTCC2083	A	RB2083_3272	RB2083_3255
<i>Ruegeria</i> sp. KLH11	A	RKLH11_1559	RKLH11_2275
<i>Roseovarius</i> sp. 217	A	ROS217_18272	ROS217_18267
<i>Roseovarius</i> sp. TM1035	A	RTM1035_10475	RTM1035_10485
<i>Ruegeria lacuscaerulensis</i> ITI-1157	A	SL1157_2477	SL1157_2476
<i>Ruegeria pomeroyi</i> DSS-3	A	SPO2286	SPO2287
<i>Ruegeria</i> sp. TW15	A	RTW15_010100013877	RTW15_010100013872
<i>Citricella</i> sp. 357	B	C357_10197	C357_10192
<i>Citricella</i> SE45	B	CSE45_4055	CSE45_4054
<i>Roseobacter denitrificans</i> OCh 114	B	RD1_1638	RD1_1639
<i>Sagittula stellata</i> E-37	B	SSE37_11169	SSE37_11164
<i>Ruegeria pomeroyi</i> DSS-3	B	SPO0371	SPO0372
<i>Dinoroseobacter shibae</i> DFL 12	B1	DSHI_2852	DSHI_2851
<i>Loktanella</i> sp. SE62	B1	LSE62_0618	LSE62_0617
<i>Phaeobacter gallaeciensis</i> 2.10	B1	PGA2_c03430	PGA2_c03440
<i>Phaeobacter gallaeciensis</i> DSM 17395	B1	PGA1_c03880	PGA1_c03890
<i>Phaeobacter gallaeciensis</i> ANG1	B1	ANG1_1316	ANG1_1315
<i>Phaeobacter</i> sp. Y4I	B1	RB4I_1689	RB4I_1631
<i>Ruegeria</i> sp. KLH11	B1	RKLH11_1933	RKLH11_260
Rhodobacterales bacterium HTCC2150	B1	RB2150_14426	RB2150_14421
<i>Roseobacter</i> sp. AzwK-3b	B1	RAZWK3B_04270	RAZWK3B_04275
<i>Roseobacter</i> sp. GAI101	B1	RGAI101_376	RGAI101_3395
<i>Roseobacter</i> sp. MED193	B1	MED193_10428	MED193_10423
<i>Ruegeria lacuscaerulensis</i> ITI-1157	B1	SL1157_0613	SL1157_0612
<i>Ruegeria</i> sp. R11	B1	RR11_2850	RR11_2520
<i>Ruegeria</i> sp. TW15	B1	RTW15_010100017779	RTW15_010100017784
<i>Roseobacter</i> sp. R2A57	B2	R2A57_2403	R2A57_2404
<i>Thalassibium</i> R2A620	B2	TR2A62_3165	TR2A62_3166 TR2A62_3167
<i>Maritimibacter alkaliphilus</i> HTCC2654	B3	RB2654_09024	RB2654_09014
Rhodobacterales bacterium HTCC2083	B4	RB2083_3265	RB2083_730
<i>Roseobacter litoralis</i> Och 149	B4	RLO149_c030690	RLO149_c030680
<i>Dinoroseobacter shibae</i> DFL 12	C	DSHI_0311	DSHI_0312
<i>Jannaschia</i> sp. CCS1	C	JANN_0619	JANN_0620
	D	SKA53_05835	SKA53_05830
<i>Loktanella vestfoldensis</i> SKA53		SKA53_05840	
<i>Loktanella</i> sp. SE62	D1	LSE62_3230 LSE62_3229	LSE62_3231
<i>Oceanicola granulosus</i> HTCC2516	D1	OG2516_02284 OG2516_02289	OG2516_02294
<i>Octadecabacter antarcticus</i> 307	D1	OA307_2044 OA307_3216	OA307_4586
<i>Roseobacter</i> sp. CCS2	D1	RCCS2_02083 RCCS2_02088	RCCS2_02078
<i>Octadecabacter arcticus</i> 238	D2	OA238_4151 OA238_3367	OA238_2886
<i>Roseobacter</i> sp. SK209-2-6	E	RSK20926_22079	RSK20926_22084

(Continued)

Table A1 | Continued

Strains	Gene orientation	<i>luxR</i> gene locus	<i>luxI</i> gene locus
<i>Sulfitobacter</i> NAS-14.1	E	NAS141_01141	NAS141_01136
<i>Maritimibacter alkaliphilus</i> HTCC2654	F	RB2654_20053	RB2654_20048
<i>Roseovarius</i> sp. 217	G	ROS217_01405	ROS217_01410
<i>Roseobacter litoralis</i> Och 149	G1	RLO149_c036220	RLO149_c036210
<i>Sulfitobacter</i> NAS-14.1	H		NAS141_00695
<i>Sulfitobacter</i> sp. EE-36	H		EE36_01635
<i>Roseovarius nubinhibens</i> ISM	I		ISM_03755
<i>Oceanibulbus indolifex</i> HEL45	I		OIH45_00955
<i>Ruegeria</i> sp. R11	J		RR11_2017
<i>Roseobacter</i> sp. MED193	J		MED193_08053
<i>Ruegeria</i> sp. TW15	J		RTW15_010100005486
<i>Dinoroseobacter shibae</i> DFL 12	K		DSH1_4152
<i>Phaeobacter gallaeciensis</i> 2.10	L	PGA2_c18970	PGA2_c18960
<i>Phaeobacter</i> sp. Y4I	L1	RB4I_1027	RB4I_3464
<i>Phaeobacter gallaeciensis</i> 2.10	M		PGA2_c07460
<i>Phaeobacter gallaeciensis</i> DSM 17395	M		PGA1_c07680
Rhodobacterales bacterium HTCC2150	N	RB2150_11281	RB2150_11291
<i>Roseobacter litoralis</i> Och 149	O		RLO149_c036590
<i>Roseobacter</i> sp. AzwK-3b	P		RAZWK3B_19371
<i>Roseobacter</i> sp. SK209-2-6	Q	RSK20926_15126	RSK20926_15131
<i>Roseobacter</i> sp. GAI101	Q1		RGAI101_1101
<i>Ruegeria lacuscaerulensis</i> ITI-1157	R		SL1157_1706
<i>Ruegeria</i> sp. TrichCH4B	S		SCH4B_1938

Homologs of *LuxI* encoding genes were determined using *BlastP* to characterized proteins^b (E -value < e^{-3}) on Roseobase (www.roseobase.org) and are consistent with the genome annotations. The *LuxR* gene loci listed do not represent all homologs within the genomes, but were determined based using *BlastP* with the autoinducer binding domain sequence from *Pfam* (PF03472) on Roseobase, and proximity to *luxI* homologs. These were also consistent with genome annotations. Gene orientations are represented in **Figure 2**.

^aOrphan *luxI* homologs are defined as those that do not have an immediately adjacent *luxR* gene. All reported orphan *luxI* genes are located and at least 100 kb from the end of the draft genome contig.

^b*Vibrio fischeri* *LuxI* (AAP22376), *Agrobacterium tumefaciens* *TraR* (AAZ50597) and *Phaeobacter gallaeciensis* *Pgal* (YP_006571842).

Table A2 | Putative orphan LuxR encoding genes that do not have an adjacent *luxI* on the chromosome.

Strains	<i>luxR</i> gene locus
<i>Citricella</i> sp. 357	C357_03001
<i>Citricella</i> sp. SE45	CSE45_1818 CSE45_4969
<i>Dinoroseobacter shibae</i> DFL 12	Dshi_1550 Dshi_1815 Dshi_1819
<i>Jannaschia</i> sp. CCS1	Jann_1153 Jann_2301 Jann_3193
<i>Loktanella</i> sp. SE62	LSE62_3779
<i>Maritimibacter alkaliphilus</i> HTCC2654	RB2654_10983 RB2654_03619
<i>Oceanibulbus indolifex</i> HEL-45	OIHEL45_01695 OIHEL45_02625 OIHEL45_13145
<i>Oceanicola batsensis</i> HTCC2597	OB2597_03302
<i>Oceanicola granulosis</i> HTCC2516	OG2516_08027
<i>Oceanicola</i> sp. S124	OS124_010100017942 OS124_010100007975
<i>Octadecabacter antarcticus</i> 238	OA238_3367 OA238_3623
<i>Octadecabacter antarcticus</i> 307	OA307_2044
<i>Pelagibaca bermudensis</i> HTCC2601	R2601_24964 R2601_10664
<i>Phaeobacter gallaeciensis</i> 2.10	PGA2_c15480 PGA2_c18970
<i>Phaeobacter gallaeciensis</i> DSM 17395	PGA1_c15590
<i>Phaeobacter gallaeciensis</i> ANG1	ANG1_869
<i>Phaeobacter</i> sp. Y4I	RB41_896
<i>Rhodobacterales bacterium</i> HTCC2083	RB2083_1776
<i>Rhodobacterales bacterium</i> HTCC2150	RB2150_02239
<i>Roseobacter denitrificans</i> OCh 114	RD1_3967
<i>Roseobacter litoralis</i> OCh 149	RLO149_c004710 RLO149_c036470
<i>Roseobacter</i> sp. AzwK-3b	RAZWK3B_15865
<i>Roseobacter</i> sp. CCS2	RCCS2_00422
<i>Roseobacter</i> sp. GAI101	RGAI101_670
<i>Roseobacter</i> sp. MED193	MED193_03932
<i>Roseobacter</i> sp. R2A57	R2A57_3570
<i>Roseobacter</i> sp. SK209-2-6	RSK20926_03972 RSK20926_18892
<i>Roseovarius nubinhibens</i> ISM	ISM_09921 ISM_15650
<i>Roseovarius</i> sp. TM1035	RTM1035_08219
<i>Roseovarius</i> sp. 217	ROS217_20327
<i>Ruegeria pomeroyi</i> DSS-3	SPO1974
<i>Ruegeria</i> sp. KLH11	RKHL11_1390
<i>Ruegeria</i> sp. R11	RR11_2316
<i>Ruegeria</i> sp. TM1040	TM1040_3102 TM1040_1212

(Continued)

Table A2 | Continued

Strains	<i>luxR</i> gene locus
<i>Ruegeria</i> sp. TW15	RTW15_010100007191
<i>Ruegeria</i> sp. TrichCH4B	SCH4B_0463 SCH4B_4179 SCH4B_4368 SCH4B_4682
<i>Ruegeria lacuscaerulensis</i> ITI-1157	SL1157_2844
<i>Sagittula stellata</i> E-37	SSE37_06082
<i>Sulfitobacter</i> sp. EE-36	EE36_03628
<i>Sulfitobacter</i> sp. NAS-14.1	NAS141_08556
<i>Thalassiosibium</i> sp. R2A62	TR2A62_0664

Homologs of *LuxR* encoding genes were determined using BlastP with the autoinducer binding domain sequence from Pfam (PF03472) on Roseobase (www.roseobase.org).