

2012

Field Application of Sustained-Yield Harvest Management for Northern Bobwhite in Texas

Joseph P. Sands
Texas A&M University

Matthew J. Schnupp
Texas A&M University

Trent W. Teinert
Texas A&M University

Stephen J. DeMaso
U.S. Fish and Wildlife Service

Fidel Hernandez
Texas A&M University

See next page for additional authors

Follow this and additional works at: <https://trace.tennessee.edu/nqsp>

Recommended Citation

Sands, Joseph P.; Schnupp, Matthew J.; Teinert, Trent W.; DeMaso, Stephen J.; Hernandez, Fidel; Brennan, Leonard A.; Rollins, Dale; and Perez, Robert M. (2012) "Field Application of Sustained-Yield Harvest Management for Northern Bobwhite in Texas," *National Quail Symposium Proceedings*: Vol. 7 , Article 83.

Available at: <https://trace.tennessee.edu/nqsp/vol7/iss1/83>

This Bobwhite Harvest Management is brought to you for free and open access by Volunteer, Open Access, Library Journals (VOL Journals), published in partnership with The University of Tennessee (UT) University Libraries. This article has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor. For more information, please visit <https://trace.tennessee.edu/nqsp>.

Field Application of Sustained-Yield Harvest Management for Northern Bobwhite in Texas

Authors

Joseph P. Sands, Matthew J. Schnupp, Trent W. Teinert, Stephen J. DeMaso, Fidel Hernandez, Leonard A. Brennan, Dale Rollins, and Robert M. Perez

FIELD APPLICATION OF SUSTAINED-YIELD HARVEST MANAGEMENT FOR NORTHERN BOBWHITE IN TEXAS

Joseph P. Sands^{1,2}

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

Matthew J. Schnupp³

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

Trent W. Teinert⁴

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

Stephen J. DeMaso

U.S. Fish and Wildlife Service, Gulf Coast Joint Venture, 700 Cajundome Boulevard, Lafayette, LA 70506, USA

Fidel Hernández

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

Leonard A. Brennan

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

Dale Rollins

Texas AgriLife Research, Texas A&M University System, San Angelo, TX 76901, USA

Robert M. Perez

Texas Parks and Wildlife Department, La Vernia, TX 78121, USA

ABSTRACT

Sustained-yield harvest (SYH) is considered a potentially viable strategy for managing harvest of northern bobwhites (*Colinus virginianus*). However, application of SYH has not been evaluated for northern bobwhites. We evaluated the application of using SYH as a harvest management strategy for bobwhite during the 2007–2008 and 2008–2009 hunting seasons in 2 ecoregions of Texas (Rolling Plains, South Texas Plains). We collected field data at 3 study sites/ecoregion (900–1,900 ha each; 2 hunted sites and 1 control) to estimate 4 demographic parameters (fall and spring density, overwinter survival in the absence of hunting, and harvest rate). We used these data to parameterize the additive harvest model for bobwhites and compare predictions of spring abundance of the model with field estimates. The additive harvest model, compared to field estimates, consistently underestimated spring population density (mean \pm SE) by $55.7 \pm 17.8\%$ (2007–2008) and $34.1 \pm 4.9\%$ (2008–2009) in the Rolling Plains, and by $26.4 \pm 25.3\%$ (2007–2008) and $49.1 \pm 2.1\%$ (2008–2009) in the South Texas Plains. Implementing SYH in the field, despite its potential benefits, will be challenging given the need for reliable estimates of 3 key population parameters (fall and spring density, and natural mortality in the absence of hunting) and the high variation often associated with them. Conservative harvest prescriptions based on the lower 95% CIs of fall density estimates may permit sustainable harvest despite variation in density estimates.

Citation: Sands, J. P., M. J. Schnupp, T. W. Teinert, S. J. DeMaso, F. Hernández, L. A. Brennan, D. Rollins, and R. M. Perez. 2012. Field application of sustained-yield harvest management for northern bobwhite in Texas. *Proceedings of the National Quail Symposium* 7:162.

Key words: *Colinus virginianus*, northern bobwhite, South Texas, sustained-yield harvest

¹ E-mail: joseph_sands@fws.gov

² U.S. Fish and Wildlife Service, Migratory Birds and Habitat Programs, 911 NE 11th Avenue, Portland OR 97232, USA.

³ Present address: King Ranch Inc., Highway 141 West, Kingsville, TX 78363, USA.

⁴ Present address: Texas Parks and Wildlife Department, 2805 North Navarro, Suite 600B, Victoria, TX 77901, USA.