



National Quail Symposium Proceedings

Volume 8

Article 51

2017

Summer Survival of Translocated Northern Bobwhite in the New Jersey Pine Barrens: Preliminary Results

Philip M. Coppola
University of Delaware

Kaili R. Stevens
University of Delaware

Christopher K. Williams
University of Delaware

Theron M. Terhune II
Tall Timbers Research Station and Land Conservancy

John P. Parke
New Jersey Audubon

See next page for additional authors

Follow this and additional works at: <http://trace.tennessee.edu/nqsp>

 Part of the [Natural Resources and Conservation Commons](#)

Recommended Citation

Coppola, Philip M.; Stevens, Kaili R.; Williams, Christopher K.; Terhune, Theron M. II; Parke, John P.; and Cecil, John (2017) "Summer Survival of Translocated Northern Bobwhite in the New Jersey Pine Barrens: Preliminary Results," *National Quail Symposium Proceedings*: Vol. 8 , Article 51.
Available at: <http://trace.tennessee.edu/nqsp/vol8/iss1/51>

This Bobwhite Translocation is brought to you for free and open access by Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Summer Survival of Translocated Northern Bobwhite in the New Jersey Pine Barrens: Preliminary Results

Authors

Philip M. Coppola, Kaili R. Stevens, Christopher K. Williams, Theron M. Terhune II, John P. Parke, and John Cecil

SUMMER SURVIVAL OF TRANSLOCATED NORTHERN BOBWHITE IN THE NEW JERSEY PINE BARRENS: PRELIMINARY RESULTS

Philip M. Coppola¹

Department of Entomology and Wildlife Ecology, University of Delaware, 250 Townsend Hall, Newark, DE 19716, USA

Kaili R. Stevens

Department of Entomology and Wildlife Ecology, University of Delaware, 250 Townsend Hall, Newark, DE 19716, USA

Christopher K. Williams

Department of Entomology and Wildlife Ecology, University of Delaware, 250 Townsend Hall, Newark, DE 19716, USA

Theron M. Terhune

Tall Timbers Research Station and Land Conservancy, 13093 Henry Beadel Drive Tallahassee, FL 32312, USA

John P. Parke

New Jersey Audubon, Wattles Stewardship Center 1024 Anderson Road, Port Murray, NJ 07865, USA

John Cecil

New Jersey Audubon, Wattles Stewardship Center 1024 Anderson Road, Port Murray, NJ 07865, USA

ABSTRACT

Northern bobwhite (*Colinus virginianus*) have declined precipitously since the 1960s, largely due to habitat deterioration and changes in land use; some of the highest declines have been observed in the Mid-Atlantic States. In other regions, attempts to augment bobwhite populations have been relatively successful using translocation. As part of a long-term restocking program, focal areas for translocation in the mid-Atlantic region were identified by biologists at a National Bobwhite Conservation Initiative (NBCI) workshop. The objective of this project is to evaluate translocation to restore bobwhite populations in the New Jersey Pine Barrens, a focal area designated with a high ranking for potential bobwhite recovery. The study site, Pine Island Cranberry Co., is the largest privately owned land tract (>6,000 hectares) in New Jersey, with a mix of shortleaf pine (*Pinus echinata*), pitch pine (*P. rigida*), scrub oak (*Quercus ilicifolia*), and early successional forbes and grasses. For three consecutive years (2015–2017) prior to breeding season, we are translocating eighty radio-collared bobwhite (40 male, 40 female) from wild populations in southwest Georgia. These individuals are radio-located 3–5 times per week, year round. We are collecting microhabitat measurements (e.g., groundcover, understory, and canopy closure) and monitoring nests to characterize habitat use, nest site selection, and nest fate. Survival is estimated using staggered-entry Kaplan-Meier analyses and a Cox proportional hazard model in R to determine covariates of daily mortality. Six of 14 nests were successful in summer 2015 (66 known hatches), and 0 of 12 nests were successful in summer 2016. Snake depredation was the cause of 41.7% of failed nests in 2016. Preliminary analyses produce a five-month adult survival rate of 0.455 (SE = 0.138) for summer 2015 and 0.270 (SE = 0.0516) for 2016. Our planned third summer (2017) of data collection will increase our understanding of these disparate survival estimates.

Citation: Coppola, P. M., K. R. Stevens, C. K. Williams, T. M. Terhune, J. P. Parke, and J. Cecil. 2017. Summer survival of translocated northern bobwhite in the New Jersey Pine Barrens: Preliminary results. National Quail Symposium Proceedings 8:185.

Key words: *Colinus virginianus*, northern bobwhite, translocation, radio-telemetry, nesting, survival, habitat use, habitat management

¹E-mail: pcoppola@udel.edu

© 2017 [Coppola, Stevens, Williams, Terhune, Parke and Cecil] and licensed under CC BY-NC 4.0.