



University of Tennessee, Knoxville
Trace: Tennessee Research and Creative Exchange

Faculty Publications and Other Works -- Biomedical
and Diagnostic Sciences

Veterinary Medicine -- Faculty Publications and
Other Works

9-15-2017

Antimicrobial resistance patterns of Staphylococcus species isolated from cats presented at a veterinary academic hospital in South Africa

Daniel Nenene Qekwana
University of Pretoria

Dikeledi Sebola
University of Pretoria

James Wabwire Oguttu
University of South Africa

Agricola Odoi
University of Tennessee, Knoxville, aodoi@utk.edu

Follow this and additional works at: http://trace.tennessee.edu/utk_compmedpubs

Recommended Citation

Qekwana, Daniel Nenene, Dikeledi Sebola, James Wabwire Oguttu, and Agricola Odoi. "Antimicrobial Resistance Patterns of Staphylococcus Species Isolated from Cats Presented at a Veterinary Academic Hospital in South Africa." *BMC Veterinary Research* 13 (2017). <https://doi.org/10.1186/s12917-017-1204-3>.

This Article is brought to you for free and open access by the Veterinary Medicine -- Faculty Publications and Other Works at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Faculty Publications and Other Works -- Biomedical and Diagnostic Sciences by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

RESEARCH ARTICLE

Open Access



Antimicrobial resistance patterns of *Staphylococcus* species isolated from cats presented at a veterinary academic hospital in South Africa

Daniel Nenene Qekwana¹, Dikeledi Sebola¹, James Wabwire Oguttu² and Agricola Odoi^{1,3*} 

Abstract

Background: Antimicrobial resistance is becoming increasingly important in both human and veterinary medicine. This study investigated the proportion of antimicrobial resistant samples and resistance patterns of *Staphylococcus* isolates from cats presented at a veterinary teaching hospital in South Africa. Records of 216 samples from cats that were submitted to the bacteriology laboratory of the University of Pretoria academic veterinary hospital between 2007 and 2012 were evaluated. Isolates were subjected to antimicrobial susceptibility testing against a panel of 15 drugs using the disc diffusion method. Chi square and Fisher's exact tests were used to assess simple associations between antimicrobial resistance and age group, sex, breed and specimen type. Additionally, associations between *Staphylococcus* infection and age group, breed, sex and specimen type were assessed using logistic regression.

Results: *Staphylococcus* spp. isolates were identified in 17.6% (38/216) of the samples submitted and 4.6% (10/216) of these were unspciated. The majority (61.1%,11/18) of the isolates were from skin samples, followed by otitis media (34.5%, 10/29). Coagulase Positive *Staphylococcus* (CoPS) comprised 11.1% (24/216) of the samples of which 7.9% (17/216) were *S. intermedius* group and 3.2% (7/216) were *S. aureus*. Among the Coagulase Negative *Staphylococcus* (CoNS) (1.9%, 4/216), *S. felis* and *S. simulans* each constituted 0.9% (2/216). There was a significant association between *Staphylococcus* spp. infection and specimen type with odds of infection being higher for ear canal and skin compared to urine specimens. There were higher proportions of samples resistant to clindamycin 34.2% (13/25), ampicillin 32.4% (2/26), lincospectin 31.6% (12/26) and penicillin-G 29.0% (11/27). Sixty three percent (24/38) of *Staphylococcus* spp. were resistant to one antimicrobial agent and 15.8% were multidrug resistant (MDR). MDR was more common among *S. aureus* 28.6% (2/7) than *S. intermedius* group isolates 11.8% (2/17). One *S. intermedius* group isolate was resistant to all β -lactam antimicrobial agents tested.

Conclusion: *S. intermedius* group was the most common cause of skin infections and antimicrobial resistance was not wide spread among cats presented at the veterinary academic hospital in South Africa. However, the presence of MDR-*Staphylococcus* spp. and isolates resistant to all β -lactams is of both public health and animal health concern.

Keywords: *Staphylococcus* spp., Antimicrobial resistance, Veterinary hospital, Cats, South Africa

* Correspondence: aodoi@utk.edu

¹University of Pretoria, Faculty of Veterinary Science, Department of Paraclinical Sciences, Section Veterinary Public Health, Pretoria, Gauteng, South Africa

³University of Tennessee, College of Veterinary Medicine, Department of Biomedical and Diagnostic Sciences, Knoxville, TN, USA

Full list of author information is available at the end of the article



Background

Although *Staphylococcus* are commensals of the skin, mucous membranes, alimentary and urogenital tracts of a diverse group of mammals and birds, they have been implicated in clinical infections of humans and animals [1–3]. Transmission of *Staphylococcus* between animals and humans are known to occur [1, 4]. Cats have been reported as carriers of both Coagulase positive (CoPS) and coagulase negative *Staphylococcus* species (CoNS) [2, 3, 5–7]. However, coagulase positive *Staphylococcus* species infections seem to be more prominent in feline medicine than CoNS infections [1]. Among the CoPS species in cats, *S. pseudintermedius* are the most common followed by *S. aureus* [5, 8]. These infections have been associated with pyoderma, postoperative wound infections and otitis [9]. In addition, *S. felis*, is a cause of urinary tract infections [10].

Although resistance to β -lactam antimicrobials among *Staphylococcus* isolates from cats has been reported [6, 8], other antimicrobial agents such as gentamycin, enrofloxacin and doxycycline have been reported to be effective against *Staphylococcus* infections in cats [5, 11, 12]. However, information on the proportion of antimicrobial resistant isolates and resistance patterns of *Staphylococcus* species in clinical cases of cats in developing economies in general and South Africa in particular is very limited. Therefore, the objective of this study was to investigate the proportion of antimicrobial resistant isolates and resistance patterns among *Staphylococcus* species isolates from cat samples submitted to a veterinary academic hospital in South Africa between 2007 and 2012.

Methods

Data collection

Data containing records of cat samples submitted to the University of Pretoria Bacteriology Laboratory at the Veterinary Teaching Hospital in South Africa between January 2007 and December 2012 for routine diagnostic tests were evaluated. The following variables were captured: breed, age, sex, specimen type, staphylococcus species isolated, antimicrobial included in the antimicrobial susceptibility test panel and the susceptibility profile of the isolates.

Staphylococcus identification and antimicrobial susceptibility testing

Culture of samples was done using sheep blood agar incubated at 37 °C for at least 24 h. All media used were quality controlled using *S. aureus* ATCC 25923. Suspected *Staphylococcus* colonies were identified based on phenotypic characteristics including colony characteristics, catalase, D-mannitol, maltose, deoxyribonuclease (DNase) tests, polymyxin-B and Gram-staining as described by Quinn

[13]. *S. intermedius* and *S. delphini* were classified as *S. intermedius* group (SIG) as described by Sasaki et al. [14].

Isolates were subjected to antimicrobial susceptibility testing against a panel of 15 drugs using the disc diffusion method (discs supplied by Oxoid) [15]. Included in the panel were the following drugs: 30 μ g amikacin (AK), 30 μ g doxycycline (DOX30), 5 μ g enrofloxacin (ENR), 10 μ g gentamicin (CN), 10 μ g penicillin G (P), 25 μ g sulphamethoxazole-trimethoprim (SXT), 30 μ g chloramphenicol (C), 30 μ g cephalothin/lexin (KF), 30 μ g kanamycin (K), 2 μ g clindamycin/lincomycin (MY), 100 μ g lincospectin (Espectinomycine-lincomycine) (LS100), 5 μ g orbifloxacin (OBX5), 20/10 μ g synulox (Amoxicillin-Clavulanic acid) (AMC20/10) and 15 μ g tylosin (TY). The results, based on the diameter of the inhibition zones, were classified as sensitive, intermediate or resistant in accordance with the Clinical and Laboratory Standards Institute [15]. For the purposes of the study, intermediate susceptibility was considered as susceptible. Multidrug resistance (MDR) was defined as resistance to at least one antimicrobial agent in three or more antimicrobial categories [16].

Data analysis

All the statistical analyses were performed using SAS 9.4 (SAS Institute Inc., Cary, NC, USA) statistical package. The dataset was assessed for missing data and inconsistencies such as improbable values. Shapiro-Wilk test of normality was used for evaluation of distributions of age that was found to be non-normally distributed and hence median and interquartile ranges were reported. Age was also categorised into two categories: <2 years and \geq 2 years. The frequencies and proportions of all categorical variables were calculated and presented in a table. Associations between antimicrobial resistance of *Staphylococcus* spp. isolates and a number of host factors (breed, age, sex, specimen type) and other categorical variables were assessed using the Chi-square and Fisher's Exact tests. Statistical significance was assessed using a critical p -value of 0.05. The variables specimen type and breed had too many categories to include in the model in their original form and hence they were re-coded (Table 1).

Univariable and multivariable models

Investigation of the predictors of *Staphylococcus* spp. infections was done in two steps. In the first step, univariable logistic regression model was fit to assess the relationships between sex, age, specimen type and breed, and the outcome variable, *Staphylococcus* status. The potential predictors of *Staphylococcus* spp. infection at this stage were identified using a relaxed $\alpha \leq 0.20$. Thus variables with $p \leq 0.20$ in the univariable model were considered for inclusion in the multivariable model in the 2nd step. Therefore, the 2nd step involved fitting a multivariable logistic regression

Table 1 Distribution of *Staphylococcus* isolates among cat specimens tested at an academic veterinary hospital laboratory, 2007–2012

| Variable | Category | Number of cats tested | | <i>Staphylococcus</i> positive | |
|-------------------------|---------------------|-----------------------|---------|--------------------------------|---------|
| | | Frequency | Percent | Frequency | Percent |
| Breed (n = 212) | | | | | |
| | Domestic Short Hair | 132 | 62.3 | 24 | 18.2 |
| | Persian | 21 | 9.9 | 4 | 19.1 |
| | Siamese | 15 | 7.1 | 1 | 6.7 |
| | Domestic Long Hair | 12 | 5.7 | 3 | 25.0 |
| | All others | 32 | 15.1 | 6 | 18.8 |
| Sex (n = 205) | | | | | |
| | Male | 122 | 59.5 | 17 | 13.9 |
| | Female | 83 | 40.5 | 20 | 24.1 |
| Specimen type (n = 215) | | | | | |
| | Urine | 95 | 44.2 | 3 | 3.2 |
| | Ear canal swab | 29 | 13.5 | 10 | 34.5 |
| | Skin | 18 | 8.4 | 11 | 61.1 |
| | All others | 73 | 34.0 | 14 | 19.2 |
| Age (n = 216) | | | | | |
| | ≥2 years | 123 | 56.9 | 21 | 17.1 |
| | <2 years | 93 | 43.1 | 17 | 18.3 |

model using manual backwards selection method with the significance set at $\alpha \leq 0.05$.

Confounding was assessed by comparing the change in model coefficients with and without the suspected confounders. If the removal of a suspected confounding variable resulted in a 20% or greater change in another model coefficient, the removed variable was considered a confounder and retained in the model regardless of its statistical significance. In addition, two-way interaction terms between variable in the final main effects model were assessed.

Odds ratios (ORs) and their 95% confidence intervals were computed for variables included in the final model. The differences between categories of statistically significant predictors for *Staphylococcus* spp. were also assessed by changing the reference categories of the predictors. Hosmer-Lemeshow goodness-of-fit test was used to assess model fit.

Results

A total of 216 samples were submitted to the bacteriology lab during the study period, of which, 17.6% (38/216) tested positive for *Staphylococcus* spp. each of which had single isolates (i.e. no mixed infections were identified). The majority of samples tested were urine (44.2%, 95/215), followed by ear canal swab (13.5%, 29/215) and skin samples (8.4%, 18/215). Significantly ($p = 0.0065$) more samples originated from males (59.5%, 122/205) than female cats (40.5%, 83/205). Similarly, a significantly ($p = 0.0412$) higher proportion of

samples came from cats ≥ 2 years (56.9%, 123/216) compared to cats < 2 years (43.1%, 93/216). The majority of samples were obtained from the domestic short hair breed (DSH) (62.3%, 132/212), followed by Persian breed (9.9%, 21/212) (Table 1).

Staphylococcus species were isolated from several cat breeds including domestic long hair (25.0% 3/12), domestic short hair (18.2%, 24/132) and persian breed (19.1%, 4/21). Skin samples yielded the highest (61.1%, 11/18) percentage of *staphylococcus* isolates followed by ear swabs (34.5%, 10/29).

Significantly ($p = 0.02$) more CoPS (11.1%, 24/216) were isolated compared to CoNS (1.9%, 4/216). Among the CoPS, *S. intermedius* group was most predominant (7.9%, 17/216) followed by *S. aureus* (3.2%, 7/216). Equal percentage of *S. felis* (0.9%, 2/216) and *S. simulans* (0.9%; 2/216) were observed among the CoNS. Five percent (4.6%, 10/216) of the *Staphylococcus* isolates identified were not characterized (Table 2).

Staphylococcus isolates exhibited relatively high levels of resistance towards ampicillin (32.4%, 12/26), penicillin-G (29.0%, 11/27), clindamycin (34.2%, 13/25) and lincospectin (31.6%, 12/26) (Table 3). Overall, 63.2% (24/38) of *Staphylococcus* spp. were resistant to at least one antimicrobial agent and 21.1% (8/38) were multi-drug resistant (MDR). *S. aureus* (85.7%, 6/7) had the highest level of resistance to at least one antimicrobial agent followed by *S. intermedius* group (52.9%, 9/17). Similarly, *S. aureus* (42.9%, 3/7) had a higher level of MDR than *S. intermedius* group (12.5%, 2/16). One *S.*

Table 2 Distribution of *Staphylococcus* species isolated from clinical specimens from cats presented at an academic veterinary hospital between 2007 and 2012 (n = 216)

| | Isolate | Frequency | Percent (%) |
|---------------------|-----------------------------------|-----------|-------------|
| CoPS n = 24 (11.1%) | <i>S. intermedius</i> group (SIG) | 17 | 7.9 |
| | <i>S. aureus</i> | 7 | 3.2 |
| CoNS n = 4 (1.9%) | <i>S. felis</i> | 2 | 0.9 |
| | <i>S. simulans</i> | 2 | 0.9 |
| Unspeciated | <i>S. spp.</i> | 10 | 4.6 |
| | Negative | 172 | 82.4 |
| | Total | 216 | 100 |

intermedius group isolate was resistant to all β -lactam antimicrobial agents tested. This isolate was also resistant to 9 out of the 15 antimicrobial agents tested. Three *S. intermedius* group isolates were resistant to both clindamycin and lincosamides. Among the *S. aureus* isolates, one was resistant to five antimicrobial agents and two to four antimicrobial agents (Table 4).

Predictors of staphylococcus infections

Based on the univariable logistic model, only sex and specimen type stood out as potential predictors of *Staphylococcus* spp. infection based on a liberal $\alpha \leq 0.20$ (Table 5). Thus, only these two variables were assessed in the multivariable model. In the final model only

specimen type was significantly associated with staphylococcus species infection based on $\alpha \leq 0.05$. The odds of testing positive for *Staphylococcus* spp. infections were significantly higher among ear canal ($p = 0.0002$) and skin samples ($p < 0.0001$) than urine samples (Table 6). However, there was no significant differences in the odds of *Staphylococcus* spp. infection between skin and ear canal samples (Table 7).

Discussion

The aim of this study was to investigate the proportion of antimicrobial resistant isolates and resistance patterns of *Staphylococcus* spp. isolates from clinical samples obtained from cats admitted to a veterinary academic

Table 3 Antimicrobial resistance profile of *Staphylococcus* isolates to antimicrobial agents from samples tested at an academic veterinary laboratory, 2007–2012

| Group | Drug | Frequency | Percent (n/N) ^b |
|----------------------------|-------------------------------|-----------|----------------------------|
| β -lactam | | 26 | 28.9 (26/90) |
| | Penicillin | | |
| | PenicillinG | 11 | 29.0 (11/27) |
| | Ampicillin | 12 | 32.4 (12/26) |
| Cephalosporin | Cephalothin | 1 | 2.7 (1/37) |
| | Ceftiofur | 1 | 2.7 (1/37) |
| Combination | Amoxicillin/Clavulanic acid | 1 | 2.7 (1/37) |
| Tetracycline | Doxycycline | 1 | 2.7 (1/37) |
| Fluoroquinolone | | 6 | 8.6 (6/70) |
| | Enrofloxacin | 3 | 7.9 (3/35) |
| | Orbifloxacin | 3 | 7.9 (3/35) |
| Aminoglycoside | | 4 | 3.6 (4/110) |
| | Gentamicin | 1 | 2.7 (1/37) |
| | Amikacin | 1 | 2.7 (1/37) |
| | Kanamycin | 2 | 5.3 (2/36) |
| Potentiated sulfonamide | Sulfamethoxazole/trimethoprim | 4 | 10.5 (4/34) |
| Amphenicols | Chloramphenicol | 2 | 5.2 (2/36) |
| Lincosamides | Clindamycin | 13 | 34.2 (13/25) |
| Aminoglycoside-lincosamide | Lincospectin | 12 | 31.6 (12/26) |
| Macrolide | Tylosin | 2 | 5.3 (2/36) |

^b = n is the number resistant, N is number tested

Table 4 Antimicrobial resistance patterns identified in *Staphylococcus* isolates from cat specimens tested at an academic veterinary hospital laboratory, 2007–2012

| Species | Antimicrobial Resistance | | | Resistance patterns |
|-----------------------------|--------------------------|------------------------------------|-----------------------------------|--|
| | Percent (n/N) | Multidrug Resistance Percent (n/N) | B-Lactam resistance Percent (n/N) | |
| <i>S. aureus</i> | 85.7 (6/7) | 42.9 (3/7) | 0 | AMP (1), AMP PEN (1), AMP PEN LIN (1), AMP PEN CLI LIN (1), AMP PEN CHL LIN (1), AMP SP KAN CLI LIN (1) |
| <i>S. felis</i> | 0 (0/2) | 0 | 0 | |
| <i>S. intermedius</i> group | 52.9 (9/17) | 11.8 (2/17) | 5.9 (1/17) | PEN (1), KAN (1), CLI LIN (3), SP LIN (1), AMP PEN SP (1), ENR LIN OR TYL (1), AMP CEF GEN PEN SP CEF KAN OR SU (1) |
| <i>S. simulans</i> | 0 (0/2) | 0 | 0 | |
| <i>S. spp.</i> | 90.0 (9/10) | 30.0 (3/10) | 0 | CLI (2), LIN (1), CHL CLI (1), AMP PEN (1), CLI LIN OR (1), AMP AMP DOX ENR PEN CLI (1), AMP PEN CLI (1), AMP ENR PEN CLI LIN TYL (1), |

n number of resistant samples, N number of samples tested, AMP ampicillin, CEF Ceftiofur, ENR Enrofloxacin, GEN Gentamicin, PEN PenicillinG, SP Sulpha/Trimethoprim, CHL Chloramphenicol, KAN Kanamycin, CLI Clindamycin/Lincomycin, AMI Amikacin, DOX Doxycycline, LIN Lincospectin, ORB Orbifloxacin, SU Amoxicillin/Clavulanic acid, TYL Tylosin

hospital in South Africa. The proportion of *Staphylococcus* spp. isolated from cat samples in this study was relatively low (17.6%). This is not directly comparable to findings from other previous studies on cats due to differences in isolation methods (use of enrichment media in particular), and differences in study designs. In the current study, we investigated *Staphylococcus* infections in hospitalised clinical cases only. However, the majority of similar published studies of *Staphylococcus* in cats have largely focused on methicillin resistance rather than *Staphylococcus* infections in general. In addition,

past studies have focused on carriage rather than infections [8, 17].

Similar to findings from other studies [8, 17], in this study we observed that skin and ear canal samples had significantly higher odds of testing positive for *Staphylococcus* spp. than other samples. These results seem to suggest that *Staphylococcus* spp. are a major cause of skin related infections in cats [18–20]. Although there tended to be a higher proportion of *Staphylococcus* spp. isolated from the domestic short hair breeds, the final model indicated no significant association between breed and odds of *Staphylococcus* spp. infection. However, the lack of significant association might be due to small sample size involved in this study. It is worth noting that, there is evidence that certain diseases are more common in certain breeds of cats and we suspect that this might be the case with *Staphylococcus* infections [17, 21].

Table 5 Results of the univariable logistic model showing predictors of *Staphylococcus* spp. infection among cats tested at an academic veterinary hospital laboratory, 2007–2012

| Variable | OR ^a | 95% CI ^b | | p-value |
|----------------------|-----------------|---------------------|-------|---------|
| Breed | | | | |
| Domestic Long Hair | 1.4 | 0.3 | 7.0 | 0.368 |
| Domestic Short Hair | 0.9 | 0.4 | 2.6 | 0.707 |
| Persian | 1.0 | 0.3 | 4.2 | 0.723 |
| Siamese | 0.3 | 0.03 | 2.8 | 0.237 |
| All others | Ref | . | . | . |
| Sex | | | | |
| Female | 1.9 | 0.9 | 4.0 | 0.066 |
| Male | Ref | . | . | . |
| Specimen type | | | | |
| Ear canal swab | 16.1 | 4.1 | 64.2 | <0.0001 |
| Skin | 48.2 | 10.9 | 213.7 | <0.0001 |
| All others | 7.3 | 2.0 | 26.4 | 0.003 |
| Urine | Ref | . | . | . |
| Age | | | | |
| < 2 years | 1.1 | 0.5 | 2.2 | 0.818 |
| > = 2 years | Ref | . | . | . |

^aOdds ratio

^b95% Confidence Interval

Consistent with other studies [3, 5, 17, 19], we observed a higher percentage of CoPS than CoNS. This is mainly due to the observed higher percentage of *S. intermedius* group, which are CoPS, isolated in this study. On the contrary, Abraham et al. [7] reported nearly equal proportions of *S. aureus* and *S. pseudintermedius*

Table 6 Multivariable logistic model showing predictors of *Staphylococcus* spp. infection among cats tested at an academic veterinary hospital laboratory, 2007–2012

| Variable | OR ¹ | 95% CI ² | | p-value |
|----------------------|-----------------|---------------------|-------|---------|
| Sex | | | | |
| Female | 1.9 | 0.8 | 4.3 | 0.117 |
| Male | Ref | . | . | . |
| Specimen type | | | | |
| Ear canal swab | 14.8 | 3.6 | 60.5 | 0.0002 |
| Skin | 52.1 | 11.3 | 240.3 | <.0001 |
| All others | 8.4 | 2.3 | 30.7 | 0.001 |
| Urine | Ref | . | . | . |

Table 7 Final multivariable logistic model showing the results of changing reference categories of specimen type

| Variable | OR ¹ | 95% CI ² | | p-value |
|----------------|-----------------|---------------------|--------|---------|
| Specimen type | | | | |
| Skin | 3.519 | 0.969 | 12.78 | 0.0559 |
| Urine | 0.068 | 0.017 | 0.276 | 0.0002 |
| All others | 0.564 | 0.208 | 1.531 | 0.2611 |
| Ear canal swab | Ref | . | . | . |
| Ear canal swab | 0.284 | 0.078 | 1.032 | 0.0559 |
| Urine | 0.019 | 0.004 | 0.089 | <.0001 |
| All others | 0.16 | 0.05 | 0.517 | 0.0022 |
| Skin | Ref | . | . | . |
| Ear canal swab | 1.773 | 0.653 | 4.81 | 0.2611 |
| Skin | 6.237 | 1.933 | 20.131 | 0.0022 |
| Urine | 0.12 | 0.033 | 0.439 | 0.0014 |
| All others | Ref | . | . | . |

isolates from asymptomatic cats. Consistent with our study, a Brazilian study by Lilenbaum et al. [8] reported a higher percentage of *S. intermedius* group (*S. pseudintermedius*) in cats compared to other *Staphylococcus* species. This may be related to the fact that SIG especially *S. pseudintermedius* is well adapted to the skin surface of dogs and cats than *S. aureus* [22–25].

The observed higher percentage of resistance towards β -lactam and lincosamide antimicrobial agents among the *Staphylococcus* isolates in cats has previously been reported [6, 8, 23]. Of particular concern is one *S. intermedius* group isolate that was resistant to all β -lactam antimicrobial agents tested in this study. Moreover, MRSA have an intrinsic resistance to β -lactams by virtue of newly acquired low-affinity penicillin-binding protein 2A (PBP2A). Therefore, it is possible that this isolate was MRSA [26, 27]. Unfortunately, we could not assess this since the lab that supplied the data used in this study did not test for methicillin resistance. Almost 16 % (15.8%) of *Staphylococcus* isolates in this study were MDR. This is close to the 14.8% reported by Gandolfi-Decristophoris et al. [23] in Switzerland.

Since this is a retrospective study, these findings should be interpreted with caution. The history of previous use of antimicrobial agents was not included in the analysis and this could have affected the recovery rates of *Staphylococcus* species. The study also suffers from low samples size which impacted the precision of some of the estimates. Nonetheless, the results provide a useful preliminary indication of the burden and antimicrobial resistance patterns of *Staphylococcus* spp. infections in cats presented to the academic veterinary hospital in South Africa.

Conclusions

As has been observed in other studies, this study suggests that *S. intermedius* group is the most common cause of skin infections in cats investigated in this study. It also suggests that antimicrobial resistance is not so wide spread among cats presented at the veterinary academic hospital in South Africa. Considering the risk of cross-transmission of resistant organisms between cats and humans, the levels of resistance to β -lactams is of great concern from both a public health and animal health point of view. However, given the limited scope of this study, there is need for larger and more detailed primary base studies to specifically assess the extent of antimicrobial resistant infections in cats in South Africa and their role in the spread of antimicrobial drug resistance to humans.

Abbreviations

AMR: Antimicrobial resistance; CoNS: Coagulase negative staphylococci species; CoPS: Coagulase Positive *Staphylococcus*; DSH: Domestic Short Hair breed; MDR: Multidrug resistant resistance; SAS: Statistical Analysis System

Acknowledgements

The authors would like to thank the Department of Tropical Diseases and Companion Animal Clinical Studies for providing access to the records used in this study. We are also grateful to Ms. S Nxumalo and Mr. W Mbethe for helping with data entry and validation.

Funding

Not applicable.

Availability of data and materials

The data that support the findings of this study are available from the bacteriology laboratory of the University of Pretoria that has legal ownership of the data. The data are not publicly available and should be requested and obtained from the above legal owner.

Authors' contributions

DNQ was involved in study design and data management and performed all statistical analyses and interpretation as well as preparation of the manuscript draft. AO was involved in study design, data analysis and interpretation as well as extensive editing of the manuscript. JWO was involved in study design and editing of the manuscript. DS was involved in data collection and interpretation of results of the manuscript. All authors read and approved the final manuscript.

Ethics approval

The study was approved by the University of Pretoria Ethics Committee (reference number S4285–15).

Consent for publication

The study does not involve human subjects and therefore no consent was required. However, the lab that supplied the study data provided consent for study results to be published.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹University of Pretoria, Faculty of Veterinary Science, Department of Paraclinical Sciences, Section Veterinary Public Health, Pretoria, Gauteng, South Africa. ²University of South Africa, College of Agriculture and

Environmental Sciences, Department of Agriculture and Animal Health, Johannesburg, South Africa. ³University of Tennessee, College of Veterinary Medicine, Department of Biomedical and Diagnostic Sciences, Knoxville, TN, USA.

Received: 6 March 2017 Accepted: 1 September 2017

Published online: 15 September 2017

References

- Morris DO, Mauldin EA, O'Shea K, Shofer FS, Rankin SC. Clinical, microbiological, and molecular characterization of methicillin-resistant *Staphylococcus Aureus* infections of cats. *Am J Vet Res*. 2006;67:1421–5.
- Hanselman BA, Kruth SA, Rousseau J, Weese JS. Coagulase positive staphylococcal colonization of humans and their household pets. *Can Vet J*. 2009;50:954–8.
- Rich M. Staphylococci in animals: prevalence, identification and antimicrobial susceptibility, with an emphasis on methicillin-resistant *Staphylococcus Aureus*. *Br J Biomed Sci*. 2005;62:98–105.
- Duquette RA, Nuttall TJ. Methicillin-resistant *Staphylococcus Aureus* in dogs and cats: an emerging problem? *J Small Anim Pract*. Blackwell Publishing Ltd. 2004;45:591–7.
- Beça N, Bessa LJ, Mendes Â, Santos J, Leite-Martins L, Matos AJF, et al. Coagulase-positive staphylococcus: prevalence and antimicrobial resistance. *J Am Anim Hosp Assoc*. 2015;51:365–71.
- Bierowiec K, Ploneczka-Janeczko K, Rypuła K. Is the colonisation of *Staphylococcus Aureus* in pets associated with their close contact with owners? *PLoS One*. 2016;11:e0156052.
- Abraham J, Morris D, Griffith G, Shofer F, Rankin S. Surveillance of healthy cats and cats with inflammatory skin disease for colonization of the skin by methicillin-resistant coagulase-positive staphylococci and staphylococcus schleiferi ssp. schleiferi. *Dermatology*. 2007;18:252–9.
- Lilenbaum W, Nunes EL, Azeredo MA. Prevalence and antimicrobial susceptibility of staphylococci isolated from the skin surface of clinically normal cats. *Lett Appl Microbiol*. 1998;27:224–8.
- Bannoehr J, Guardabassi L. *Staphylococcus pseudintermedius* in the dog: taxonomy, diagnostics, ecology, epidemiology and pathogenicity. *Vet Dermatol*. 2012;23:253–66.
- Litster A, Moss SSM, Honnery M, Rees B, Trott DJD. Prevalence of bacterial species in cats with clinical signs of lower urinary tract disease: recognition of *Staphylococcus Felis* as a possible feline urinary tract pathogen. *Vet Microbiol*. 2007;121:182–8.
- Ganiere JP, Medaille C, Mangion C. Antimicrobial drug susceptibility of *Staphylococcus Intermedius* clinical isolates from canine pyoderma. *J Vet Med B Infect Dis Vet Public Health*. 2005;52:25–31.
- Lilenbaum W, Esteves AL, Souza GN. Prevalence and antimicrobial susceptibility of staphylococci isolated from saliva of clinically normal cats. *Lett Appl Microbiol*. Blackwell Science Ltd. 1999;28:448–52.
- Quinn PJ, Carter ME, Markey B, Carter GR. *Clinical veterinary microbiology*. Edinburgh: Mosby Wolfe; 1994.
- Sasaki T, Kikuchi K, Tanaka Y, Takahashi N, Kamata S, Hiramatsu K. Reclassification of phenotypically identified *Staphylococcus Intermedius* strains. *J Clin Microbiol*. 2007;45:2770–8.
- Clinical Institute Laboratory Standards. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals; approved standard – fourth edition. CLSI document VET01-A. 2013.
- Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect*. 2012;18:268–81.
- Loeffler A, Boag AK, Sung J, Lindsay JA, Guardabassi L, Dalsgaard A, et al. Prevalence of methicillin-resistant *Staphylococcus Aureus* among staff and pets in a small animal referral hospital in the UK. *J Antimicrob Chemother*. 2005;56:692–7.
- Scott DW, Paradis M. A survey of canine and feline skin disorders seen in a university practice: small animal clinic, University of Montréal, Saint-Hyacinthe, Québec (1987-1988). *Can Vet J*. 1990;31:830–5.
- Devriese LA, Nzuambe D, Godard C. Identification and characteristics of staphylococci isolated from lesions and normal skin of horses. *Vet Microbiol*. 1985;10:269–77.
- Biberstein EL, Jang SS, Hirsh DC. Species distribution of coagulase-positive staphylococci in animals. *J Clin Microbiol*. 1984;19:610–5.
- Lekcharoensuk C, Osborne CA, Lulich JP. Epidemiologic study of risk factors for lower urinary tract diseases in cats. *J Am Vet Med Assoc*. 2001;218:1429–35.
- Kawakami T, Shibata S, Murayama N, Nagata M, Nishifuji K, Iwasaki T, et al. Antimicrobial susceptibility and methicillin resistance in staphylococcus pseudintermedius and staphylococcus schleiferi subsp. coagulans isolated from dogs with pyoderma in Japan. *J Vet Med Sci*. 2010;72:1615–9.
- Gandolfi-Decristophoris P, Regula G, Petriani O, Zinsstag J, Schelling E. Prevalence and risk factors for carriage of multi-drug resistant staphylococci in healthy cats and dogs. *J Vet Sci*. 2013;14:449–56.
- Schmidt VM, Williams NJ, Pinchbeck G, Corless CE, Shaw S, McEwan N, et al. Antimicrobial resistance and characterisation of staphylococci isolated from healthy Labrador retrievers in the United Kingdom. *BMC Vet Res*. 2014;10:17.
- Paul NC, Bärghman SC, Moodley A, Nielsen SS, Guardabassi L. *Staphylococcus pseudintermedius* colonization patterns and strain diversity in healthy dogs: a cross-sectional and longitudinal study. *Vet Microbiol*. 2012;160:420–7.
- Guignard B, Entenza JM, Moreillon P. β -Lactams against methicillin-resistant *Staphylococcus Aureus*. *Curr Opin Pharmacol*. 2005;5:479–89.
- Lim D, Strynadka NCJ. Structural basis for the beta lactam resistance of PBP2a from methicillin-resistant *Staphylococcus Aureus*. *Nat Struct Biol*. 2002;9:870–6.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

