

University of Tennessee, Knoxville Trace: Tennessee Research and Creative Exchange

Faculty Publications and Other Works -- Biomedical and Diagnostic Sciences

Veterinary Medicine -- Faculty Publications and Other Works

Winter 9-27-2018

Complete Genome Sequences of Four Staphylococcus aureus Sequence Type 398 Isolates from Four Goats with Osteomyelitis

Mohamed A. Abouelkhair University of Tennessee, Knoxville

Stephen A. Kania University of Tennessee, Knoxville

David A. Bemis University of Tennessee, Knoxville

Rebecca E. Rifkin University of Tennessee, Knoxville

Remiqiusz M. Grzeskowiak University of Tennessee, Knoxville

See next page for additional authors

Follow this and additional works at: https://trace.tennessee.edu/utk_compmedpubs Part of the <u>Veterinary Medicine Commons</u>

Recommended Citation

Abouelkhair, Mohamed A.; Kania, Stephen A.; Bemis, David A.; Rifkin, Rebecca E.; Grzeskowiak, Remiqiusz M.; Biris, Alexandru S.; and Anderson, David A., "Complete Genome Sequences of Four Staphylococcus aureus Sequence Type 398 Isolates from Four Goats with Osteomyelitis" (2018). *Faculty Publications and Other Works -- Biomedical and Diagnostic Sciences*. https://trace.tennessee.edu/utk_compmedpubs/116

This Article is brought to you for free and open access by the Veterinary Medicine -- Faculty Publications and Other Works at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Faculty Publications and Other Works -- Biomedical and Diagnostic Sciences by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Authors

Mohamed A. Abouelkhair, Stephen A. Kania, David A. Bemis, Rebecca E. Rifkin, Remiqiusz M. Grzeskowiak, Alexandru S. Biris, and David A. Anderson



Complete Genome Sequences of Four *Staphylococcus aureus* Sequence Type 398 Isolates from Four Goats with Osteomyelitis

Mohamed A. Abouelkhair,^a Rebecca E. Rifkin,^b Remiqiusz M. Grzeskowiak,^b Alexandru S. Biris,^c David E. Anderson,^b David A. Bemis,^a ⁽ⁱ⁾ Stephen A. Kania^a

Microbiology

Resource Announcements

^aDepartment of Biomedical and Diagnostic Sciences, College of Veterinary Medicine, University of Tennessee, Knoxville, Tennessee, USA

^bDepartment of Large Animal Clinical Sciences, College of Veterinary Medicine, University of Tennessee, Knoxville, Tennessee, USA

^cCenter for Integrative Nanotechnology Sciences, University of Arkansas at Little Rock, Little Rock, Arkansas, USA

ABSTRACT Staphylococcus aureus is the causative agent of multiple infections, including bacteremia, infective endocarditis, osteomyelitis, septic arthritis, and prosthetic device infections. We report here the first whole-genome sequence for four *S. aureus* sequence type 398 isolates from clinical cases of osteomyelitis in four goats with a history of orthopedic surgery.

S nent of the human and animal normal flora (1–3). *S. aureus* is an important opportunistic pathogen in human medicine and is an emerging problem in veterinary practice (1, 2, 4).

Here, we report draft genome sequences of four methicillin-susceptible *S. aureus* (MSSA) sequence type 398 (ST398) isolates obtained from goats that underwent orthopedic surgery at the College of Veterinary Medicine, University of Tennessee (Table 1). The goats were 5 to 6 years old, weighed between 41 and 61 kg, and were free of underlying disease at the time of surgery. The goats had undergone orthopedic surgery a mean of 45 days prior (range, 30 to 90 days) to clinical manifestation of osteomyelitis. Clinical signs included nonhealing wounds, deep surgical site infections, and prolific bone formation. This work was approved by the University of Tennessee IACUC and conducted under protocol 2383-1215.

A single bacterial colony of each strain grown on blood agar plates was inoculated into 5 ml of sterile Trypticase soy broth (TSB) (BD Biosciences, USA; cat. no. RS1-011-21) and incubated overnight at 37°C with shaking at 225 rpm using an Excella E24 incubator shaker (New Brunswick Scientific, USA). DNA was extracted using the MasterPure DNA purification kit (Epicentre, USA; cat. no. MCD85201) according to the manufacturer's instructions.

Sequencing libraries were constructed using the Nextera DNA sample prep kit (Illumina, Inc., USA) according to the manufacturer's instructions. The genomes were sequenced using a MiSeq platform (Illumina, Inc.) with a single-end read length of 150 bp at the University of Tennessee Genomics Core facility. Sequences were trimmed using BBDuk and *de novo* assembled using Geneious version 11.0.3 (5). Annotation was performed by the NCBI Prokaryotic Genome Annotation Pipeline version 4.6 (https:// www.ncbi.nlm.nih.gov/genome/annotation_prok) using the best-placed reference protein set with GenMarkS+. The numbers of reads and contigs, *N*₅₀ values, G+C content values, and the total lengths of the draft genome sequences are listed in (Table 1).

In the genome sequence of S. aureus strain MI18-33, there are 2,793 putative

Received 22 August 2018 Accepted 4 September 2018 Published 27 September 2018 Citation Abouelkhair MA, Rifkin RE,

Grzeskowiak RM, Biris AS, Anderson DE, Bemis DA, Kania SA. 2018. Complete genome sequences of four *Staphylococcus aureus* sequence type 398 isolates from four goats with osteomyelitis. Microbiol Resour Announc 7:e01174-18. https://doi.org/10.1128/MRA .01174-18.

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2018 Abouelkhair et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Stephen A. Kania, skania@utk.edu.

AMERICAN SOCIETY FOR

MICROBIOLOGY

TABLE 1 Strain characteristics

	Data for isolate:			
Characteristic	UTCVM 1	UTCVM 2	UTCVM 3	UTCVM 4
Laboratory accession no.	MI18-33	MI18-34	MI18-935	MI18-1974
GenBank accession no.	QQZR0000000	QQZS0000000	QQZU00000000	QQZX0000000
No. of reads	552,318	760,362	735,277	920,750
No. of contigs	29	28	27	28
N ₅₀ (bp)	193,883	227,498	380,234	228,010
Genome length (bp)	2,835,200	2,834,945	2,835,359	2,836,056
G+C content (%)	32.9	32.9	32.9	32.9
No. of predicted coding sequences	2,793	2,793	2,793	2,790
No. of predicted RNAs	76	75	77	78
SRA accession no.	SRR7765197	SRR7765198	SRR7765195	SRR7765196

protein-coding genes out of 2,905 predicted genes with 112 pseudogenes. Strain MI18-34 contained 2,793 putative protein-coding genes and 109 pseudogenes out of 2,902 predicted genes. Strain MI18-935 had 2,793 putative protein-coding genes out of 2,904 predicted genes with 111 pseudogenes. In strain MI18-1974, there are 2,790 putative protein-coding genes and 111 pseudogenes out of 2,901 predicted genes.

This work used the *S. aureus* MLST database at the University of Oxford (https://pubmlst.org/saureus) for whole-genome multilocus sequence typing (6). All isolates were identified as ST398.

The genome sequences will facilitate identification of the genetic relatedness of these isolates with previously published *S. aureus* strains and may help to elucidate a pathomechansim involved in bone infections, healing, and regeneration (7–9).

Data availability. The whole-genome shotgun sequencing projects of isolates *S. aureus* MI18-33 (UTCVM 1), MI18-34 (UTCVM 2), MI18-935 (UTCVM 3), and MI18-1974 (UTCVM 4) have been deposited at DDBJ/ENA/GenBank and the Sequence Read Archive (SRA) under the accession numbers listed in Table 1.

ACKNOWLEDGMENTS

These studies were partially funded by U.S. Army Medical Research and Materiel Command (USAMRMC) grant W81XWH-15-1-0666.

We acknowledge the Bacteriology Lab, College of Veterinary Medicine, University of Tennessee, for technical assistance.

REFERENCES

- 1. Weese JS. 2010. Methicillin-resistant *Staphylococcus aureus* in animals. ILAR J 51:233–244. https://doi.org/10.1093/ilar.51.3.233.
- Burstiner LC, Faires M, Weese JS. 2010. Methicillin-resistant *Staphylococcus aureus* colonization in personnel attending a veterinary surgery conference. Vet Surg 39:150–157. https://doi.org/10.1111/j.1532-950X.2009.00638.x.
- Weese JS, van Duijkeren E. 2010. Methicillin-resistant Staphylococcus aureus and Staphylococcus pseudintermedius in veterinary medicine. Vet Microbiol 140:418–429. https://doi.org/10.1016/j.vetmic.2009.01.039.
- Kottler S, Middleton JR, Perry J, Weese JS, Cohn LA. 2010. Prevalence of *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* car- riage in three populations. J Vet Intern Med 24:132–139. https://doi.org/ 10.1111/j.1939-1676.2009.0424.x.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28:1647–1649. https://doi.org/10.1093/bioinformatics/bts199.
- Jolley KA, Maiden MCJ. 2010. BIGSdb: scalable analysis of bacterial genome variation at the population level. BMC Bioinformatics 11:595. https://doi.org/10.1186/1471-2105-11-595.

- Baba T, Bae T, Schneewind O, Takeuchi F, Hiramatsu K. 2008. Genome sequence of *Staphylococcus aureus* strain Newman and comparative analysis of staphylococcal genomes: polymorphism and evolution of two major pathogenicity islands. J Bacteriol 190:300–310. https://doi.org/10 .1128/JB.01000-07.
- Holden MT, Feil EJ, Lindsay JA, Peacock SJ, Day NP, Enright MC, Foster TJ, Moore CE, Hurst L, Atkin R, Barron A, Bason N, Bentley SD, Chillingworth C, Chillingworth T, Churcher C, Clark L, Corton C, Cronin A, Doggett J, Dowd L, Feltwell T, Hance Z, Harris B, Hauser H, Holroyd S, Jagels K, James KD, Lennard N, Line A, Mayes R, Moule S, Mungall K, Ormond D, Quail MA, Rabbinowitsch E, Rutherford K, Sanders M, Sharp S, Simmonds M, Stevens K, Whitehead S, Barrell BG, Spratt BG, Parkhill J. 2004. Complete genomes of two clinical *Staphylococcus aureus* strains: evidence for the rapid evolution of virulence and drug resistance. Proc Natl Acad Sci U S A 101:9786–9791. https://doi.org/10.1073/pnas.0402521101.
- Zubair S, Fischer A, Liljander A, Meens J, Hegerman J, Gourle H, Bishop RP, Roebbelen I, Younan M, Mustafa MI, Mushtaq M, Bongcam-Rudloff E, Jores J. 2015. Complete genome sequence of *Staphylococcus aureus*, strain ILRI_Eymole1/1, isolated from a Kenyan dromedary camel. Stand Genomic Sci 10:109. https://doi.org/10.1186/s40793-015-0098-6.