



University of Tennessee, Knoxville Trace: Tennessee Research and Creative Exchange

Middle Atlantic States Mycological Conference
2019

Conferences at UT

4-2019

A community survey of symbiotic mine site fungi along a gradient of recovery on mine sites in the Carolinas

Eudora Miao
Duke University

Rytas Vilgalys
Duke University

Follow this and additional works at: <https://trace.tennessee.edu/masmc>

Recommended Citation

Miao, Eudora and Vilgalys, Rytas, "A community survey of symbiotic mine site fungi along a gradient of recovery on mine sites in the Carolinas" (2019). *Middle Atlantic States Mycological Conference 2019*.
<https://trace.tennessee.edu/masmc/2>

This Poster is brought to you for free and open access by the Conferences at UT at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Middle Atlantic States Mycological Conference 2019 by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Mid-Atlantic States Mycological Conference (MASMC)
University of Tennessee – Knoxville
12-14 April 2019

ABSTRACTS – Posters

A community survey of symbiotic mine site fungi along a gradient of recovery on mine sites in the Carolinas

Eudora Miao, Rytas Vilgalys

Department of Biology, Duke University

In the Anthropocene an increasing part of terrestrial environments is losing ecosystem services and function. This negatively affects both human economics and ecological systems. Phytoremediation, the use of plants to reverse degradation and to restore ecological function, has been a promising approach. However, the symbiotic soil microbiota that influence the effectiveness of this method is not fully understood. I sampled the soil and roots of *Pinus* spp. (pines) at four sites along a gradient of vegetation recovery on the Superfund Site Brewer Gold Mine (SC), the Henry Knob Mine (SC), and Russell Gold Mine. Acidity, nutrient profile, and heavy-metal contamination of collected soil were determined. DNA was extracted from soil and root samples followed by preparation of multiplex PCR samples of the ITS region. Sequence reads generated through Illumina Miseq were processed through the QIIME pipeline. Taxonomy was assigned through the UNITE database. Results showed a pattern of succession in fungal communities along a recovery gradient. While mycorrhizal fungi on the least recovered site were dominated by *Acephala* and *Pisolithus*, sites with more recovered vegetation revealed a more diverse array of symbiotic fungi, including *Rhizopogon*, *Tomentella*, *Tricholoma*, and *Laccaria*. These fungi, although later in the succession pattern, likely add more diverse benefits to help their plant associates cope with the stressful environment. This “bioprospecting” method could be applied to extract and amplify symbiotic fungi to facilitate re-vegetation efforts.