



University of Tennessee, Knoxville
Trace: Tennessee Research and Creative Exchange

Middle Atlantic States Mycological Conference
2019

Conferences at UT

4-2019

Detecting polyploidy and its role in diversification across *Hydnum* (Cantharellales)

Rachel Swenie

University of Tennessee, Knoxville

P. Brandon Matheny

University of Tennessee, Knoxville

Follow this and additional works at: <https://trace.tennessee.edu/masmc>

Recommended Citation

Swenie, Rachel and Matheny, P. Brandon, "Detecting polyploidy and its role in diversification across *Hydnum* (Cantharellales)" (2019). *Middle Atlantic States Mycological Conference 2019*.
<https://trace.tennessee.edu/masmc/23>

This Presentation is brought to you for free and open access by the Conferences at UT at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Middle Atlantic States Mycological Conference 2019 by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Mid-Atlantic States Mycological Conference (MASMC)
University of Tennessee – Knoxville
12-14 April 2019

ABSTRACTS - Oral Presentations

Detecting polyploidy and its role in diversification across *Hydnum* (Cantharellales)

Rachel Swenie, P. Brandon Matheny

Department of Ecology and Evolutionary Biology, University of Tennessee

Whole genome duplication and gene-loss events leading to changes in ploidy levels are well-characterized in plants and animals. However, the frequency of ploidy changes and subsequent effects on diversification are poorly understood in mushroom-forming fungi. Previous research on the genus *Hydnum* (Cantharellales) has suggested that some species vary in ploidy level and differ by as much as a four-fold increase in chromosome sets. If so, ploidy level could correlate with changes in morphology observed across the phylogeny of *Hydnum*. Here, we present a framework to estimate ploidy levels in *Hydnum* and discuss hypotheses regarding the effect of polyploidy on morphology and diversification in the genus. To estimate ploidy levels in *Hydnum*, we collected dense basidiospore deposits from a range of species to compare relative gene content using flow cytometry. To study the phylogenetic distribution of ploidy level and other characters, we constructed a three-gene (ITS, *rbp2*, *tef1*) phylogeny from more than 50 samples of *Hydnum*, which provides a robust evolutionary framework for comparative analysis.