

## APPENDICES

## Appendix A

GenBank Entry for *Escherichia coli* partial PMI mRNA

---

LOCUS ECOMANAA 1604 bp DNA linear BCT  
26-APR-1993

DEFINITION *E.coli* manA gene encoding phosphomannose isomerase, complete cds.

ACCESSION M15380

VERSION M15380.1

KEYWORDS isomerase; mannose; phosphomannose isomerase.

SOURCE *Escherichia coli*

ORGANISM [Escherichia coli](#)  
Bacteria; Proteobacteria; Gammaproteobacteria; Enterobacteriales; Enterobacteriaceae; *Escherichia*.

REFERENCE 1 (bases 1 to 1604)

AUTHORS Miles, J.S. and Guest, J.R.

TITLE Nucleotide sequence and transcriptional start point of the phosphomannose isomerase gene (manA) of *Escherichia coli*

JOURNAL *Gene* 32 (1-2), 41-48 (1984)

PUBMED 6397402

COMMENT Original source text: *E.coli* (strain GM242) DNA, clone pGS57.

FEATURES Location/Qualifiers

source	1..1604 /organism=" <a href="#">Escherichia coli</a> " /mol_type="genomic DNA" /db_xref=" <a href="#">taxon:562</a> "
mRNA	1..1604 /product="PMI mRNA"
gene	392..1567 /gene="manA"
CDS	392..1567 /gene="manA" /note="phosphomannose isomerase" /codon_start=1 /transl_table=11 /protein_id=" <a href="#">AAA24109.1</a> " /db_xref="GI:146722"

/translation="MQKLINSVQNYAWGSKTALTELYGMENPSSQPMAELWMGAHPKS  
SSRVQNAAGDIVSLRDVIESDKSTLLGEAVAKRFGELPFLFKVLCQAQPLSIQVHPNK  
HNSEIGFAKENAAGIPMDAAERNYKDPNHKPELVFALTPFLAMNAFREFSEIVSLLQP  
VAGAHPAIAHF<sup>1</sup>LQQPDAERLSELFASLLNMQGEEKSRALAILKSALDSQQGEPWQTIR  
LISEFYPEDSGLFSPLLLNVVKLNPGEAMFLFAETPHAYLQGVALEVMANS<sup>409</sup>DNVLRAG  
LTPKYIDIPELVANVKFEAKPANQLLTQPVKQGAELDFPIPVDDFAFSLHDLSDKETT  
ISQQAAILFCVEGDATLWKGSQQQLKPGESAFIAANESPVTVKGGHRLARVYNKL"  
BASE COUNT 409 a 391 c 392 g 412 t


ORIGIN 1 bp upstream of MspI site.

```

1 ccggacgcag catgaacgac gcatcatgaa acgcctggcg cgccaacaga
gttaacgctt
61 cgggtgcgac tttcaaaatc tcctgccctt caaattcaga tacgctaacg
tgttcgctgg
121 ttagcaggta atactcagta tcactctttt tgagtggaaa aggagcctga
taatgaaagg
181 gtttgtttga cattgttctc tcacttactg cctggtttgg ttatgctctg
ggcgggtggt
241 ccgttgccct gttaaaagcg agtaacaata tcctacacac tttttaaca
aaaactgaga
301 ctagtacgac tttttgcggc tccaggttac ttcccgtagg attcctgctt
taatagtggg
361 attaatctcc acattaaaac agggattgat catgcaaaaa ctcattaact
cagtgcaaaa
421 ctatgcctgg ggcagcaaaa cggcgttgac tgaactttat ggtatggaaa
atccgtccag
481 ccagccgatg gccgagctgt ggatggggcg acatccgaaa agcagttcac
gagtgcagaa
541 tgccgccgga gatatcgttt cactgcgtga tgtgattgag agtgataaat
cgactctgct
601 cggagaggcc gttgccaaac gctttggcga actgcctttc ctgttcaaag
tattatgcgc
661 agcacagcca ctctccattc aggttcatcc aaacaacac aattctgaaa
tcggttttgc
721 caaagaaaat gccgcaggta tcccgatgga tgccgccgag cgtaactata
aagatcctaa
781 ccacaagccg gagctggttt ttgcgctgac gcctttcctt gcgatgaacg
cgtttcgtga
841 attttccgag attgtctccc tactccagcc ggtcgcagggt gcacatccgg
cgattgctca
901 ctttttacia cagcctgatg ccgaacgttt aagcgaactg ttcgccagcc
tgttgaatat
961 gcagggtgaa gaaaaatccc gcgcgctggc gattttaaaa tcggcctcg
atagccagca
1021 gggtgaaacc tggcaaacga ttcgtttaat ttctgaattt taccgggaag
acagcggctc
1081 gttctccccg ctattgctga atgtggtgaa attgaaccct ggcgaagcga
tgttcctggt
1141 cgctgaaaca ccgcacgctt acctgcaagg cgtggcgctg gaagtgatgg
caaactccga
1201 taacgtgctg cgtgcgggtc tgacgcctaa atacattgat attccggaac
tggttgccaa
1261 tgtgaaatcc gaagccaaac cggctaacca gttggtgacc cagccggtga
aacaaggtgc
1321 agaactggac ttcccgatc cagtggatga tttgccttc tcgctgcatg
accttagtga
1381 taaagaaacc accattagcc agcagagtgc cgccattttg ttctgcgtcg
aaggcgatgc
1441 aacgttgagg aaaggttctc agcagttaca gcttaaaccg ggtgaatcag
cgtttattgc
1501 cgccaacgaa tcaccggtga ctgtcaaagg ccacggccgt ttagcgcgtg
tttacaacaa
1561 gctgtaagag cttactgaaa aaattaacat ctcttgctaa gctt


```

## Appendix B




Restriction map of pMI3

## Appendix C




Restriction map of pMI11

## Appendix D



Restriction map of pMI3G


## Appendix E



Restriction map of pMI11G

## Appendix F

## Fructose and mannose metabolism



### Appendix G

Effect of mannose on selection and regeneration of oil palm embryogenic calli. EC were cultured for 5 months on different concentrations selection medium and 30 g/l sucrose used as a control.

No.	Man:Suc (g/l)	Average fresh weight (g) in 5 months						% proliferation
		0	1	2	3	4	5	
1	0:30	0.54	2.48 ± 0.08	13.04 ± 0.50	47.98 ± 0.16	100.26 ± 0.58	154.45 ± 0.29	100
2	5:25	0.56	2.21 ± 0.08	11.09 ± 0.18	31.39 ± 0.41	77.60 ± 0.47	130.51 ± 0.26	84.43
3	10:20	0.54	2.25 ± 0.15	11.71 ± 0.35	49.80 ± 0.28	84.75 ± 0.28	101.13 ± 0.32	65.36
4	15:15	0.57	2.08 ± 0.24	11.34 ± 0.49	44.81 ± 1.52	75.43 ± 1.71	126.03 ± 3.70	81.52
5	20:10	0.56	2.82 ± 0.18	8.10 ± 0.23	26.81 ± 0.49	66.71 ± 0.97	111.48 ± 4.00	72.07
6	25:5	0.56	2.58 ± 0.16	11.08 ± 0.41	49.35 ± 1.08	79.31 ± 0.50	107.12 ± 2.90	69.24
7	30:0	0.53	2.38 ± 0.09	12.42 ± 1.19	43.35 ± 1.83	72.26 ± 2.10	98.46 ± 0.97	63.63

Values represent the mean ± S.E. of five replications.



**LIST OF PUBLICATIONS**

## NATIONAL PROCEEDINGS

1. Bahariah, B., Parveez, G. K. A., Norzulaani, K and Rofina, Y. O. (2009). The use of a non-antibiotic selection system, phosphomannose isomerase in oil palm transformation. In 34<sup>th</sup> Annual Conference of the MSBMB held in Kuala Lumpur, 7-8 October 2009, pp 82.
  
2. Bahariah, B., Parveez, G. K. A., Norzulaani, K and Rofina, Y. O. (2009). Phosphomannose isomerase as an alternative selectable marker gene for oil palm transformation. In 2009 PIPOC International Palm Oil Congress, Kuala Lumpur, 9-15 November 2009, pp 1254-1266.