

## REFERENCES

- Abdullah ,N, A, B (2008) : Development of recombinant *Mycobacterium bovis* Bacille calmette guerin(rBCG) expressing the 19kDa-c terminus of merozoite surface protein-1(*MSP-1C*) and the 22kDa of serine repeat antigen (SE22) of *plasmodium falciparum* as a potential blood-stage malaria vaccine , PHD thesis .
- Aikawa, M., Iseki, M., Barnwell, J. W., Taylor, D., Oo, M. M., & Howard, R. J. (1990). The pathology of human cerebral malaria. *Am J Trop Med Hyg*, 43(2 Pt 2), 30-37.
- Al Yaman, F. M., Mokela, D., Genton, B., Rockett, K. A., Alpers, M. P., & Clark, I. A. (1996). Association between serum levels of reactive nitrogen intermediates and coma in children with cerebral malaria in Papua New Guinea. *Trans R Soc Trop Med Hyg*, 90(3), 270-273.
- Al-Yaman, F., Genton, B., Reeder, J. C., Mokela, D., Anders, R. F., & Alpers, M. P. (1997). Humoral response to defined *Plasmodium falciparum* antigens in cerebral and uncomplicated malaria and their relationship to parasite genotype. *Am J Trop Med Hyg*, 56(4), 430-435.
- Anderson, T. J., Haubold, B., Williams, J. T., Estrada-Franco, J. G., Richardson, L., Mollinedo, R., et al. (2000). Microsatellite markers reveal a spectrum of population structures in the malaria parasite *Plasmodium falciparum*. *Mol Biol Evol*, 17(10), 1467-1482.
- Arez, A. P., Snounou, G., Pinto, J., Sousa, C. A., Modiano, D., Ribeiro, H., et al. (1999). A clonal *Plasmodium falciparum* population in an isolated outbreak of malaria in the Republic of Cabo Verde. *Parasitology*, 118 (Pt 4), 347-355.

- Ariey, F., Chalvet, W., Hommel, D., Peneau, C., Hulin, A., Mercereau-Puijalon, O., et al. (1999). *Plasmodium falciparum* parasites in French Guiana: limited genetic diversity and high selfing rate. *Am J Trop Med Hyg*, 61(6), 978-985.
- Ariey, F., Hommel, D., Le Scanf, C., Duchemin, J. B., Peneau, C., Hulin, A., et al. (2001). Association of severe malaria with a specific *Plasmodium falciparum* genotype in French Guiana. *J Infect Dis*, 184(2), 237-241.
- Arnot, D. (1998). Unstable malaria in Sudan: the influence of the dry season. Clone multiplicity of *Plasmodium falciparum* infections in individuals exposed to variable levels of disease transmission. *Trans R Soc Trop Med Hyg*, 92(6), 580-585.
- Aubouy, A., Migot-Nabias, F., & Deloron, P. (2003). Polymorphism in two merozoite surface proteins of *Plasmodium falciparum* isolates from Gabon. *Malar J*, 2, 12.
- Babiker, H. A., Charlwood, J. D., Smith, T., & Walliker, D. (1995). Gene flow and cross-mating in *Plasmodium falciparum* in households in a Tanzanian village. *Parasitology*, 433-442.
- Babiker, H. A., Lines, J., Hill, W. G., & Walliker, D. (1997). Population structure of *Plasmodium falciparum* in villages with different malaria endemicity in east Africa. *Am J Trop Med Hyg*, 56(2), 141-147.
- Babiker, H. A., Ranford-Cartwright, L. C., & Walliker, D. (1999). Genetic structure and dynamics of *Plasmodium falciparum* infections in the Kilombero region of Tanzania. *Trans R Soc Trop Med Hyg*, 93 Suppl 1, 11-14.

- Babiker, H. A., Satti, G., & Walliker, D. (1995). Genetic changes in the population of *Plasmodium falciparum* in a Sudanese village over a three-year period. *Am J Trop Med Hyg*, 53(1), 7-15.
- Basco, L. K., Tahar, R., & Escalante, A. (2004). Molecular epidemiology of malaria in Cameroon. XVIII. Polymorphisms of the *Plasmodium falciparum* merozoite surface antigen-2 gene in isolates from symptomatic patients. *Am J Trop Med Hyg*, 70(3), 238-244.
- Beck, H. P., Felger, I., Huber, W., Steiger, S., Smith, T., Weiss, N., et al. (1997). Analysis of multiple *Plasmodium falciparum* infections in Tanzanian children during the phase III trial of the malaria vaccine SPf66. *J Infect Dis*, 175(4), 921-926.
- Bendixen, M., Msangeni, H. A., Pedersen, B. V., Shayo, D., & Bodker, R. (2001). Diversity of *Plasmodium falciparum* populations and complexity of infections in relation to transmission intensity and host age: a study from the Usambara Mountains, Tanzania. *Trans R Soc Trop Med Hyg*, 95(2), 143-148.
- Black, C. G., Wu, T., Wang, L., Hibbs, A. R., & Coppel, R. L. (2001). Merozoite surface protein 8 of *Plasmodium falciparum* contains two epidermal growth factor-like domains. *Mol Biochem Parasitol*, 114(2), 217-226.
- Blackman, M. J., & Holder, A. A. (1992). Secondary processing of the *Plasmodium falciparum* merozoite surface protein-1 (*MSP-1*) by a calcium-dependent membrane-bound serine protease: shedding of *MSP-133* as a noncovalently associated complex with other fragments of the *MSP-1*. *Mol Biochem Parasitol*, 50(2), 307-315.

- Bogreau, H., Renaud, F., Bouchiba, H., Durand, P., Assi, S. B., Henry, M. C., et al. (2006). Genetic diversity and structure of African *Plasmodium falciparum* populations in urban and rural areas. *Am J Trop Med Hyg*, 74(6), 953-959.
- Bruce, M. C., Galinski, M. R., Barnwell, J. W., Donnelly, C. A., Walmsley, M., Alpers, M. P., et al. (2000). Genetic diversity and dynamics of *plasmodium falciparum* and *P. vivax* populations in multiply infected children with asymptomatic malaria infections in Papua New Guinea. *Parasitology*, 121 (Pt 3), 257-272.
- Cattamanchi, A., Kyabayinze, D., Hubbard, A., Rosenthal, P. J., & Dorsey, G. (2003). Distinguishing recrudescence from reinfection in a longitudinal antimalarial drug efficacy study: comparison of results based on genotyping of *MSP-1*, *MSP-2*, and glurp. *Am J Trop Med Hyg*, 68(2), 133-139.
- Chappel, J. A., & Holder, A. A. (1993). Monoclonal antibodies that inhibit *Plasmodium falciparum* invasion in vitro recognise the first growth factor-like domain of merozoite surface protein-1. *Mol Biochem Parasitol*, 60(2), 303-311.
- Chenet, S. M., Branch, O. H., Escalante, A. A., Lucas, C. M., & Bacon, D. J. (2008). Genetic diversity of vaccine candidate antigens in *Plasmodium falciparum* isolates from the Amazon basin of Peru. *Malar J*, 7, 93.
- Clark, J. T., Donachie, S., Anand, R., Wilson, C. F., Heidrich, H. G., & McBride, J. S. (1989). 46-53 kilodalton glycoprotein from the surface of *Plasmodium falciparum* merozoites. *Mol Biochem Parasitol*, 32(1), 15-24.
- Conway, D. J., Greenwood, B. M., & McBride, J. S. (1991). The epidemiology of multiple-clone *Plasmodium falciparum* infections in Gambian patients. *Parasitology*, 103 Pt 1, 1-6.

- Coppel, R. L., Cowman, A. F., Lingelbach, K. R., Brown, G. V., Saint, R. B., Kemp, D. J., et al. (1983). Isolate-specific S-antigen of *Plasmodium falciparum* contains a repeated sequence of eleven amino acids. *Nature*, 306(5945), 751-756.
- Daubersies, P., Sallenave-Sales, S., Magne, S., Trape, J. F., Contamin, H., Fandeur, T., et al. (1996). Rapid turnover of *Plasmodium falciparum* populations in asymptomatic individuals living in a high transmission area. *Am J Trop Med Hyg*, 54(1), 18-26.
- Davis, T. M., Supanaranond, W., Pukrittayakamee, S., Silamut, K., & White, N. J. (2003). Evidence for continued two-brood replication of *Plasmodium falciparum* in vivo during quinine treatment. *Acta Trop*, 89(1), 41-45.
- David, P. H., Hadley, T. J., Aikawa, M., & Miller, L. H. (1984). Processing of a major parasite surface glycoprotein during the ultimate stages of differentiation in *Plasmodium knowlesi*. *Mol Biochem Parasitol*, 11, 267-282.
- Decuyper, S., Elinck, E., Van Overmeir, C., Talisuna, A. O., D'Alessandro, U., & Dujardin, J. C. (2003). Pathogen genotyping in polyclonal infections: application of a fluorogenic polymerase-chain-reaction assay in malaria. *J Infect Dis*, 188(8), 1245-1249.
- Deleersnijder, W., Hendrix, D., Bendahman, N., Hanegreefs, J., Brijs, L., Hamers-Casterman, C., et al. (1990). Molecular cloning and sequence analysis of the gene encoding the major merozoite surface antigen of *Plasmodium chabaudi* IP-PC1. *Mol Biochem Parasitol*, 43(2), 231-244.
- Dent, A. E., Yohn, C. T., Zimmerman, P. A., Vulule, J., Kazura, J. W., & Moormann, A. M. (2007). A polymerase chain reaction/ligase detection reaction fluorescent

- microsphere assay to determine *Plasmodium falciparum* MSP-119 haplotypes. *Am J Trop Med Hyg*, 77(2), 250-255.
- Eliades, M. J., Wolkon, A., Morgah, K., Crawford, S. B., Dorkenoo, A., Sodahlon, Y., et al. (2006). Burden of malaria at community level in children less than 5 years of age in Togo. *Am J Trop Med Hyg*, 75(4), 622-629.
- Engelbrecht, F., Togel, E., Beck, H. P., Enwezor, F., Oettli, A., & Felger, I. (2000). Analysis of *Plasmodium falciparum* infections in a village community in Northern Nigeria: determination of MSP-2 genotypes and parasite-specific IgG responses. *Acta Trop*, 74(1), 63-71.
- Epstein, N., Miller, L. H., Kaushel, D. C., Udeinya, I. J., Rener, J., Howard, R. J., et al. (1981). Monoclonal antibodies against a specific surface determinant on malarial (*Plasmodium knowlesi*) merozoites block erythrocyte invasion. *J Immunol*, 127(1), 212-217.
- Farnert, A., Arez, A. P., Babiker, H. A., Beck, H. P., Benito, A., Bjorkman, A., et al. (2001). Genotyping of *Plasmodium falciparum* infections by PCR: a comparative multicentre study. *Trans R Soc Trop Med Hyg*, 95(2), 225-232.
- Farnert, A., Rooth, I., Svensson, Snounou, G., & Bjorkman, A. (1999). Complexity of *Plasmodium falciparum* infections is consistent over time and protects against clinical disease in Tanzanian children. *J Infect Dis*, 179(4), 989-995.
- Farnert, A., Snounou, G., Rooth, I., & Bjorkman, A. (1997). Daily dynamics of *Plasmodium falciparum* subpopulations in asymptomatic children in a holoendemic area. *Am J Trop Med Hyg*, 56(5), 538-547.

- Felger, I., Marshal, V. M., Reeder, J. C., Hunt, J. A., Mgone, C. S., & Beck, H. P. (1997). Sequence diversity and molecular evolution of the merozoite surface antigen 2 of *Plasmodium falciparum*. *J Mol Evol*, 45(2), 154-160.
- Felger, I., Smith, T., Edoh, D., Kitua, A., Alonso, P., Tanner, M., et al. (1999). Multiple *Plasmodium falciparum* infections in Tanzanian infants. *Trans R Soc Trop Med Hyg*, 93 Suppl 1, 29-34.
- Felger, I., Tavul, L., Kabintik, S., Marshall, V., Genton, B., Alpers, M., et al. (1994). *Plasmodium falciparum*: extensive polymorphism in merozoite surface antigen 2 alleles in an area with endemic malaria in Papua New Guinea. *Exp Parasitol*, 79(2), 106-116.
- Fenton, B., Clark, J. T., Wilson, C. F., McBride, J. S., & Walliker, D. (1989). Polymorphism of a 35-48 kDa *Plasmodium falciparum* merozoite surface antigen. *Mol Biochem Parasitol*, 34(1), 79-86.
- Garcia, J. E., Puentes, A., & Patarroyo, M. E. (2006). Developmental biology of sporozoite-host interactions in *Plasmodium falciparum* malaria: implications for vaccine design. *Clin Microbiol Rev*, 19(4), 686-707.
- Gardner, M. J., Hall, N., Fung, E., White, O., Berriman, M., Hyman, R. W., et al. (2002). Genome sequence of the human malaria parasite *Plasmodium falciparum*. *Nature*, 419(6906), 498-511.
- Genton, B., Betuela, I., Felger, I., Al-Yaman, F., Anders, R. F., Saul, A., et al. (2002). A recombinant blood-stage malaria vaccine reduces *Plasmodium falciparum* density and exerts selective pressure on parasite populations in a phase 1-2b trial in Papua New Guinea. *J Infect Dis*, 185(6), 820-827.

- Gerold, P., Schofield, L., Blackman, M. J., Holder, A. A., & Schwarz, R. T. (1996). Structural analysis of the glycosyl-phosphatidylinositol membrane anchor of the merozoite surface proteins-1 and -2 of *Plasmodium falciparum*. *Mol Biochem Parasitol*, 75(2), 131-143.
- Ghanchi, N. K., Martensson, A., Ursing, J., Jafri, S., Bereczky, S., Hussain, R. (2010). Genetic diversity among *Plasmodium falciparum* field isolates in Pakistan measured with PCR genotyping of the merozoite surface protein 1 and 2. *Malar J*, 9, 1.
- Gibson, H. L., Tucker, J. E., Kaslow, D. C., Krettli, A. U., Collins, W. E., Kiefer, M. C., et al. (1992). Structure and expression of the gene for Pv200, a major blood-stage surface antigen of *Plasmodium vivax*. *Mol Biochem Parasitol*, 50(2), 325-333.
- Gilles hm, Warrell, da (1993) Bruce-Chwatt's essential malariology. 3rd ed. Edward Arnold, Hodder and Stoughton, London.
- Gomez, D., Chaparro, J., Rubiano, C., Rojas, M. O., & Wasserman, M. (2002). Genetic diversity of *Plasmodium falciparum* field samples from an isolated Colombian village. *Am J Trop Med Hyg*, 67(6), 611-616.
- Guinovart, C., Navia, M. M., Tanner, M., & Alonso, P. L. (2006). Malaria: burden of disease. *Curr Mol Med*, 6(2), 137-140.
- Haddad, D., Snounou, G., Mattei, D., Enamorado, I. G., Figueroa, J., Stahl, S., et al. (1999). Limited genetic diversity of *Plasmodium falciparum* in field isolates from Honduras. *Am J Trop Med Hyg*, 60(1), 30-34.
- Happi, C. T., Gbotosho, G. O., Sowunmi, A., Falade, C. O., Akinboye, D. O., Gerena, L., et al. (2004). Molecular analysis of *Plasmodium falciparum* recrudescence malaria

- infections in children treated with chloroquine in Nigeria. *Am J Trop Med Hyg*, 70(1), 20-26.
- Healer, J., Murphy, V., Hodder, A. N., Masciantonio, R., Gemmill, A. W., Anders, R. F., et al. (2004). Allelic polymorphisms in apical membrane antigen-1 are responsible for evasion of antibody-mediated inhibition in *Plasmodium falciparum*. *Mol Microbiol*, 52(1), 159-168.
- Heidari, A., Keshavarz, H., Rokni, M. B., & Jelinek, T. (2007). Genetic diversity in merozoite surface protein (MSP)-1 and MSP-2 genes of *Plasmodium falciparum* in a major endemic region of Iran. *Korean J Parasitol*, 45(1), 59-63.
- Herrera, S., Rudin, W., Herrera, M., Clavijo, P., Mancilla, L., de Plata, C., et al. (1993). A conserved region of the MSP-1 surface protein of *Plasmodium falciparum* contains a recognition sequence for erythrocyte spectrin. *EMBO J*, 12(4), 1607-1614.
- Holder, A. A., & Freeman, R. R. (1984). The three major antigens on the surface of *Plasmodium falciparum* merozoites are derived from a single high molecular weight precursor. *J Exp Med*, 160(2), 624-629.
- Holder, A. A., & Blackman, M. J. (1994). What is the function of MSP-I on the malaria merozoite? *Parasitol Today*, 10(5), 182-184.
- Irion, A., Beck, H. P., & Felger, I. (1997). New repeat unit and hot spot of recombination in FC27-type alleles of the gene coding for *Plasmodium falciparum* merozoite surface protein 2. *Mol Biochem Parasitol*, 90(1), 367-370.

Irion, A. (2000). Molecular diversity and immunological properties of the *Plasmodium falciparum* Merozoite Surface Protein 2 (*MSP-2*). PhD Thesis ,University of Basel, Switzerland.

Issifou, S., Djikou, S., Sanni, A., Lekoulou, F., & Ntoumi, F. (2001). No influence of season of transmission nor age of patients on the complexity and genetic diversity of *Plasmodium falciparum* infection in Cotonou, Benin. *Bull Soc Pathol Exot*, 94(2 Pt 2), 195-198.

Joshi, H., Valecha, N., Verma, A., Kaul, A., Mallick, P. k.Shalini, S. et al. (2007). Genetic structure of *Plasmodium falciparum* field isolates in eastern and north-eastern India. *Malar J*, 6, 60.

Kang, J. M., Moon, S. U., Kim, J. Y., Cho, S. H., Lin, K., Sohn, W. M., (2010). Genetic polymorphism of merozoite surface protein-1 and merozoite surface protein-2 in *Plasmodium falciparum* field isolates from Myanmar. *Malar J*, 9, 131.

Khatoon, L., Baliraine, F. N., Bonizzoni, M., Malik, S. A., & Yan, G (2010). Genetic structure of *Plasmodium vivax* and *Plasmodium falciparum* in the Bannu district of Pakistan. *Malar J*, 9, 112.

Konate, L., Zwetyenga, J., Rogier, C., Bischoff, E., Fontenille, D. Tall, A. (1999). Variation of *Plasmodium falciparum* *MSP-1* block 2 and *MSP-2* allele prevalence and of infection complexity in two neighbouring Senegalese villages with different transmission conditions. *Trans R Soc Trop Med Hyg*, 93 Suppl 1, 21-28.

Krajden, S., Panisko, D. M., Tobe, B., Yang, J., & Keystone, J. S. (1991). Prolonged infection with *Plasmodium falciparum* in a semi-immune patient. *Trans R Soc Trop Med Hyg*, 85(6), 731-732.

- Kumar, S., Epstein, J. E., & Richie, T. L. (2002). Vaccines against asexual stage malaria parasites. *Chem Immunol*, 80, 262-286.
- Kushwaha, A., Rao, P. P., Suresh, R. P., & Chauhan, V. S. (2001). Immunogenicity of recombinant fragments of *Plasmodium falciparum* acidic basic repeat antigen produced in Escherichia coli. *Parasite Immunol*, 23(8), 435-444.
- Kyes, S., Harding, R., Black, G., Craig, A., Peshu, N., Newbold, C., et al. (1997). Limited spatial clustering of individual *Plasmodium falciparum* alleles in field isolates from coastal Kenya. *Am J Trop Med Hyg*, 57(2), 205-215.
- Lawrence, N., Stowers, A., Mann, V., Taylor, D., & Saul, A. (2000). Recombinant chimeric proteins generated from conserved regions of *Plasmodium falciparum* merozoite surface protein 2 generate antiparasite humoral responses in mice. *Parasite Immunol*, 22(5), 211-221.
- Lengeler, C. (2004). Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Syst Rev*(2), CD000363.
- Lewis, A. P. (1989). Cloning and analysis of the gene encoding the 230-kilodalton merozoite surface antigen of *Plasmodium yoelii*. *Mol Biochem Parasitol*, 36(3), 271-282.
- Liljander, A. (2010). *Malaria : multiclonal infections and protective immunity*. Karolinska Institutet, Stockholm, Sweden.
- MacLeod, A. (2004). Minisatellites and MVR-PCR for the individual identification of parasite isolates. *Methods Mol Biol*, 270, 187-202.
- Marks, F., Meyer, C. G., Sievertsen, J., Timmann, C., Evans, J., Horstmann, R. D., et al. (2004). Genotyping of *Plasmodium falciparum* pyrimethamine resistance by

- matrix-assisted laser desorption-ionization time-of-flight mass spectrometry. *Antimicrob Agents Chemother*, 48(2), 466-472.
- Marshall, V. M., Anthony, R. L., Bangs, M. J., Purnomo, Anders, R. F., & Coppel, R. L. (1994). Allelic variants of the *Plasmodium falciparum* merozoite surface antigen 2 (MSA-2) in a geographically restricted area of Irian Jaya. *Mol Biochem Parasitol*, 63(1), 13-21.
- Marshall, V. M., Coppel, R. L., Anders, R. F., & Kemp, D. J. (1992). Two novel alleles within subfamilies of the merozoite surface antigen 2 (MSA-2) of *Plasmodium falciparum*. *Mol Biochem Parasitol*, 50(1), 181-184.
- Marshall, V. M., Silva, A., Foley, M., Cranmer, S., Wang, L., McColl, D. J., et al. (1997). A second merozoite surface protein (*MSP-4*) of *Plasmodium falciparum* that contains an epidermal growth factor-like domain. *Infect Immun*, 65(11), 4460-4467.
- Marshall, V. M., Tieqiao, W., & Coppel, R. L. (1998). Close linkage of three merozoite surface protein genes on chromosome 2 of *Plasmodium falciparum*. *Mol Biochem Parasitol*, 94(1), 13-25.
- Mayor, A., Saute, F., Aponte, J. J., Almeda, J., Gomez-Olive, F. X., Dgedge, M., et al. (2003). *Plasmodium falciparum* multiple infections in Mozambique, its relation to other malariological indices and to prospective risk of malaria morbidity. *Trop Med Int Health*, 8(1), 3-11.
- McColl, D. J., Silva, A., Foley, M., Kun, J. F., Favaloro, J. M., Thompson, J. K., et al. (1994). Molecular variation in a novel polymorphic antigen associated with *Plasmodium falciparum* merozoites. *Mol Biochem Parasitol*, 68(1), 53-67.

- McKean, P. G., O'Dea, K., & Brown, K. N. (1993). Nucleotide sequence analysis and epitope mapping of the merozoite surface protein 1 from *Plasmodium chabaudi chabaudi* AS. *Mol Biochem Parasitol*, 62(2), 199-209.
- Menegon, M., Severini, C., Sannella, A., Paglia, M. G., Sangare, D., Abdel-Wahab, A., et al. (2000). Genotyping of *Plasmodium falciparum* gametocytes by reverse transcriptase polymerase chain reaction. *Mol Biochem Parasitol*, 111(1), 153-161.
- Miller, L. H., Roberts, T., Shahabuddin, M., & McCutchan, T. F. (1993). Analysis of sequence diversity in the *Plasmodium falciparum* merozoite surface protein-1 (MSP-1). *Mol Biochem Parasitol*, 59(1), 1-14.
- Montoya, L., Maestre, A., Carmona, J., Lopes, D., Do Rosario, V., & Blair, S. (2003). *Plasmodium falciparum*: diversity studies of isolates from two Colombian regions with different endemicity. *Exp Parasitol*, 104(1-2), 14-19.
- Mugittu, K., Adjuik, M., Snounou, G., Ntoumi, F., Taylor, W., Mshinda, H., et al. (2006). Molecular genotyping to distinguish between recrudescents and new infections in treatment trials of *Plasmodium falciparum* malaria conducted in Sub-Saharan Africa: adjustment of parasitological outcomes and assessment of genotyping effectiveness. *Trop Med Int Health*, 11(9), 1350-1359.
- Muller, O., Traore, C., Becher, H., & Kouyate, B. (2003). Malaria morbidity, treatment-seeking behaviour, and mortality in a cohort of young children in rural Burkina Faso. *Trop Med Int Health*, 8(4), 290-296.
- Nwagwu, M., Haynes, J. D., Orlandi, P. A., & Chulay, J. D. (1992). *Plasmodium falciparum*: chymotryptic-like proteolysis associated with a 101-kDa acidic-basic repeat antigen. *Exp Parasitol*, 75(4), 399-414.

- O'Dempsey, T. J., McArdle, T. F., Laurence, B. E., Lamont, A. C., Todd, J. E., & Greenwood, B. M. (1993). Overlap in the clinical features of pneumonia and malaria in African children. *Trans R Soc Trop Med Hyg*, 87(6), 662-665.
- Oeuvray, C., Bouharoun-Tayoun, H., Gras-Masse, H., Bottius, E., Kaidoh, T., Aikawa, M., et al. (1994). Merozoite surface protein-3: a malaria protein inducing antibodies that promote *Plasmodium falciparum* killing by cooperation with blood monocytes. *Blood*, 84(5), 1594-1602.
- Pachebat, J. A., Ling, I. T., Grainger, M., Trucco, C., Howell, S., Fernandez-Reyes, D., et al. (2001). The 22 kDa component of the protein complex on the surface of *Plasmodium falciparum* merozoites is derived from a larger precursor, merozoite surface protein 7. *Mol Biochem Parasitol*, 117(1), 83-89.
- Paul, R. E., & Day, K. P. (1998). Mating patterns of *Plasmodium falciparum*. *Parasitol Today*, 14(5), 197-202.
- Paul, R. E., Hackford, I., Brockman, A., Muller-Graf, C., Price, R., Luxemburger, C., et al. (1998). Transmission intensity and *Plasmodium falciparum* diversity on the northwestern border of Thailand. *Am J Trop Med Hyg*, 58(2), 195-203.
- Paul, R. E., Packer, M. J., Walmsley, M., Lagog, M., Ranford-Cartwright, L. C., Paru, R., et al. (1995). Mating patterns in malaria parasite populations of Papua New Guinea. *Science*, 269(5231), 1709-1711.
- Pearce, J. A., Mills, K., Triglia, T., Cowman, A. F., & Anders, R. F. (2005). Characterisation of two novel proteins from the asexual stage of *Plasmodium falciparum*, H101 and H103. *Mol Biochem Parasitol*, 139(2), 141-151.

- Peyerl-Hoffmann, G., Jelinek, T., Kilian, A., Kabagambe, G., Metzger, W. G., & von Sonnenburg, F. (2001). Genetic diversity of *Plasmodium falciparum* and its relationship to parasite density in an area with different malaria endemicities in West Uganda. *Trop Med Int Health*, 6(8), 607-613.
- Ranjit, M. R., Das, A., Das, B. P., Das, B. N., Dash, B. P., & Chhotray, G. P. (2005). Distribution of *Plasmodium falciparum* genotypes in clinically mild and severe malaria cases in Orissa, India. *Trans R Soc Trop Med Hyg*, 99(5), 389-395.
- Robert, F., Ntoumi, F., Angel, G., Candito, D., Rogier, C., Fandeur, T., et al. (1996). Extensive genetic diversity of *Plasmodium falciparum* isolates collected from patients with severe malaria in Dakar, Senegal. *Trans R Soc Trop Med Hyg*, 90(6), 704-711.
- Rubio, J. M., Berzosa, P. J., & Benito, A. (2001). Amplified fragment length polymorphism (AFLP) protocol for genotyping the malarial parasite *Plasmodium falciparum*. *Parasitology*, 123(Pt 4), 331-336.
- Sachs, J., & Malaney, P. (2002). The economic and social burden of malaria. *Nature*, 415(6872), 680-685.
- Sakihama, N., Ohmae, H., Bakote'e, B., Kawabata, M., Hirayama, K., & Tanabe, K. (2006). Limited allelic diversity of *Plasmodium falciparum* merozoite surface protein 1 gene from populations in the Solomon Islands. *Am J Trop Med Hyg*, 74(1), 31-40.
- Sallenave-Sales, S., Ferreira-da-Cruz, M. F., Faria, C. P., Cerruti, C., Jr., Daniel-Ribeiro, C. T., & Zalis, M. G. (2003). *Plasmodium falciparum*: limited genetic diversity of

- MSP-2* in isolates circulating in Brazilian endemic areas. *Exp Parasitol*, 103(3-4), 127-135.
- Schoepflin, S., Valsangiacomo, F., Lin, E., Kiniboro, B., Mueller, I., & Felger, I. (2009). Comparison of *Plasmodium falciparum* allelic frequency distribution in different endemic settings by high-resolution genotyping. *Malar J*, 8, 250.
- Silva, N. S., Silveira, L. A., Machado, R. L., Povoa, M. M., & Ferreira, M. U. (2000). Temporal and spatial distribution of the variants of merozoite surface protein-1 (*MSP-1*) in *Plasmodium falciparum* populations in Brazil. *Ann Trop Med Parasitol*, 94(7), 675-688.
- Smith, T., Beck, H. P., Kitua, A., Mwankusye, S., Felger, I., Fraser-Hurt, N., et al. (1999). Age dependence of the multiplicity of *Plasmodium falciparum* infections and of other malariological indices in an area of high endemicity. *Trans R Soc Trop Med Hyg*, 93 Suppl 1, 15-20.
- Smythe, J. A., Coppel, R. L., Brown, G. V., Ramasamy, R., Kemp, D. J., & Anders, R. F. (1988). Identification of two integral membrane proteins of *Plasmodium falciparum*. *Proc Natl Acad Sci U S A*, 85(14), 5195-5199.
- Smythe, J. A., Peterson, M. G., Coppel, R. L., Saul, A. J., Kemp, D. J., & Anders, R. F. (1990). Structural diversity in the 45-kilodalton merozoite surface antigen of *Plasmodium falciparum*. *Mol Biochem Parasitol*, 39(2), 227-234.
- Snewin, V. A., Herrera, M., Sanchez, G., Scherf, A., Langsley, G., & Herrera, S. (1991). Polymorphism of the alleles of the merozoite surface antigens MSA1 and MSA2 in *Plasmodium falciparum* wild isolates from Colombia. *Mol Biochem Parasitol*, 49(2), 265-275.

- Snounou, G., Viriyakosol, S., Jarra, W., Thaithong, S., & Brown, K. N. (1993). Identification of the four human malaria parasite species in field samples by the polymerase chain reaction and detection of a high prevalence of mixed infections. *Mol Biochem Parasitol*, 58(2), 283-292.
- Snounou, G., Zhu, X., Siripoon, N., Jarra, W., Thaithong, S., Brown, K. N., et al. (1999). Biased distribution of *MSP-1* and *MSP-2* allelic variants in *Plasmodium falciparum* populations in Thailand. *Trans R Soc Trop Med Hyg*, 93(4), 369-374.
- Snow, R. W., Guerra, C. A., Noor, A. M., Myint, H. Y., & Hay, S. I. (2005). The global distribution of clinical episodes of *Plasmodium falciparum* malaria. *Nature*, 434(7030), 214-217.
- Soulama, I., Nebie, I., Ouedraogo, A., Gansane, A., Diarra, A., Tiono, A. B., et al. (2009). *Plasmodium falciparum* genotypes diversity in symptomatic malaria of children living in an urban and a rural setting in Burkina Faso. *Malar J*, 8, 135.
- Soulama, I., Sawadogo, M., Nebie, I., Diarra, A., Tiono, A. B., Konate, A., et al. (2006). Genetic diversity of *Plasmodium falciparum* and pathogenesis of the severe malarial anaemia in children under 5 years old in the province of Boulgou, Burkina Faso. *Bull Soc Pathol Exot*, 99(3), 166-170.
- Subbarao, S. K., & Sharma, V. P. (1997). Anopheline species complexes & malaria control. *Indian J Med Res*, 106, 164-173.
- Takala, S., Branch, O., Escalante, A. A., Kariuki, S., Wootton, J., & Lal, A. A. (2002). Evidence for intragenic recombination in *Plasmodium falciparum*: identification of a novel allele family in block 2 of merozoite surface protein-1: Asembo Bay Area Cohort Project XIV. *Mol Biochem Parasitol*, 125(1-2), 163-171

- Tibayrenc, M. (1998). Beyond strain typing and molecular epidemiology: integrated genetic epidemiology of infectious diseases. *Parasitol Today*, 14(8), 323-329.
- Toebe, C. S., Clements, J. D., Cardenas, L., Jennings, G. J., & Wiser, M. F. (1997). Evaluation of immunogenicity of an oral *Salmonella* vaccine expressing recombinant *Plasmodium berghei* merozoite surface protein-1. *Am J Trop Med Hyg*, 56(2), 192-199.
- Uthaipibull, C., Aufiero, B., Syed, S. E., Hansen, B., Guevara Patino, J. A., Angov, E., et al. (2001). Inhibitory and blocking monoclonal antibody epitopes on merozoite surface protein 1 of the malaria parasite *Plasmodium falciparum*. *J Mol Biol*, 307(5), 1381-1394.
- Vafa, M., Troye-Blomberg, M., Anchang, J., Garcia, A., & Migot-Nabias, F. (2008). Multiplicity of *Plasmodium falciparum* infection in asymptomatic children in Senegal: relation to transmission, age and erythrocyte variants. *Malar J*, 7, 17.
- Vargas-Serrato, E., Barnwell, J. W., Ingravallo, P., Perler, F. B., & Galinski, M. R. (2002). Merozoite surface protein-9 of *Plasmodium vivax* and related simian malaria parasites is orthologous to p101/ABRA of *P. falciparum*. *Mol Biochem Parasitol*, 120(1), 41-52.
- Walliker, D. (1983). The genetic basis of diversity in malaria parasites. *Adv Parasitol*, 22, 217-259.
- White, N. J. (2008). *Plasmodium knowlesi*: the fifth human malaria parasite. *Clin Infect Dis*, 46(2), 172-173.
- World Health Organization (2008). "World Malaria Report 2008.
- World Health Organization (2004). "World Malaria Report 2004.

World Health Organization (2009). "World Malaria Report 2009.

World Health Organization (2010). "World Malaria Report 2010.

Wongsrichanalai, C., Barcus, M. J., Muth, S., Sutamihardja, A., & Wernsdorfer, W. H. (2007). A review of malaria diagnostic tools: microscopy and rapid diagnostic test (RDT). *Am J Trop Med Hyg*, 77(6 Suppl), 119-127.

Zakeri, S., Bereczky, S., Naimi, P., Pedro Gil, J., Djadid, N. D., Farnert, A., et al. (2005). Multiple genotypes of the merozoite surface proteins 1 and 2 in *Plasmodium falciparum* infections in a hypoendemic area in Iran. *Trop Med Int Health*, 10(10), 1060-1064.

Zwetyenga, J., Rogier, C., Tall, A., Fontenille, D., Snounou, G., Trape, J. F., et al. (1998). No influence of age on infection complexity and allelic distribution in *Plasmodium falciparum* infections in Ndiop, a Senegalese village with seasonal, mesoendemic malaria. *Am J Trop Med Hyg*, 59(5), 726-735.