

Journal of Bioresource Management

Volume 5 | Issue 4

Article 2

Status and Food Preferences of Bears in Sub Alpine Scrub Forests, AJK

Madeeha Manzoor

Bioresource Research Centre, Islamabad, madeemanzoor@hotmail.com

Safdar Ali Shah

Khyber Pakhtunkhwa (KP) Wildlife Department, Peshawar, Pakistan

Jibran Haider

Forest and wildlife Department, Gilgit-Baltistan, Pakistan

Follow this and additional works at: <https://corescholar.libraries.wright.edu/jbm>



Part of the [Biodiversity Commons](#), [Forest Biology Commons](#), and the [Zoology Commons](#)

Recommended Citation

Manzoor, M., Shah, S. A., & Haider, J. (2018). Status and Food Preferences of Bears in Sub Alpine Scrub Forests, AJK, *Journal of Bioresource Management*, 5 (4).

DOI: <https://doi.org/10.35691/JBM.8102.0096>

ISSN: 2309-3854 online

(Received: Dec 10, 2019; Accepted: Dec 10, 2019; Published: Dec 10, 2018)

This Article is brought to you for free and open access by CORE Scholar. It has been accepted for inclusion in *Journal of Bioresource Management* by an authorized editor of CORE Scholar. For more information, please contact library-corescholar@wright.edu.

Status and Food Preferences of Bears in Sub Alpine Scrub Forests, AJK

© Copyrights of all the papers published in Journal of Bioresource Management are with its publisher, Center for Bioresource Research (CBR) Islamabad, Pakistan. This permits anyone to copy, redistribute, remix, transmit and adapt the work for non-commercial purposes provided the original work and source is appropriately cited. Journal of Bioresource Management does not grant you any other rights in relation to this website or the material on this website. In other words, all other rights are reserved. For the avoidance of doubt, you must not adapt, edit, change, transform, publish, republish, distribute, redistribute, broadcast, rebroadcast or show or play in public this website or the material on this website (in any form or media) without appropriately and conspicuously citing the original work and source or Journal of Bioresource Management's prior written permission.

STATUS AND FOOD PREFERENCES OF BEARS IN SUB ALPINE SCRUB FORESTS, AJK

MADEEHA MANZOOR^{1*}, SAFDAR ALI SHAH², JIBRAN HAIDER³

¹Center for Bioresource Research (CBR), Pakistan

²Khyber Pakhtunkhwa (KP) Wildlife Department, Peshawar, Pakistan

³Forest and wildlife Department, Gilgit-Baltistan, Pakistan

*Corresponding author: madeemanzoor@hotmail.com

ABSTRACT

Bears belong to the family Ursidae, under the order Carnivora with eight extant species (Grzimek, 2003). Roberts (1997) cite the presence of Himalayan brown bear, Himalayan black bear and Balochistan black bear in Pakistan. Dhirkot National Park (2008), Banjosa Nature Reserve (2009), Tolipir National Park (2008), Pir Chanasi National Park (2010) and Pir Lasura National Park (2009) were surveyed. A broad-spectrum investigation was conducted where the number of bears observed directly; pictures from surveillance cameras (Arc View GIS 3.3, 1992-2002) and at the same time indirect indicators such as bear scats, walking trails, paw prints and fur samples were noted for presence of bears. Only two types of bears, the Asiatic black bear and brown bear were reported from the study area. the Asiatic black bear was observed in four of the five study areas (Banjosa Nature Reserve, Dhirkot National Park, Tolipir National Park and Pir Chanasi National Park). Pir Chanasi had the highest numbers of bears observed while only four bears were reported from Dhirkot National Park. From the whole study site, seven brown bears were sighted only in Pir Chanasi National Park.

Keywords: Asiatic black bear, brown bear, carnivore, omnivore, forest

INTRODUCTION

Bears belong to the family Ursidae, under the order Carnivora with eight extant species (Grzimek, 2003). These include the giant panda, Malayan sun bear, Asiatic black bear, American black bear, brown bear, spectacled bear (*Tremarcto ornatus*), sloth bear (*Melursus ursinus*) and polar bear (American Bear Association, n.d.). Roberts (1997) cite the presence of Himalayan brown bear, Himalayan black bear and Balochistan black bear in Pakistan. The IUCN redlist (2018) places the *Helarctos malayanus*, *Melursus ursinus*, *Tremarcto ornatus*, *Ursus maritimus* and *Ursus thibetanus* in the vulnerable category with a decreasing trend of population except for polar bears. Due to the extreme geographic location that this species inhabits, only limited data is available regarding their population. The brown bear is

categorized under the least concern category and with a stable population. The American black bear is also categorized under least concern but with an increasing trend in population.

The habitat preferences of these species are very diverse. Polar bears prefer the cold arctic tundra whereas pandas inhabit Chinese bamboo forests. Sloth bears (*Melursus ursinus*) prefer grassy lowlands. Brown bears inhabit a variety of biomes such as the tundra, forests and grasslands.

Bears are predominantly nocturnal but have also been observed catching prey in daylight. Most bears remain active all year round. Polar bears and the Asiatic black bear do not hibernate but their sleeping patterns change and become longer in colder weather. Their body temperature lowers during this time. According to Grzimek (2003) Ursids

are omnivores and select their diet based on location and availability.

For a long time, this family has been hunted mainly as a show of power but also for their fur and meat. Their body parts have been exploited for their medicinal value. Their wild community has also been affected by their capture for entertainment in zoos or by gypsies that train bears for dancing. This study was conducted to report the number of bears present in the study area and their feeding preferences using physical observation and previously published literature.

MATERIALS AND METHOD

Dhirkot National Park (2008), Banjosa Nature Reserve (2009), Tolipir National Park (2008), Pir Chanasi National Park (2010) and Pir Lasura National Park (2009) were surveyed. A broad-spectrum investigation was conducted where the number of bears observed directly; pictures from surveillance cameras (Arc View GIS 3.3, 1992-2002) and at the same time indirect indicators such as bear scats, walking trails, paw prints and fur samples were noted for presence of bears. Data was also collected by interviewing locals and forest rangers for information along with review of literature to determine possible locations the bears may be found.

RESULTS

Table 1. List of mammalian species of family Ursidae recorded from AJK

S. #	Names	Food	Remarks	Local sightings (%)
1.	<i>Ursus thibetanus</i> Asiatic Black Bear	Acorns, mulberries, apricots, Rosehips, carrion, goat, sheep, small crustaceans, Fungus or forest mushrooms, maize, grass, ber (<i>Ziziphus nummularia</i>), Russian olive (<i>Eleagnus hortensis</i>), lizards, orthopterous insects including locusts, rhizomatous stems, sorghum (<i>Sorghum suanense</i>) and ripe dates (<i>Phoenix dactylifera</i>).	Widely distributed mammal, occupying wide variety of habitat, and not nocturnal, acute sense of smell. Breeds from March-April Liters size: 1-2	(12- BNR) (4- DNP) (22-TNP) (24-PCNP)
2.	<i>Ursus arctos</i> (Brown Bear or Snow Bear)	vegetable matter with rodent bones (Omnivorous, feed on alpine bulbs and insects crustaceans etc)	There is continuous threat to its survival in Pakistan due to increasing demand by gypsy tribes. Breeds from March-April Liters size: 1-2	(7-PCNP)



a



b

Figure 1. (a) Asiatic Black bear (*Ursus thibetanus*) (b) Brown bear (*Ursus arctos*)

DISCUSSION

Only two types of bears, the Asiatic black bear and brown bear were reported from the study area.

The Asian Black Bear *Ursus thibetanus* is native to Asia (the Indian subcontinent, Korean peninsula, China, Russia, Taiwan and Japan) (Heptner and Naumov, 1992; Montgomery, 2002). Since these bears prefer to inhabit moist forests, it is important to conserve the woods to protect their habitat and as a result their population. *Ursus thibetanus* was observed in four of the five study areas (Banjosa Nature Reserve, Dhirkot National Park, Tolipir National Park and Pir Chanasi National Park). Pir Chanasi had the highest numbers of bears observed while only four bears were reported from Dhirkot National Park.

According to Abbas et al. (2015) *U thibetanus* may be found in the highlands ranging from 1,500 m to 4,000 m above sea level. However, they note a possible decrease in their numbers during the 1900s. Sheikh and Molur (2005) report the status of Asiatic black bears to be vulnerable in Pakistan. Ali et al. (2017) suggest reduced food availability as one of the reasons which influences bears to attack crops and livestock reared for food by humans, which would result in a competition for food.

Bears belonging to this species have a wide-ranging diet. From berries and wild mushrooms to crustaceans and small animals, the diet of *U thibetanus* varies with changing seasons. This finding is concurrent with the research by Ali et al. (2017) where they found remnants of 21 different food items in diets of bears in their study site.

Nawaz (2007) report that the number of Brown bears were on a fast decline, to the point where they have completely vanished from some parts of Pakistan. A similar result was observed in the current study, where the population of brown bears was only sighted in Pir Chanasi National Park and the number of brown bears seen there was also only seven. To protect these bears, efforts need to be expended to conserve their numbers. A good example of such measures is Deosai National Park where Nawaz et al. (2008) conducted a study on brown bears and reported a five percent annual increase in their population. Qureshi et al. (2011) conducted a study in Khunjerab National Park and observed the presence of Brown bears. Some of the factors that may affect the population of brown bears in Khunjerab National Park included trophy hunting and carcass poisoning. There is no confirmed evidence that this is also true for the current study site, but may be a possible reason for the decline in brown bear population.

According to McLellan et al. (2017) near Bhimbar, Kotli and Peshawar the population of brown bears have become extinct.

CONCLUSION

Only two types of bears, the Asiatic black bear and brown bear were reported from the study area. *Ursus thibetanus* was observed in four of the five study areas (Banjosa Nature Reserve, Dhirkot National Park, Tolipir National Park and Pir Chanasi National Park). Pir Chanasi had the highest numbers of bears observed while only four bears were reported from Dhirkot National Park. From berries and wild mushrooms to crustaceans and small animals, the diet of *U thibetanus* varies with changing seasons. The population of brown bears was only sighted in Pir Chanasi National Park.

REFERENCES

- Ali A, Zhixiang Z, Waseem M, Khan MF, Ali I, Asad M, Qashqaei AT (2017). An assessment of food habits and altitudinal distribution of the Asiatic black bear (*Ursus thibetanus*) in the Western Himalayas, Pakistan. *Journal of Natural History*, 51 (11–12): 689–701. <http://dx.doi.org/10.1080/00222933.2017.1303097>
- Arc View GIS 3.3 (1992-2002). Arc View GIS 3.3 software, Environmental Systems Research Institute, Inc. HCL Technologies LTD, New Delhi, India.
- Garshelis DL, Steinmetz R (2016). "Ursus thibetanus". The IUCN Red List of Threatened Species.
- Google Earth 5.0. (2009). Google Earth 5.0. programme. Europa Technologies. Data SIO, NOAA, U.S. Navy, NGS, GEBCO. Tele Atlas.
- Grzimek (2003). Grzimek's Animal Life Encyclopedia. Vol 14. Mammals III, 2nd ed. Thomson Gale London. 295-306.
- Heptner VG, Naumov NP (1992). "White-chested, black bear". *Mlekopitajušćie Sovetskogo Soiuza*. Moskva: Vysšaia Škola [Mammals of the Soviet Union. Vol. II Part 1a, Sirenia and Carnivora (Sea cows; Wolves and Bears)]. Washington DC: Smithsonian Institution and the National Science Foundation. pp. 713–733.
- McLellan BN, Proctor MF, Huber D, Michel S (2017). *Ursus arctos*. The IUCN Red List of Threatened Species. <http://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T41688A121229971.en>.
- Montgomery S (2002). Search for the golden moon bear: science and adventure in Southeast Asia. Simon & Schuster.
- Nawaz MA (2007). Status of the brown bear in Pakistan. *Ursus*, 18(1): 89-100. [https://doi.org/10.2192/1537-6176\(2007\)18\[89:SOTBBI\]2.0.CO;2](https://doi.org/10.2192/1537-6176(2007)18[89:SOTBBI]2.0.CO;2)
- Oishimaya SN (2017) How Many Types Of Bears Are There Living In The World Today?, WorldAtlas, <https://www.worldatlas.com/articles/how-many-types-of-bears-are-there.html>.
- Roberts TJ (2005 a). Field Guide to the Large and Medium-sized Mammals of Pakistan. 1st ed., Oxford University Press New York. Pp 280.
- Roberts TJ (2005 b). The Mammals of Pakistan. 1st ed. Oxford University Press New York. Pp 559.
- Scotson L, Fredriksson G, Augeri D, Cheah C, Ngoprasert D, Wai-Ming W (2017). *Helarctos malayanus*. The IUCN Red List of Threatened Species 2017. <http://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T9760A45033547.en>.

Sokal RR, Rohlf FJ (2000). *Biometry: The Principles and Practice of Statistics in Biological Research*. W. H. Freeman and Co., New York.

Than U, Simcharoen S, Kanchanasakha B (1998). *Carnivores of Mainland South East Asia*. Bangkok: Siam Tong Kit Printing Co., Ltd.