

SYMMETRY AND UNIQUENESS OF SOLUTIONS TO SOME LIOUVILLE-TYPE EQUATIONS AND SYSTEMS

CHANGFENG GUI, ALEKS JEVIKAR, AMIR MORADIFAM

ABSTRACT. We prove symmetry and uniqueness results for three classes of Liouville-type problems arising in geometry and mathematical physics: asymmetric Sinh-Gordon equation, cosmic string equation and Toda system, under certain assumptions on the mass associated to these problems. The argument is in the spirit of the Sphere Covering Inequality which for the first time is used in treating different exponential nonlinearities and systems.

1. INTRODUCTION

In this paper, we shall consider three classes of Liouville-type equations and systems: asymmetric Sinh-Gordon equation, cosmic string equation and Toda system. These problems arise in geometry and mathematical physics. We are mainly concerned about the symmetry and uniqueness questions under certain assumptions on the mass associated to these problems.

1.1. Asymmetric Sinh-Gordon equation. Consider the following version of the asymmetric Sinh-Gordon equation

$$(1) \quad \begin{cases} -\Delta u = \rho \frac{e^u + \frac{\alpha}{|\alpha|} e^{\alpha u}}{\int_{\Omega} (e^u + e^{\alpha u}) dx} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\alpha \in [-1, 1], \alpha \neq 0, \rho > 0$ is a parameter and $\Omega \subset \mathbb{R}^2$ is a bounded domain with smooth boundary $\partial\Omega$. Equation (1) is known also as Neri's mean field equation and arises in the context of the statistical mechanics description of $2D$ -turbulence introduced in [43]. In the model where the circulation number density is subject to a probability measure, under a *stochastic* assumption on the vortex intensities one obtains the following equation (see [42]):

$$(2) \quad \begin{cases} -\Delta u = \rho \int_{[-1,1]} \beta \frac{e^{\beta u} \mathcal{P}(d\beta)}{\iint_{[-1,1] \times \Omega} e^{\beta u} \mathcal{P}(d\beta) dx} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

2000 *Mathematics Subject Classification.* 35J61, 35R01, 35A02, 35B06.

Key words and phrases. Geometric PDEs, Sinh-Gordon equation, Cosmic string equation, Toda system, Sphere covering inequality, Symmetry results, Uniqueness results.

The first author is partially supported by NSF grant DMS-1601885 and NSFC grant No 11371128. The second author is supported by PRIN12 project: *Variational and Perturbative Aspects of Nonlinear Differential Problems* and FIRB project: *Analysis and Beyond*. The third author is partially supported by NSF grant DMS-1715850.

where u stands for the stream function of a turbulent Euler flow, \mathcal{P} is a Borel probability measure defined in $[-1, 1]$ describing the point vortex intensity distribution and $\rho > 0$ is a physical constant associated to the inverse temperature. Equation (1) is related to the latter model when \mathcal{P} is supported in two points.

On the other hand, a *deterministic* assumption on the vortex intensities yields the following model (see [51]):

$$(3) \quad \begin{cases} -\Delta u = \rho \left(\frac{e^u}{\int_{\Omega} e^u dx} + \frac{\alpha}{|\alpha|} \frac{e^{\alpha u}}{\int_{\Omega} e^{\alpha u} dx} \right) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Concerning the analysis of the latter equation we refer the interested readers to [24, 25, 26, 27, 29, 30, 31, 32, 47, 49]. The arguments presented here do not apply to (3), and we postpone its analysis to a forthcoming paper.

Observe that by taking $\alpha = -1$ in (1) we end up with the standard Sinh-Gordon equation, while for \mathcal{P} supported in a single point we derive the standard mean field equation

$$(4) \quad \begin{cases} -\Delta u = \rho \frac{e^u}{\int_{\Omega} e^u dx} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

which is related to the prescribed Gaussian curvature problem and Euler flows (see [2, 13, 14] and [9, 35], respectively). The latter equation has been widely studied and we refer to the surveys [40, 54]. We note that even though the equation in (4) is related to geometric applications, the Dirichlet boundary conditions are usually not natural in this geometric setting. Recently in [21, 22, 23] the authors proved the Sphere Covering Inequality (see Theorem 2.5 below) which leads to several symmetry and uniqueness results for the latter equation. The Sphere Covering Inequality [21] will also be a crucial tool in this paper.

Returning to (1), some partial existence results and blow-up analysis was carried out in [48, 50], while a complete existence result for (2) with $\text{supp } \mathcal{P} \subset [0, 1]$ was given in [18]. On the other hand, we are not aware of any symmetry or uniqueness results for the latter equation with the only exception of [52] where (3) is considered. We present here several results in this direction, under natural assumptions both on the parameter ρ and the domain Ω . Due to different features of problem (2) depending on whether $\text{supp } \mathcal{P} \subset [0, 1]$ or $\text{supp } \mathcal{P} \subset [-1, 1]$ we will distinguish these two cases in the discussion below. In the first situation we may rewrite (1) as

$$(5) \quad \begin{cases} -\Delta u = \rho \frac{e^u + e^{au}}{\int_{\Omega} (e^u + e^{au}) dx} & \text{in } \Omega, \\ u = g(x) \geq 0 & \text{on } \partial\Omega, \end{cases}$$

with $a \in (0, 1)$. Our first result is the following.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^2$ be a bounded, simply-connected domain and $g \in C(\partial\Omega)$ be a non-negative function. Suppose $\rho \leq 4\pi$. If u_1 and u_2 are two solutions of (5) such that*

$$(6) \quad \int_{\Omega} (e^{u_1} + e^{au_1}) dx = \int_{\Omega} (e^{u_2} + e^{au_2}) dx,$$

then $u_1 \equiv u_2$.

Corollary 1.2. *Under the condition of Theorem 1.1, assume further Ω and g are evenly symmetric about a line. Then, any solution of (5) must be evenly symmetric about that line. In particular, if Ω is radially symmetric and g is a non-negative constant, then u is radially symmetric.*

We will exploit the fact that for $\text{supp } \mathcal{P} \subset [0, 1]$ equation (2) shares some features with the mean field equation (4). Indeed we shall rewrite (2) in the form of (4) and apply the Sphere Covering Inequality (see [21]) to get the desired results.

Remark 1.3. *The argument for Theorem 1.1 can be adapted to treat the more general case where the probability measure \mathcal{P} in (2) is supported at $(m + 1)$ points, i.e.*

$$\begin{cases} -\Delta u = \rho \frac{e^u + e^{a_1 u} + \dots + e^{a_m u}}{\int_{\Omega} (e^u + e^{a_1 u} + \dots + e^{a_m u}) dx} & \text{in } \Omega, \\ u = g \geq 0 & \text{on } \partial\Omega, \end{cases}$$

with $a_i \in (0, 1)$ for all i . Indeed if $\rho \leq \frac{8\pi}{m+1}$ and

$$\int_{\Omega} (e^{u_1} + e^{a_1 u_1} + \dots + e^{a_m u_1}) dx = \int_{\Omega} (e^{u_2} + e^{a_1 u_2} + \dots + e^{a_m u_2}) dx,$$

then we must necessarily have $u_1 \equiv u_2$. In particular, Corollary 1.2 also generalizes to the above equation. The case where $a_i > 1$ for some i can be carried out as well and we refer to Remark 1.5 for more details.

On the other hand, for the general case $\text{supp } \mathcal{P} \subset [-1, 1]$, the problem (2) substantially differs from the standard equation (4). In this case we may rewrite (1) as

$$(7) \quad \begin{cases} -\Delta u = \rho \frac{e^u - e^{-au}}{\int_{\Omega} (e^u + e^{-au}) dx} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

with $a \in (0, 1]$. Observe that $u \equiv 0$ is a solution of the latter problem. We indeed show that for $\rho \leq \frac{8\pi}{1+a}$ the trivial solution is the only solution.

Theorem 1.4. *Suppose $\rho \leq \frac{8\pi}{1+a}$ and $\Omega \subset \mathbb{R}^2$ bounded, simply-connected. Then, equation (7) admits only the trivial solution $u \equiv 0$.*

The proof is based on the Sphere Covering Inequality (see Section 2 in [21]). Roughly speaking, letting $v_1 = u$, $v_2 = -au$ we will consider a symmetrization of $v_2 - v_1$ with respect to two suitable measures to get the conclusion.

Remark 1.5. *Let us point out that in equations (5) and (7) we are considering $a < 1$ and $a \leq 1$ (respectively) due to the physical motivations. However, we can treat the case $a > 1$ as well. More precisely, letting $v = au$ in (5) we may rewrite the latter equation in a form to which we can apply Theorem 1.2 with a new parameter $\tilde{\rho} = a\rho$. Therefore, the conclusions of Theorem 1.1 and Corollary 1.2 still hold true for $\rho \leq \frac{4\pi}{a}$ and $a > 1$. On the other hand, one can easily see from the proof of Theorem 1.4 that the assumption $a \leq 1$ is not needed and we get the same conclusion for $a > 1$.*

Remark 1.6. *The same arguments clearly apply to the following version of (1):*

$$(8) \quad \begin{cases} -\Delta u = e^u + \frac{\alpha}{|\alpha|} e^{\alpha u} & \text{in } \Omega, \\ u = g(x) \geq 0 & \text{on } \partial\Omega. \end{cases}$$

We have:

1. Let $\alpha = a \in (0, 1)$. Suppose $\Omega \subset \mathbb{R}^2$ is a bounded, simply-connected domain and $g \in C(\partial\Omega)$ is a non-negative function. If u_1 and u_2 are two solutions of (8) such that

$$\int_{\Omega} e^{u_1} dx = \int_{\Omega} e^{u_2} dx \leq 4\pi,$$

then $u_1 \equiv u_2$.

Moreover, suppose that Ω and g are evenly symmetric about a line. Let u be a solution of (8). Then, u is evenly symmetric about that line. In particular, if Ω is radially symmetric and g is a non-negative constant, then u is radially symmetric.

2. Let $\alpha = -a$, $a \in (0, 1]$. Suppose $\Omega \subset \mathbb{R}^2$ bounded, simply-connected. If u is a solution of (8) with $g = 0$ such that

$$\int_{\Omega} (e^u + e^{-au}) dx \leq \frac{8\pi}{1+a},$$

then $u \equiv 0$.

Moreover, similar results hold for $a > 1$ (see Remark 1.5).

The above results follow by suitably adapting the proofs of Theorem 1.1, Corollary 1.2 and Theorem 1.4 and we omit the details here.

Finally, we have the following remark concerning the sharpness of the above results.

Remark 1.7. *Consider for simplicity the standard Sinh-Gordon equation with $\alpha = -1$ in (1). Even though the associated energy functional is coercive for $\rho < 8\pi$ (see [50]), we can not extend Theorem 1.1, Corollary 1.2 and Theorem 1.4 to the range $\rho \leq 8\pi$ (as it holds for the standard mean field equation (4)). In [52] (Section 2) the authors provide non-trivial solutions for (3) with $\rho < 8\pi$.*

1.2. Cosmic String Equation. We will next discuss the following problem to which we will refer to as the cosmic string equation:

$$(9) \quad \begin{cases} -\Delta u = e^{au} + h(x)e^u & \text{in } \Omega, \\ u = g(x) \geq 0 & \text{on } \partial\Omega, \end{cases}$$

with $a > 0$, and $\Omega \subset \mathbb{R}^2$ is a smooth bounded domain containing the origin and h is of the form

$$(10) \quad h(x) = e^{-4\pi N G_0(x)},$$

where $N \in \mathbb{N}$ and G_0 is the Green's function with pole at 0, i.e.

$$(11) \quad \begin{cases} -\Delta G_0(x) = \delta_0 & \text{in } \Omega, \\ G_0(x) = 0 & \text{on } \partial\Omega. \end{cases}$$

Observe that

$$h > 0 \quad \text{in } \Omega \setminus \{0\} \quad \text{and} \quad h(x) \cong |x|^{2N} \quad \text{near } 0.$$

Equation (9) describes the behavior of selfgravitating cosmic strings for a massive W-boson model coupled with Einstein's equation where a is a physical parameter and N the string's multiplicity (see [44, 57]). Observe that for $a = 1$ the equation (9) is also related to the Gaussian curvature with conic singularities (see [54] and references therein).

Many results concerning (9) have been established especially for the full plane case. We refer to [11, 12, 57] for existence results, to [44, 45] for what concerns symmetry issues, and to [56] for blow-up analysis. In particular, in [44, 45] the authors provide necessary and sufficient conditions for the solvability of (9) in the full plane in the context of radially symmetric solutions, depending on the values of the total mass $\beta = \int_{\mathbb{R}^2} (e^{au} + |x|^{2N} e^u) dx$. For $N \in (-1, 0]$ it follows from a moving plane argument that all the solutions to (9) are radially symmetric, under suitable assumptions on the domain Ω . However, it remains an open problem if the results in [44, 45] are sharp for the non-radial framework. We prove the following result.

Theorem 1.8. *Let $\Omega \subset \mathbb{R}^2$ be a bounded, simply-connected domain, $a > 0$, $N \geq 0$ and $g \in C(\partial\Omega)$ be non-negative. Suppose u_1 and u_2 are two distinct solutions of (9) such that*

$$(12) \quad \begin{cases} \int_{\Omega} (e^{au_1} + e^{au_2}) dx \leq \frac{8\pi}{a} & \text{if } a \geq 1, \\ \int_{\Omega} (e^{u_1} + e^{u_2}) dx \leq 8\pi & \text{if } a < 1 \end{cases}$$

Then u_1 and u_2 can not intersect, i.e. either

$$(13) \quad u_2 > u_1 \quad \text{or} \quad u_2 < u_1 \quad \text{in } \Omega.$$

Corollary 1.9. *Let $\Omega \subset \mathbb{R}^2$ be a bounded, simply-connected domain, $a > 0$, $N \geq 0$ and $g \in C(\partial\Omega)$ be non-negative. Assume*

$$(14) \quad \gamma := \begin{cases} \int_{\Omega} e^{au} dx \leq \frac{4\pi}{a} & \text{if } a \geq 1, \\ \int_{\Omega} e^u dx \leq 4\pi & \text{if } a < 1. \end{cases}$$

Then (9) has a unique solution u for any γ satisfying (14). In particular, if $0 \in \Omega$ and Ω, g are evenly symmetric about a line passing through the origin, then u is evenly symmetric about that line. Consequently, if Ω is radially symmetric about the origin and g is a non-negative constant, then u is radially symmetric about the origin.

The proof is based on a simple manipulation of equation (9) and the Sphere Covering Inequality (see Theorem 2.5 below or [21]).

Remark 1.10. *Theorem 1.8 and Corollary 1.9 can be generalized for the following more general equation (we refer to [45] for applications of this equation)*

$$\begin{cases} -\Delta u = \sum_{i=0}^m h_i(x) e^{a_i u} & \text{in } \Omega, \\ u = g(x) \geq 0 & \text{on } \partial\Omega, \end{cases}$$

where $a_i > 0$ and

$$h_i(x) = e^{-4\pi N_i G_0(x)},$$

with $N_i \geq 0$ for all i . Let $a_M = \max_i \{a_i\}$. Using similar arguments as in the proofs of Theorem 1.8, one can check the assumptions (12) and (14) (where $m = 1$) should be replaced by

$$\int_{\Omega} (e^{a_M u_1} + e^{a_M u_2}) dx \leq \frac{16\pi}{a_M(m+1)},$$

and

$$\int_{\Omega} e^{a_M u} dx \leq \frac{8\pi}{a_M(m+1)},$$

respectively.

1.3. Liouville-Type Systems. We also study the following class of Liouville-type systems:

$$(15) \quad \begin{cases} -\Delta u_1 = A e^{u_1} - B e^{u_2} & \text{in } \Omega, \\ -\Delta u_2 = B' e^{u_2} - A' e^{u_1} & \text{in } \Omega, \\ u_1 = u_2 = g(x) & \text{on } \partial\Omega, \end{cases}$$

with $g \in C(\partial\Omega)$ and

$$(16) \quad A, A', B, B' > 0, \quad A + A' = B + B' := M > 0.$$

Observe that we allow some of the above coefficients to be zero.

The latter system is deeply connected both with geometry and mathematical physics. For example, by taking $A = B' = 2$, $B = A' = 1$ we recover the 2×2 Toda system which has been extensively studied in the literature. This equation appears in the description of holomorphic curves in $\mathbb{C}\mathbb{P}^N$ (see [8, 10, 39]). It also arises in the non-abelian Chern-Simons theory in the context of high critical temperature superconductivity (see [19, 56, 57]). The case $A = B' = 1$ and $B = A' = \tau$ with a singular source was considered in [46] in unbounded domains.

For what concerns Toda-type systems we refer to [33, 37, 38] for blow-up analysis, to [39] for classification issues, and to [6, 28, 41] for existence results. On the other hand, we are not aware of any symmetry or uniqueness results for Liouville-type systems alike (15). In this direction we provide the following result.

Theorem 1.11. *Let (u_1, u_2) be a solution of (15) and (16). Let M be as defined in (16). Suppose that Ω is a bounded, simply-connected domain and*

$$\int_{\Omega} (e^{u_1} + e^{u_2}) dx \leq \frac{8\pi}{M}.$$

Then $u_1 \equiv u_2 \equiv u$, where u is the unique solution to

$$\begin{cases} -\Delta u = De^u & \text{in } \Omega, \\ u = g(x) & \text{on } \partial\Omega, \end{cases}$$

and $D := A - B = B' - A'$.

Remark 1.12. For Toda-type systems where $A = B' = 2$, $B = A' = 1$, the above result asserts that if Ω is bounded, simply-connected and

$$\int_{\Omega} (e^{u_1} + e^{u_2}) dx \leq \frac{8\pi}{3},$$

then $u_1 \equiv u_2 \equiv u$, where u is the unique solution to

$$\begin{cases} -\Delta u = e^u & \text{in } \Omega, \\ u = g(x) & \text{on } \partial\Omega. \end{cases}$$

Arguing as in the proof of the Sphere Covering Inequality (see Section 2 below or [21]), we will consider a symmetrization of $u_2 - u_1$ with respect to two suitable measures to get the latter result. The uniqueness property will then follow by applying the Sphere Covering inequality to the scalar equation.

A similar argument can be carried out for the following singular version of (15):

$$(17) \quad \begin{cases} -\Delta u_1 = Ae^{u_1} - Be^{u_2} - 4\pi\alpha\delta_0 & \text{in } \Omega, \\ -\Delta u_2 = B'e^{u_2} - A'e^{u_1} - 4\pi\alpha\delta_0 & \text{in } \Omega, \\ u_1 = u_2 = g(x) & \text{on } \partial\Omega, \end{cases}$$

where $\alpha \geq 0$ and $0 \in \Omega$. Recall the definitions of M , D in (16) and in Theorem 1.11, respectively. By using the Green's function G_0 with pole at 0 as in (11) we may consider

$$(18) \quad \tilde{u}_i(x) = u(x) + 4\pi\alpha G_0(x)$$

which satisfies

$$\begin{cases} -\Delta \tilde{u}_1 = Ah(x)e^{\tilde{u}_1} - Bh(x)e^{\tilde{u}_2} & \text{in } \Omega, \\ -\Delta \tilde{u}_2 = B'h(x)e^{\tilde{u}_2} - A'h(x)e^{\tilde{u}_1} & \text{in } \Omega, \\ \tilde{u}_1 = \tilde{u}_2 = g(x) & \text{on } \partial\Omega, \end{cases}$$

with $h(x) = e^{-4\pi\alpha G_0(x)}$. We have the following result.

Theorem 1.13. Let (u_1, u_2) be a solution of (17) with $\alpha \geq 0$ and (16). Let \tilde{u}_i be as in (18). Suppose Ω is bounded, simply-connected and

$$\int_{\Omega} (e^{\tilde{u}_1} + e^{\tilde{u}_2}) dx \leq \frac{8\pi}{M}$$

Then $u_1 \equiv u_2 \equiv u$, where u is the unique solution to

$$\begin{cases} -\Delta u = De^u - 4\pi\alpha\delta_0 & \text{in } \Omega, \\ u = g(x) & \text{on } \partial\Omega. \end{cases}$$

The next remark concerns a possible generalization of the results we have obtained so far for multiply-connected domains.

Remark 1.14. *All the previous results hold for multiply-connected domains with constant boundary condition, i.e. $g(x) = c \in \mathbb{R}$. This follows from the same arguments and the Sphere Covering Inequality (Theorem 2.5) for multiply-connected domains. See Remark 2.6 below.*

The paper is organized as follows. In Section 2 we recall the main ingredients of the Sphere Covering Inequality. In Section 3 we present our strategy for proving the uniqueness result of Theorem 1.1, the symmetry result of Corollary 1.2, and the uniqueness result of Theorem 1.4. In Section 4 we show how to get the no intersection property of Theorem 1.8 and the symmetry property of Corollary 1.9. In Section 5 we provide the proof of the uniqueness result in Theorems 1.11 and 1.13.

Notation

The symbol $B_r(p)$ will denote the open metric ball of radius r and center p . Where there is no ambiguity, with a little abuse of notation we will write x and dx to denote $(x, y) \in \mathbb{R}^2$ and the integration with respect to (x, y) , respectively.

2. THE SPHERE COVERING INEQUALITY

In this section we recall the main ingredients of the Sphere Covering Inequality proved in [21] as we will need them in the sequel. Roughly speaking, the latter result asserts that the total area of two distinct surfaces with Gaussian curvature equal to 1, conformal to the Euclidean unit disk with the same conformal factor on the boundary, must cover the whole unit sphere after a proper rearrangement. See [21] for more details. Let us start by recalling the standard Bol's isoperimetric inequality as in [53, 55] (see also [7] in its original form).

Proposition 2.1. *Let $\Omega \subset \mathbb{R}^2$ be a simply-connected set and $u \in C^2(\Omega)$ be such that*

$$\Delta u + e^u \geq 0 \quad \text{and} \quad \int_{\Omega} e^u dx \leq 8\pi.$$

Then, for any $\omega \subset\subset \Omega$ of class C^1 it holds

$$\left(\int_{\partial\omega} e^{\frac{u}{2}} d\sigma \right)^2 \geq \frac{1}{2} \left(\int_{\omega} e^u dx \right) \left(8\pi - \int_{\omega} e^u dx \right).$$

The basic function, which satisfies the above properties and will be used in the sequel, is the following:

$$(19) \quad U_{\lambda}(x) = -2 \ln \left(1 + \frac{\lambda^2 |x|^2}{8} \right) + 2 \ln \lambda,$$

for $\lambda > 0$. Observe that

$$\Delta U_{\lambda} + e^{U_{\lambda}} = 0 \quad \text{and} \quad \int_{B_r(0)} e^{U_{\lambda}} dx = 8\pi \frac{\lambda^2 r^2}{8 + \lambda^2 r^2},$$

for all $r > 0$.

Now the idea is to consider symmetric rearrangements with respect to two distinct measures. More precisely, let $w \in C^2(\overline{\Omega})$ be such that

$$(20) \quad \Delta w + e^w \geq 0.$$

Then, any function $\phi \in C^2(\overline{\Omega})$ can be equimeasurably rearranged with respect to the measures $e^w dx$ and $e^{U_\lambda} dx$ (see [3]). Indeed, for $t > \min_{x \in \overline{\Omega}} \phi(x)$ let \mathcal{B}_t^* be the ball centered at the origin such that

$$\int_{\mathcal{B}_t^*} e^{U_\lambda} dx = \int_{\{\phi > t\}} e^w dx.$$

Then, if we let $\phi^* : \mathcal{B}_t^* \rightarrow \mathbb{R}$ to be $\phi^*(x) = \sup\{t \in \mathbb{R} : x \in \mathcal{B}_t^*\}$, it holds that ϕ^* is a symmetric equimeasurable rearrangement of ϕ with respect to the measures $e^w dx$ and $e^{U_\lambda} dx$, i.e.

$$(21) \quad \int_{\{\phi^* > t\}} e^{U_\lambda} dx = \int_{\{\phi > t\}} e^w dx,$$

for all $t > \min_{x \in \overline{\Omega}} \phi(x)$. Moreover, by using Bol's inequality stated in Proposition 2.1 we get the following estimate on the gradient of the rearrangement (see [21]).

Proposition 2.2. *Let $w \in C^2(\overline{\Omega})$ be such that it satisfies (20) with $\Omega \subset \mathbb{R}^2$ being simply-connected. Let U_λ be as in (19). Suppose $\phi \in C^2(\overline{\Omega})$ is such that $\phi \equiv C$ on $\partial\Omega$. If ϕ^* is the equimeasurable symmetric rearrangement of ϕ with respect to the measures $e^w dx$ and $e^{U_\lambda} dx$, then*

$$\int_{\{\phi^* = t\}} |\nabla \phi^*| d\sigma \leq \int_{\{\phi = t\}} |\nabla \phi| d\sigma,$$

for all $t > \min_{x \in \overline{\Omega}} \phi(x)$.

We shall also need the following counterpart of Bol's inequality in the radial setting (see [21]).

Proposition 2.3. *Let $\psi \in C^{0,1}(\overline{B_R(0)})$ be a strictly decreasing radial function satisfying*

$$\int_{\partial B_r(0)} |\nabla \psi| d\sigma \leq \int_{B_r(0)} e^\psi dx \quad \text{for a.e. } r \in (0, R) \quad \text{and} \quad \int_{B_R(0)} e^\psi dx \leq 8\pi.$$

Then

$$\left(\int_{\partial B_R(0)} e^{\frac{\psi}{2}} d\sigma \right)^2 \geq \frac{1}{2} \left(\int_{B_R(0)} e^\psi dx \right) \left(8\pi - \int_{B_R(0)} e^\psi dx \right).$$

The main idea is then to relate the strictly decreasing radial function ψ with two radial solutions $U_{\lambda_1}, U_{\lambda_2}$ defined in (19) with $\lambda_2 > \lambda_1$, such that $\psi = U_{\lambda_1} = U_{\lambda_2}$ on $\partial B_R(0)$.

Proposition 2.4. *$U_{\lambda_1}, U_{\lambda_2}$ defined in (19) with $\lambda_2 > \lambda_1$. Let $\psi \in C^{0,1}(\overline{B_R(0)})$ be a strictly decreasing radial function satisfying*

$$(22) \quad \int_{\partial B_r(0)} |\nabla \psi| d\sigma \leq \int_{B_r(0)} e^\psi dx \quad \text{for a.e. } r \in (0, R)$$

and $\psi = U_{\lambda_1} = U_{\lambda_2}$ on $\partial B_R(0)$. Then, either

$$\int_{B_R(0)} e^\psi dx \leq \int_{B_R(0)} e^{U_{\lambda_1}} dx \quad \text{or} \quad \int_{B_R(0)} e^\psi dx \geq \int_{B_R(0)} e^{U_{\lambda_2}} dx.$$

Moreover, we have

$$\int_{B_R(0)} (e^{U_{\lambda_1}} + e^{U_{\lambda_2}}) dx = 8\pi.$$

We can now state the Sphere Covering Inequality as in [21].

Theorem 2.5. *Let $\Omega \subset \mathbb{R}^2$ be a simply-connected set and let $w_i \in C^2(\overline{\Omega})$, $i = 1, 2$ be such that*

$$(23) \quad \Delta w_i + e^{w_i} = f_i(x) \quad \text{in } \Omega,$$

where $f_2 \geq f_1 \geq 0$ in Ω . Suppose

$$\begin{cases} w_2 \geq w_1, w_2 \neq w_1 & \text{in } \Omega, \\ w_2 = w_1 & \text{on } \partial\Omega, \end{cases}$$

Then, it holds

$$\int_{\Omega} (e^{w_1} + e^{w_2}) dx \geq 8\pi.$$

Moreover, if some $f_i \neq 0$ then the latter inequality is strict.

The idea is to consider a symmetric rearrangement φ of $w_2 - w_1$ with respect to the measures $e^{w_1} dx$ and $e^{U_{\lambda_1}} dx$ for a suitable λ_2 . Then, by using equation (23) and the properties of the rearrangements (see also Proposition 2.2), it is possible to show that (22) holds true for $\psi = U_{\lambda_1} + \varphi$. Applying then Proposition 2.4 one can deduce that

$$\int_{\Omega} (e^{w_1} + e^{w_2}) dx \geq \int_{B_R(0)} (e^{U_{\lambda_1}} + e^{U_{\lambda_2}}) dx = 8\pi.$$

See [21] for full details.

Remark 2.6. *We point out that the Sphere Covering Inequality holds as long as Bol's inequality holds. Indeed, if $\Delta w + e^w \geq 0$ in Ω which is simply-connected, then Bol's and Sphere Covering Inequalities hold in any region $\Omega_1 \subset \Omega$ for general boundary data. In particular, Ω_1 does not need to be simply-connected. Moreover, Bol's inequality and Sphere Covering inequalities hold for a multiply-connected domain Ω , provided that we have constant boundary conditions (see [5]).*

3. ASYMMETRIC SINH-GORDON EQUATION

In this section we study uniqueness and symmetry of solutions of asymmetric Sinh-Gordon equation (1), and prove Theorem 1.1 and Theorem 1.4. The first one relies mainly on the Sphere Covering Inequality (see Theorem 2.5). On the other hand, the second one is based on the arguments which yield the Sphere Covering Inequality, which we collected in Section 2.

Let us start with the case $\text{supp } \mathcal{P} \subset [0, 1]$ which we recall here for convenience

$$(24) \quad \begin{cases} -\Delta u = \rho \frac{e^u + e^{au}}{\int_{\Omega} (e^u + e^{au}) dx} & \text{in } \Omega, \\ u = g(x) \geq 0 & \text{on } \partial\Omega, \end{cases}$$

with $a \in (0, 1)$, $\rho > 0$, and $g \in C(\partial\Omega)$.

Proof of Theorem 1.1. Let u_1 and u_2 be solutions of equation (24) satisfying the assumptions of Theorem (1.1). We aim to show that $u_1 \equiv u_2$. We proceed by contradiction by assuming that this is not the case. Rewrite equation (24) as

$$\Delta u + \rho \frac{2e^u}{\int_{\Omega} (e^u + e^{au}) dx} = \rho \frac{e^u - e^{au}}{\int_{\Omega} (e^u + e^{au}) dx}.$$

Let

$$(25) \quad v = u + \log 2 + \log \rho - \log \left(\int_{\Omega} (e^u + e^{au}) dx \right).$$

Then v satisfies

$$(26) \quad \Delta v + e^v = f(u) := \rho \frac{e^u - e^{au}}{\int_{\Omega} (e^u + e^{au}) dx}.$$

It follows from (6) that there exists two regions $\Omega_1, \Omega_2 \subset \Omega$ (not necessarily simply-connected) such that $u_1 > u_2$ in Ω_1 , $u_2 > u_1$ in Ω_2 , and $u_1 = u_2$ on $\partial\Omega_1 \cup \partial\Omega_2$. We have that v_1, v_2 defined by (25) satisfy

$$\Delta v_i + e^{v_i} = f(u_i) \quad \text{in } \Omega.$$

Moreover

$$v_1 > v_2 \quad \text{in } \Omega_1, \quad v_2 > v_1 \quad \text{in } \Omega_2 \quad \text{and} \quad v_1 = v_2 \quad \text{on } \partial\Omega_1 \cup \partial\Omega_2.$$

Since $g \geq 0$, both solutions u_1 and u_2 are positive in Ω by the maximum principle. By the latter fact it is also easy to see that

$$f(u_1) > f(u_2) > 0 \quad \text{in } \Omega_1 \quad \text{and} \quad f(u_2) > f(u_1) > 0 \quad \text{in } \Omega_2.$$

Therefore, by applying the Sphere Covering Inequality (Theorem 2.5, see also Remark 2.6), we get (observe that $f_i \neq 0$)

$$\int_{\Omega} (e^{v_1} + e^{v_2}) dx \geq \int_{\Omega_1} (e^{v_1} + e^{v_2}) dx + \int_{\Omega_2} (e^{v_1} + e^{v_2}) dx > 16\pi.$$

Recalling now the definition of v in (25) and (6) we have

$$\begin{aligned} 4\rho &= \frac{2\rho}{\int_{\Omega} (e^{u_1} + e^{au_1}) dx} \left(\int_{\Omega} (e^{u_1} + e^{au_1}) dx + \int_{\Omega} (e^{u_1} + e^{au_1}) dx \right) \\ &\geq \frac{2\rho}{\int_{\Omega} (e^{u_1} + e^{au_1}) dx} \int_{\Omega} (e^{u_1} + e^{u_2}) dx = \int_{\Omega} (e^{v_1} + e^{v_2}) dx > 16\pi. \end{aligned}$$

Hence $\rho > 4\pi$, which is a contradiction. The proof is now complete. \square

Proof of Corollary 1.2. Without loss of generality we can assume that Ω and g are evenly symmetric with respect to the line $y = 0$. Suppose u is a solution of (5), which is not evenly symmetric about $y = 0$. Then $u_1 = u$ and $u_2(x, y) = u(x, -y)$ are two distinct solutions of (5) satisfying the condition (6). Thus it follows from Theorem 1.1 that $\rho > 4\pi$. \square

We consider now the general case $\text{supp } \mathcal{P} \subset [-1, 1]$ which yields to (7), i.e.:

$$(27) \quad \begin{cases} -\Delta u = \rho \frac{e^u - e^{-au}}{\int_{\Omega} (e^u + e^{-au}) dx} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

with $a \in (0, 1)$, $\rho > 0$. We give here the proof of the uniqueness result for the trivial solution $u \equiv 0$.

Proof of Theorem 1.4. Let u be a solution of (27). We will show that $u \equiv 0$ in Ω . Assume by contradiction this is not the case and let

$$(28) \quad \begin{aligned} v_1 &= -au + \log \rho - \log \left(\int_{\Omega} (e^u + e^{-au}) dx \right), \\ v_2 &= u + \log \rho - \log \left(\int_{\Omega} (e^u + e^{-au}) dx \right). \end{aligned}$$

Then we have

$$\Delta(v_2 - v_1) + (1 + a)(e^{v_2} - e^{v_1}) = 0.$$

Letting further

$$(29) \quad w_i = v_i + \log(1 + a), \quad i = 1, 2,$$

we deduce

$$(30) \quad \Delta(w_2 - w_1) + (e^{w_2} - e^{w_1}) = 0.$$

Since $u = 0$ on $\partial\Omega$, we get

$$(31) \quad w_1 = w_2 = \log(1 + a) + \log \rho - \log \left(\int_{\Omega} (e^u + e^{-au}) dx \right) \quad \text{on } \partial\Omega.$$

It follows that there exists at least one region $\tilde{\Omega} \subseteq \Omega$ (not necessarily simply-connected) such that

$$(32) \quad \begin{cases} w_1 \neq w_2 & \text{in } \tilde{\Omega}, \\ w_1 = w_2 & \text{on } \partial\tilde{\Omega}, \end{cases}$$

and

$$(33) \quad \Delta(w_2 - w_1) + (e^{w_2} - e^{w_1}) = 0 \quad \text{in } \tilde{\Omega}.$$

We point out that $\tilde{\Omega}$ may coincide with Ω . Without loss of generality we may assume $w_2 > w_1$. From equation (27) and the definitions of w_i in (28) and (29) we derive that

$$\Delta v_1 + ae^{v_1} = ae^{v_2}$$

and thus

$$(34) \quad \Delta w_1 + e^{w_1} = \left(\frac{1}{1+a} e^{w_1} + ae^{v_2} \right) > 0 \quad \text{in } \Omega.$$

We now proceed as in the proof of the Sphere Covering Inequality. Let $\lambda_2 > \lambda_1$ be such that $U_{\lambda_2} > U_{\lambda_1}$ in $B_1(0)$ and $U_{\lambda_1} = U_{\lambda_2}$ on $\partial B_1(0)$, where U_{λ} is given as in (19), and such that

$$\int_{\tilde{\Omega}} e^{w_1} dx = \int_{B_1(0)} e^{U_{\lambda_1}} dx.$$

Since w_1 satisfies (34) we can find a symmetric equimeasurable rearrangement φ^* of $w_2 - w_1$ with respect to the two measures $e^{w_1} dx$ and $e^{U_{\lambda_1}} dx$. See the discussion after (20). In particular we have

$$\int_{\{\varphi^* > t\}} e^{U_{\lambda_1}} dx = \int_{\{w_2 - w_1 > t\}} e^{w_1} dx$$

for $t \geq 0$. We first estimate the gradient of the rearrangement by Proposition 2.2, then exploit equation (33), the equation satisfied by U_{λ_1} and the properties of the rearrangements to obtain

$$\begin{aligned} \int_{\{\varphi^*=t\}} |\nabla \varphi^*| d\sigma &\leq \int_{\{w_2-w_1=t\}} |\nabla(w_2 - w_1)| d\sigma \\ &= \int_{\{w_2-w_1>t\}} (e^{w_2} - e^{w_1}) dx \\ &= \int_{\{\varphi^*>t\}} e^{U_{\lambda_1}+\varphi^*} dx - \int_{\{\varphi^*>t\}} e^{U_{\lambda_1}} dx \\ &= \int_{\{\varphi^*>t\}} e^{U_{\lambda_1}+\varphi^*} dx - \int_{\{\varphi^*=t\}} |\nabla U_{\lambda_1}| d\sigma, \end{aligned}$$

for a.e. $t > 0$. Therefore

$$\int_{\{\varphi^*=t\}} |\nabla(U_{\lambda_1} + \varphi^*)| d\sigma \leq \int_{\{\varphi^*>t\}} e^{U_{\lambda_1}+\varphi^*} dx,$$

for a.e. $t > 0$. Since φ^* is decreasing by construction, $U_{\lambda_1} + \varphi^*$ is a strictly decreasing function. Moreover, by the above estimate we derive

$$(35) \quad \int_{\partial B_r(0)} |\nabla(U_{\lambda_1} + \varphi^*)| d\sigma \leq \int_{B_r(0)} e^{U_{\lambda_1}+\varphi^*} dx \quad \text{for a.e. } r > 0.$$

Furthermore, since $\varphi^* \geq 0$, we clearly have

$$\int_{B_1(0)} e^{U_{\lambda_1}+\varphi^*} dx \geq \int_{B_1(0)} e^{U_{\lambda_1}} dx.$$

By the latter estimate, (34) and (35) we can exploit Proposition 2.4 with $\psi = U_{\lambda_1} + \varphi^*$ to get

$$\int_{B_1(0)} e^{U_{\lambda_1}+\varphi^*} dx \geq \int_{B_1(0)} e^{U_{\lambda_2}} dx.$$

Thus

$$\int_{\tilde{\Omega}} (e^{w_1} + e^{w_2}) dx = \int_{B_1(0)} (e^{U_{\lambda_1}} + e^{U_{\lambda_1}+\varphi^*}) dx \geq \int_{B_1(0)} (e^{U_{\lambda_1}} + e^{U_{\lambda_2}}) dx = 8\pi.$$

Recall now the definitions of w_i in (28) and (29). We have

$$\frac{\rho(1+a)}{\int_{\Omega} (e^u + e^{-au}) dx} \int_{\tilde{\Omega}} (e^u + e^{-au}) dx \geq 8\pi,$$

and hence

$$\frac{8\pi}{1+a} \leq \frac{\rho}{\int_{\Omega} (e^u + e^{-au}) dx} \int_{\tilde{\Omega}} (e^u + e^{-au}) dx \leq \rho.$$

The above inequality is indeed strict. To see this, we note that the equality would yield the equality in (35) which corresponds to equality in Bol's inequality in Proposition 2.1 for w_1 and consequently w_1 should satisfy $\Delta w_1 + e^{w_1} = 0$, which contradicts (34). In view of the assumption $\rho \leq \frac{8\pi}{1+a}$, we therefore have shown $u \equiv 0$ in Ω as desired. \square

4. COSMIC STRING EQUATION

In this section we study the cosmic string equation

$$(36) \quad \begin{cases} -\Delta u = e^{au} + h(x)e^u & \text{in } \Omega, \\ u = g(x) \geq 0 & \text{on } \partial\Omega, \end{cases}$$

with $a > 0$ and h as in (10). We will rewrite this equation in a form suitable to apply the Sphere Covering Inequality, Theorem 2.5, to prove Theorem 1.8.

Proof of Theorem 1.8. First suppose $a > 1$. Let u_1 and u_2 be two solutions of (36) with $a > 1$, $N \geq 0$ satisfying (12). We proceed by contradiction. Suppose there exists $\Omega_1, \Omega_2 \subset \Omega$ (not necessarily simply-connected) such that

$$u_1 > u_2 \text{ in } \Omega_1 \quad \text{and} \quad u_2 > u_1 \text{ in } \Omega_2.$$

The equation (36) can be rewritten as

$$\Delta u + 2e^{au} = e^{au} - h(x)e^u.$$

Multiply this equation by a and let

$$(37) \quad v = au + \log(2a).$$

Then v satisfies

$$(38) \quad \Delta v + e^v = f(u) := a(e^{au} - h(x)e^u).$$

Let v_1, v_2 be defined by (37) (u replaced by u_1 and u_2 , respectively). Then we have

$$\Delta v_i + e^{v_i} = f(u_i) \text{ in } \Omega.$$

Furthermore, we get

$$v_1 > v_2 \text{ in } \Omega_1, \quad v_2 > v_1 \text{ in } \Omega_2 \quad \text{and} \quad v_1 = v_2 \text{ on } \partial\Omega_1 \cup \partial\Omega_2.$$

Since $g \geq 0$, it follows from the maximum principle that both solutions u_1 and u_2 are positive inside Ω . Note also that $h(x) \leq 1$. It is now easy to see that

$$f(u_1) > f(u_2) > 0 \text{ in } \Omega_1 \quad \text{and} \quad f(u_2) > f(u_1) > 0 \text{ in } \Omega_2.$$

By the Sphere Covering Inequality (Theorem 2.5, see also Remark 2.6) we conclude that

$$\int_{\Omega} (e^{v_1} + e^{v_2}) dx \geq \int_{\Omega_1} (e^{v_1} + e^{v_2}) dx + \int_{\Omega_2} (e^{v_1} + e^{v_2}) dx > 16\pi.$$

Using the expression of v in (37) we deduce

$$2a \int_{\Omega} (e^{au_1} + e^{au_2}) dx > 16\pi,$$

which contradicts the assumption

$$\int_{\Omega} (e^{au_1} + e^{au_2}) dx \leq \frac{8\pi}{a}.$$

For what concerns the case $a < 1$ we write (36) in the form

$$\Delta u + 2e^u = (e^u - e^{au}) + (e^u - h(x)e^u).$$

The argument is then developed as before so we skip the details. The proof is now complete. \square

Proof of Corollary 1.9. Without loss of generality we assume that Ω and g are evenly symmetric with respect to the line $y = 0$. Observe that the associated Green's function (and hence h , see (10)) is evenly symmetric with respect to the line $y = 0$. We consider just the case $a > 1$ since for $a < 1$ one can proceed in the same way. Suppose u is a solution of (5) satisfying (14), which is not evenly symmetric about $y = 0$. Then $u_1 = u$ and $u_2(x, y) = u(x, -y)$ are two distinct intersecting solutions of (9). It follows from Theorem 1.8 that

$$2 \int_{\Omega} e^{au} dx = \int_{\Omega} (e^{au_1} + e^{au_2}) dx > \frac{8\pi}{a}.$$

which contradicts (14). \square

5. LIOUVILLE-TYPE SYSTEMS IN DOMAINS

In this section we consider the class of Liouville-type systems

$$(39) \quad \begin{cases} -\Delta u_1 = Ae^{u_1} - Be^{u_2} \\ -\Delta u_2 = B'e^{u_2} - A'e^{u_1} \\ u_1 = u_2 = g(x) \end{cases} \quad \begin{array}{l} \text{in } \Omega, \\ \text{in } \Omega, \\ \text{on } \partial\Omega, \end{array}$$

where A, A', B, B' satisfy condition (16), and prove Theorem 1.11.

Proof of Theorem 1.11. Let (u_1, u_2) be a solution of (39). We will prove that there exists a unique u solving a mean field equation as stated in Theorem 1.11 such that $u_1 \equiv u_2 \equiv u$ in Ω . Assume by contradiction $u_1 \not\equiv u_2$. As in the proof of Theorem 1.4, the strategy is to apply the argument of the Sphere Covering Inequality in Theorem 2.5 (see Section 2) to the functions u_1 and u_2 . We start by recalling that the coefficients in (39) are such that $A + A' = B + B' := M$. Hence

$$\Delta(u_2 - u_1) + M(e^{u_2} - e^{u_1}) = 0.$$

Letting

$$(40) \quad w_i = u_i + \log M, \quad i = 1, 2,$$

we deduce that

$$(41) \quad \Delta(w_2 - w_1) + (e^{w_2} - e^{w_1}) = 0,$$

and

$$(42) \quad w_1 = w_2 = \log M + g(x) \quad \text{on } \partial\Omega.$$

It follows that there exists at least one region $\tilde{\Omega} \subseteq \Omega$ (not necessarily simply-connected) such that

$$(43) \quad \begin{cases} w_1 \neq w_2 & \text{in } \tilde{\Omega}, \\ w_1 = w_2 & \text{on } \partial\tilde{\Omega}, \end{cases}$$

and

$$(44) \quad \Delta(w_2 - w_1) + (e^{w_2} - e^{w_1}) = 0 \quad \text{in } \Omega.$$

Without loss of generality we can assume $w_2 > w_1$ in $\tilde{\Omega}$.

Using the first equation in (39), the definitions of w_i in (29), and the fact that $M = A + A'$ we get

$$\Delta u_1 + Ae^{u_1} = Be^{u_2}$$

and hence

$$(45) \quad \Delta w_1 + e^{w_1} = \left(\frac{A'}{A + A'} e^{w_1} + B e^{u_2} \right) \geq 0 \quad \text{in } \Omega.$$

The rest of the argument is very similar to the proof of Theorem 1.4 so we will skip the details. Let $\lambda_2 > \lambda_1$ be such that $U_{\lambda_2} > U_{\lambda_1}$ in $B_1(0)$ and $U_{\lambda_1} = U_{\lambda_2}$ on $\partial B_1(0)$, where U_λ is given as in (19), and

$$\int_{\tilde{\Omega}} e^{w_1} dx = \int_{B_1(0)} e^{U_{\lambda_1}} dx.$$

Recalling (45) we can find a symmetric equimeasurable rearrangement φ^* of $w_2 - w_1$ with respect to the two measures $e^{w_1} dx$ and $e^{U_{\lambda_1}} dx$. Reasoning as in the proof of Theorem 1.4 we get

$$\int_{\partial B_r(0)} |\nabla(U_{\lambda_1} + \varphi^*)| d\sigma \leq \int_{B_r(0)} e^{U_{\lambda_1} + \varphi^*} dx \quad \text{for a.e. } r > 0.$$

Furthermore $U_{\lambda_1} + \varphi^*$ is a strictly decreasing function. Hence from Proposition 2.4 to $\psi = U_{\lambda_1} + \varphi^*$ we deduce

$$\int_{B_1(0)} e^{U_{\lambda_1} + \varphi^*} dx \geq \int_{B_1(0)} e^{U_{\lambda_2}} dx.$$

Therefore

$$\int_{\tilde{\Omega}} (e^{w_1} + e^{w_2}) dx = \int_{B_1(0)} (e^{U_{\lambda_1}} + e^{U_{\lambda_1} + \varphi^*}) dx \geq \int_{B_1(0)} (e^{U_{\lambda_1}} + e^{U_{\lambda_2}}) dx = 8\pi.$$

It follows from the definitions of w_i that

$$M \int_{\tilde{\Omega}} (e^{u_1} + e^{u_2}) dx \geq 8\pi.$$

Thus

$$\frac{8\pi}{M} \leq \int_{\tilde{\Omega}} (e^{u_1} + e^{u_2}) dx \leq \int_{\Omega} (e^{u_1} + e^{u_2}) dx.$$

Arguing as in the proof of Theorem 1.4 it is easy to show that the latter inequality is strict, which is a contradiction. Hence $u_1 \equiv u_2$ in Ω . Letting $u := u_1 = u_2$ and using the system (39) we get

$$\begin{cases} -\Delta u = D e^u & \text{in } \Omega, \\ u = g(x) & \text{on } \partial\Omega, \end{cases}$$

where we recall $D := A - B = A' - B'$. Note that $M := A + A' = B + B'$ and hence

$$\int_{\Omega} D e^u dx \leq 4\pi \frac{D}{M} = 4\pi \frac{A - B}{A + A'} < 4\pi.$$

Since Ω is simply-connected and the latter bound holds true, by the Sphere Covering Inequality of Theorem 2.5 we deduce that u is unique. This concludes the proof of Theorem 1.11. \square

We conclude this section by giving the proof of Theorem 1.13 regarding the uniqueness of solutions of the system

$$(46) \quad \begin{cases} -\Delta u_1 = Ae^{u_1} - Be^{u_2} - 4\pi\alpha\delta_0 & \text{in } \Omega, \\ -\Delta u_2 = B'e^{u_2} - A'e^{u_1} - 4\pi\alpha\delta_0 & \text{in } \Omega, \\ u_1 = u_2 = g(x) & \text{on } \partial\Omega. \end{cases}$$

Proof of Theorem 1.13. Let (u_1, u_2) be a solution of (46) with $\alpha \geq 0$. By using the Green's function G_0 with pole in 0 as in (11) we desingularize the problem by setting

$$\tilde{u}_i(x) = u(x) + 4\pi\alpha G_0(x).$$

Indeed (46) is equivalent to

$$(47) \quad \begin{cases} -\Delta \tilde{u}_1 = Ah(x)e^{\tilde{u}_1} - Bh(x)e^{\tilde{u}_2} & \text{in } \Omega, \\ -\Delta \tilde{u}_2 = B'h(x)e^{\tilde{u}_2} - A'h(x)e^{\tilde{u}_1} & \text{in } \Omega, \\ \tilde{u}_1 = \tilde{u}_2 = g(x) & \text{on } \partial\Omega, \end{cases}$$

where

$$(48) \quad h(x) = e^{-4\pi\alpha G_0(x)}.$$

Observe that

$$h > 0 \quad \text{in } \Omega \setminus \{0\} \quad \text{and} \quad h(x) \cong |x|^{2\alpha} \quad \text{near } 0.$$

Assume now by contradiction that $\tilde{u}_1 \not\equiv \tilde{u}_2$ and suppose, without loss of generality, that $\tilde{u}_2 > \tilde{u}_1$ in $\tilde{\Omega} \subseteq \Omega$. Recall that $A + A' = B + B' := M$. Therefore, by (47) we have

$$\Delta(\tilde{u}_2 - \tilde{u}_1) + Mh(x)(e^{\tilde{u}_2} - e^{\tilde{u}_1}) = 0.$$

Note also that $h(x) \leq 1$. Since $\tilde{u}_2 > \tilde{u}_1$ in $\tilde{\Omega}$ we deduce

$$\Delta(\tilde{u}_2 - \tilde{u}_1) + M(e^{\tilde{u}_2} - e^{\tilde{u}_1}) \geq 0 \quad \text{in } \tilde{\Omega}.$$

With an argument similar to the one in the proof of Theorem 1.11 we get a contradiction. Thus $\tilde{u}_1 \equiv \tilde{u}_2 := \tilde{u}$ and \tilde{u} satisfies

$$\begin{cases} -\Delta \tilde{u} = Dh(x)e^{\tilde{u}} & \text{in } \Omega, \\ \tilde{u} = g(x) & \text{on } \partial\Omega, \end{cases}$$

where $D := A - B = A' - B'$. Arguing as in the proof of Theorem 1.11 we deduce that \tilde{u} is unique. \square

Acknowledgements

The authors would like to thank Prof. G. Tarantello and Dr. W. Yang for the discussions concerning the topic of this paper.

REFERENCES

- [1] W. Ao, A. Jevnikar and W. Yang, *On the boundary behavior for the blow up solutions of the sinh-Gordon equation and B_2, G_2 Toda systems in bounded domain*. Preprint. <http://cvgmt.sns.it/paper/3215/>.
- [2] T. Aubin, *Nonlinear analysis on manifolds. Monge-Ampere equations*, Springer-Verlag, Berlin, 1982.
- [3] C. Bandle. *Isoperimetric inequalities and applications*. Volume 7 of Monographs and Studies in Mathematics. Pitman (Advanced Publishing Program), Boston, Mass. London, 1980.
- [4] D. Bartolucci and C.S. Lin, *Uniqueness results for mean field equations with singular data*, *Comm. in Partial Differential Equations* 34 (2009), 676-702.
- [5] D. Bartolucci and C.S. Lin, *Existence and uniqueness for Mean Field Equations on multiply connected domains at the critical parameter*, *Math. Ann.* 359 (2014), 1-44.
- [6] L. Battaglia, A. Jevnikar, A. Malchiodi and D. Ruiz, A general existence result for the Toda system on compact surfaces. *Adv. Math.* 285 (2015), 937-979.
- [7] G. Bol, Isoperimetrische Ungleichungen für Bereiche auf Flächen. *Jber. Deutsch. Math. Verein.* 51 (1941), 219-257.
- [8] J. Bolton and L.M. Woodward, Some geometrical aspects of the 2-dimensional Toda equations. In *Geometry, topology and physics* (Campinas, 1996), pages 69-81. de Gruyter, Berlin, 1997.
- [9] E. Caglioti, P.L. Lions, C. Marchioro and M. Pulvirenti, A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description. *Comm. Math. Phys.* 143 (1992), no. 3, 501-525.
- [10] E. Calabi, Isometric imbedding of complex manifolds. *Ann. of Math. (2)*, 58 (1953), 1-23.
- [11] D. Chae, Existence of a semilinear elliptic system with exponential nonlinearities, *Discrete Contin. Dyn. Syst.* 18 (2007), 709-718.
- [12] D. Chae, Existence of multistring solutions of a selfgravitating massive Wboson. *Lett. Math. Phys.* 73 (2005), 123-134.
- [13] Sun-Yung A. Chang and Paul C. Yang, Prescribing Gaussian curvature on S^2 , *Acta Math.*, 159(3-4) (1987), 215-179.
- [14] Sun-Yung A. Chang and Paul C. Yang. Conformal deformation of metrics on S^2 , *J. Differential Geom.* 27(2) (1988), 259-176.
- [15] W. Chen and C. Li, Classification of solutions of some nonlinear elliptic equations. *Duke Math. J.* 63 (1991), no. 3, 615-622.
- [16] W. Chen and C. Li, Qualitative properties of solutions to some nonlinear elliptic equations in R^2 , *Duke Math. J.* 71 (1993), 427-439.
- [17] K.S. Cheng and C.S. Lin, On the asymptotic behavior of solutions of the conformal Gaussian curvature equations in R^2 , *Math. Ann.* 308 (1997), 119-139.
- [18] F. De Marchis and T. Ricciardi, Existence of stationary turbulent flows with variable positive vortex intensity. Preprint. Arxiv <https://arxiv.org/abs/1607.07051>.
- [19] G. Dunne, *Self-dual Chern-Simons Theories*. Lecture notes in physics. New series m: Monographs. Springer, 1995.
- [20] N. Ghoussoub and C.S. Lin, On the best constant in the Moser-Onofri-Aubin inequality, *Comm. Math. Phys.* 298 (2010), no. 3, 869-878.
- [21] C. Gui and A. Moradifam, The Sphere Covering Inequality and its applications. Preprint. Arxiv <https://arxiv.org/abs/1605.06481>.
- [22] C. Gui and A. Moradifam, Symmetry of solutions of a mean field equation on flat tori. *International Mathematics Research Notices*, to appear. Arxiv <https://arxiv.org/abs/1605.06905>.
- [23] C. Gui and A. Moradifam, Uniqueness of solutions of mean field equations in R^2 , *Proc. AMS* to appear. Arxiv <https://arxiv.org/pdf/1612.08403.pdf>.
- [24] A. Jevnikar, An existence result for the mean field equation on compact surfaces in a doubly supercritical regime. *Proc. Royal Soc. Edimb. Sect. A* 143 (2013), no. 5, 1021-1045.
- [25] A. Jevnikar, New existence results for the mean field equation on compact surfaces via degree theory. *Rend. Semin. Mat. Univ. Padova* 136 (2016), 11-17.
- [26] A. Jevnikar, A note on a multiplicity result for the mean field equation on compact surfaces. *Adv. Nonlinear Stud.* 16 (2016), no. 2, 221-229.
- [27] A. Jevnikar, Blow-up analysis and existence results in the supercritical case for an asymmetric mean field equation with variable intensities. *J. Diff. Eq.* 263 (2017), 972-1008.

- [28] A. Jevnikar, S. Kallel and A. Malchiodi, A topological join construction and the Toda system on compact surfaces of arbitrary genus. *Anal. PDE* 8 (2015), no. 8, 1963-2027.
- [29] A. Jevnikar, J. Wei and W. Yang, Classification of blow-up limits for the sinh-Gordon equation. To appear in *Differential and Integral Equations*.
- [30] A. Jevnikar, J. Wei and W. Yang, On the Topological degree of the Mean field equation with two parameters. To appear in *Indiana Univ. Math. J.*
- [31] A. Jevnikar and W. Yang, Analytic aspects of the Tzitzéica equation: blow-up analysis and existence results. *Calc. Var. and PDEs* 56 (2017), no. 2, 56:43.
- [32] A. Jevnikar and W. Yang, A mean field equation involving positively supported probability measures: blow-up phenomena and variational aspects. To appear in *Proc. Royal Soc. Edinb. Sect. A*.
- [33] J. Jost, C.S. Lin and G. Wang, Analytic aspects of the Toda system. II. Bubbling behavior and existence of solutions. *Comm. Pure Appl. Math.* 59 (2006), no. 4, 526-558.
- [34] J. Kazdan and F. Warner, Curvature functions for compact 2-manifolds, *Ann. of Math.* (99), (1974), 1417.
- [35] M.K.H. Kiessling, Statistical mechanics of classical particles with logarithmic interactions. *Comm. Pure Appl. Math.* 46 (1993), no. 1, 27-56.
- [36] C.S. Lin, Uniqueness of solutions to the mean field equations for the spherical Onsager vortex, *Arch. Ration. Mech. Anal.*, 153 (2000), no. 2, 153-176.
- [37] C.S. Lin, J.C. Wei and W. Yang, Degree counting and shadow system for SU(3) Toda system: one bubbling. Preprint. ArXiv <http://arxiv.org/pdf/1408.5802v1>.
- [38] C.S. Lin, J.C. Wei, W. Yang and L. Zhang, On Rank Two Toda System with Arbitrary Singularities: Local Mass and New Estimates. Preprint. Arxiv <https://arxiv.org/pdf/1609.02772v1>.
- [39] C.S. Lin, J. C. Wei and D. Ye, Classification and nondegeneracy of SU(n+1) Toda system. *Invent. Math.* 190 (2012), no. 1, 169-207.
- [40] A. Malchiodi, Topological methods for an elliptic equation with exponential nonlinearities. *Discrete Contin. Dyn. Syst.* 21 (2008), no. 1, 277-294.
- [41] A. Malchiodi and D. Ruiz, A variational Analysis of the Toda System on Compact Surfaces. *Comm. Pure Appl. Math.* 66 (2013), no. 3, 332-371
- [42] C. Neri, Statistical mechanics of the N-point vortex system with random intensities on a bounded domain. *Ann Inst. H. Poincaré Anal. Non Linéaire* 21 (2004), 381-399.
- [43] L. Onsager, Statistical hydrodynamics. *Nuovo Cimento Suppl.* 6 (1949), 279-287.
- [44] A. Poliakovsky and G. Tarantello, On a planar Liouvilletype problem in the study of self-gravitating strings. *J. Diff. Equations* 252, (2012), no. 5, 3668-3693.
- [45] A. Poliakovsky and G. Tarantello, *On singular Liouville systems*. Analysis and Topology in Nonlinear Differential Equations, PNDLE 85 Birkhauser Basel, 2014.
- [46] A. Poliakovsky and G. Tarantello, On non-topological solutions for planar Liouville Systems of Toda-type. *Comm. Math. Phys.* 347 (2016), 223-270.
- [47] T. Ricciardi, R. Takahashi, G. Zecca and X. Zhang, On the existence and blow-up of solutions for a mean field equation with variable intensities. Preprint. Arxiv <http://arxiv.org/abs/1509.05204v1>.
- [48] T. Ricciardi and G. Zecca, Mass quantization and minimax solutions for Neri's mean field equation in 2D-turbulence, *J. Differential Equations* 260 (2016), 339-369.
- [49] T. Ricciardi and G. Zecca, Minimal blow-up masses and existence of solutions for an asymmetric sinh-Poisson equation. Preprint. Arxiv <http://arxiv.org/abs/1605.05895>.
- [50] T. Ricciardi, G. Zecca, Blow-up analysis for some mean field equations involving probability measures from statistical hydrodynamics. *Differential and Integral Equations* 25 (2012), no. 3-4 201-222.
- [51] K. Sawada, and T. Suzuki, Derivation of the equilibrium mean field equations of point vortex and vortex filament system. *Theoret. Appl. Mech. Japan* 56 (2008), 285-290.
- [52] K. Sawada, T. Suzuki and F. Takahashi, Mean field equation for equilibrium vortices with neutral orientation. *Nonlinear Analysis: TMA* 66 (2007), no. 2, 509-526.
- [53] T. Suzuki, Global analysis for a two-dimensional elliptic eigenvalue problem with the exponential nonlinearity. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 9. (1992), no. 4, 367-397.
- [54] G. Tarantello, Analytical, geometrical and topological aspects of a class of mean field equations on surfaces. *Discrete Contin. Dyn. Syst.* 28 (2010), no. 3, 931-973.
- [55] G. Tarantello, Blow up analysis for a cosmic strings equation, *J. Funct. Anal.* 272 (2017), no. 1, 255178

- [56] G. Tarantello, *Selfdual gauge field vortices: an analytical approach*. Progress in Nonlinear Differential Equations and their Applications, 72. Birkhäuser Boston Inc., Boston, MA, 2008.
- [57] Y. Yang, *Solitons in field theory and nonlinear analysis*, Springer Monographs in Mathematics, SpingerVerlag, New York (2001).

CHANGFENG GUI, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT SAN ANTONIO,
TEXAS, USA

E-mail address: `changfeng.gui@utsa.edu`

ALEKS JEVNIKAR, UNIVERSITY OF ROME 'TOR VERGATA', VIA DELLA RICERCA SCIENTIFICA 1,
00133 ROMA, ITALY

E-mail address: `jevnikar@mat.uniroma2.it`

AMIR MORADIFAM, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE,
CALIFORNIA, USA

E-mail address: `amirm@ucr.edu`