

Implication Zroupoids and Identities of Associative Type

Juan M. Cornejo and Hanamantagouda P. Sankappanavar

Abstract. An algebra $\mathbf{A} = \langle A, \rightarrow, 0 \rangle$, where \rightarrow is binary and 0 is a constant, is called an *implication zroupoid* (\mathcal{I} -zroupoid, for short) if \mathbf{A} satisfies the identities: $(x \rightarrow y) \rightarrow z \approx [(z' \rightarrow x) \rightarrow (y \rightarrow z)]'$ and $0'' \approx 0$, where $x' := x \rightarrow 0$, and \mathcal{I} denotes the variety of all \mathcal{I} -zroupoids. An \mathcal{I} -zroupoid is *symmetric* if it satisfies $x'' \approx x$ and $(x \rightarrow y) \rightarrow z \approx (y \rightarrow x) \rightarrow z$. The variety of symmetric \mathcal{I} -zroupoids is denoted by \mathcal{S} . An identity $p \approx q$, in the groupoid language $\langle \rightarrow \rangle$, is called an *identity of associative type of length 3* if p and q have exactly 3 (distinct) variables, say x, y, z , and are grouped according to one of the two ways of grouping: (1) $\star \rightarrow (\star \rightarrow \star)$ and (2) $(\star \rightarrow \star) \rightarrow \star$, where \star is a place holder for a variable. A subvariety of \mathcal{I} is said to be of *associative type of length 3*, if it is defined, relative to \mathcal{I} , by a single identity of associative type of length 3. In this paper we give a complete analysis of the mutual relationships of all subvarieties of \mathcal{I} of associative type of length 3. We prove, in our main theorem, that there are exactly 8 such subvarieties of \mathcal{I} that are distinct from each other and describe explicitly the poset formed by them under inclusion. As an application of the main theorem, we derive that there are three distinct subvarieties of the variety \mathcal{S} , each defined, relative to \mathcal{S} , by a single identity of associative type of length 3.

1 Introduction

In 1934, Bernstein gave a system of axioms for Boolean algebras in [3] using implication alone. Even though his system was not equational, it is not hard to see that one could easily convert it into an equational one by using an additional constant. In 2012, the second author extended this “modified Bernstein’s theorem” to De Morgan algebras in [25] by showing that the variety of De Morgan algebras, is term-equivalent to the variety \mathcal{DM} (defined below) whose defining axioms use only an implication \rightarrow and a constant 0 .

The primary role played by the identity (I): $(x \rightarrow y) \rightarrow z \approx [(z' \rightarrow x) \rightarrow$

The first author wishes to dedicate this work to the memory of Oscar Foresi, his second dad.

The second author wishes to dedicate this work to the memory of his sister, Paddawwa R. Hooli.

2000 Mathematics Subject Classification: 06D30, 08B15, 20N02, 03G10.

Keywords: implication zroupoid, variety, identity of associative type

$(y \rightarrow z)']'$, where $x' := x \rightarrow 0$, in the axiomatization of each of those new varieties motivated the second author to study the identity (I) in its own right and led him to introduce a new equational class of algebras called “implication zroupoids” in [25] (also called “implicator groupoids” in [7]).

An algebra $\mathbf{A} = \langle A, \rightarrow, 0 \rangle$, where \rightarrow is binary and 0 is a constant, is called a zroupoid. A zroupoid $\mathbf{A} = \langle A, \rightarrow, 0 \rangle$ is an implication zroupoid (\mathcal{I} -zroupoid, for short) if \mathbf{A} satisfies:

$$(I) \quad (x \rightarrow y) \rightarrow z \approx [(z' \rightarrow x) \rightarrow (y \rightarrow z)']', \text{ where } x' := x \rightarrow 0,$$

$$(I_0) \quad 0'' \approx 0.$$

\mathcal{I} denotes the variety of implication zroupoids. The varieties \mathcal{DM} and \mathcal{SL} are defined relative to \mathcal{I} , respectively, by the following identities:

$$(DM) \quad (x \rightarrow y) \rightarrow x \approx x \text{ (De Morgan Algebras);}$$

$$(SL) \quad x' \approx x \text{ and } x \rightarrow y \approx y \rightarrow x \text{ (semilattices with the least element 0).}$$

The variety \mathcal{BA} of Boolean algebras is defined relative to \mathcal{DM} by the following identity:

$$(BA) \quad x \rightarrow x \approx 0'.$$

The variety \mathcal{I} exhibits (see [25]) several interesting properties; for example, the identity $x''' \rightarrow y \approx x' \rightarrow y$ holds in \mathcal{I} ; in particular, \mathcal{I} satisfies $x'''' \approx x''$. Two of the subvarieties of \mathcal{I} that are of interest in this paper are: $\mathcal{I}_{2,0}$ and \mathcal{MC} which are defined relative to \mathcal{I} , respectively, by the following identities, where $x \wedge y := (x \rightarrow y)'$:

$$(I_{2,0}) \quad x'' \approx x;$$


$$(MC) \quad x \wedge y \approx y \wedge x.$$

The (still largely unexplored) lattice of subvarieties of \mathcal{I} seems to be fairly complex. In fact, Problem 6 of [25] calls for an investigation of the structure of the lattice of subvarieties of \mathcal{I} .

The papers [5], [6], [7], [8], [9], [10] and [11] have addressed further the above-mentioned problem, but still partially, by introducing several new subvarieties of \mathcal{I} and investigating relationships among them. The (currently known) size of the poset of subvarieties of \mathcal{I} is at least 30; but it is still unknown whether the lattice of subvarieties is finite or infinite. We conjecture that its cardinality is 2^ω .

Motivated by the fact that not all algebras in \mathcal{I} are associative with respect to the operation \rightarrow , the quest for finding more new subvarieties of \mathcal{I} led us naturally to consider the question as to whether generalizations of the associative law would yield some new subvarieties of \mathcal{I} and thereby reveal further insight into the structure of the lattice of subvarieties of \mathcal{I} . This quest led to the results in [9], [10] and this paper, which will show that this is indeed the case.

The poset of the (then) known varieties that appears in [8] is given below for the reader's convenience (for the definitions of the varieties in the picture, see [8]).



A look at the associative law would reveal the following characteristics:

- (1) Length of the left side term = length of the right side term = 3,
- (2) The number of distinct variables on the left = the number of distinct variables

on right = the number of occurrences of variables on either side,

- (3) The order of the variables on the left side is the same as the order of the variables on the right side,
- (4) The bracketings used in the left side term and in the right side term are different from each other.

One way to generalize the associative law is to relax somewhat the restrictions (1) and (2) by choosing m distinct variables and setting the length of the left term = that of right term = n , with $n \geq m$, and keeping (3) and (4). But then, for $n \geq 4$, there will be more than two possible bracketings. So, we order all possible bracketings and assign a number to each, called bracketing number. Such identities are called “weak associative identities of length n ”. For a precise definition and notation of weak associative identities, we refer the reader to [10] and the references therein.

A second way to generalize the associative law is to relax (3) and to keep (2), (4) and the first half of (1). So, we consider the laws of the form $p \approx q$ of length n such that (a) each of p and q contains the n (an integer ≥ 3) distinct variables, say, x_1, x_2, \dots, x_n , (b) p and q are terms obtained by distinct bracketings of permutations of the n variables. Let us call such laws as “identities of associative type of length n ”.

A third way to generalize the associative law is to relax all of four features mentioned above by allowing number of occurrences of variables on one side be different from the number on the other side. Let us refer to these as “identities of mixed type”.

Specific instances of all such generalizations of the associative law have already occurred in the literature at least since late 19th century. We mention below a few such instances.

Weak associative identities of length 4 with 3 distinct variables, called “identities of Bol-Moufang type”, have been investigated in the literature quite extensively for the varieties of quasigroups and loops. In fact, the first systematic analysis of the relationships among the identities of Bol-Moufang type appears to be in [12] in the context of loops. For more information about these identities in the context of quasigroups and loops, see [12], [15], [20], [21]) and the references therein.

More recently, in [9] and [10], we **have** made a complete analysis of relationships among weak associative identities of length ≤ 4 , relative to the variety \mathcal{S} of symmetric I-zroupoids (i.e., satisfying $x'' \approx x$ and $(x \rightarrow y)' \approx (y \rightarrow x)'$). We have shown that 6 of the 155 subvarieties of \mathcal{S} , **each being defined by a single weak associative law of length $m \leq 4$ (including the Bol-Moufang type)**, are distinct. Furthermore, we describe explicitly by a Hasse diagram the poset formed by them, together with the varieties \mathcal{BA} and \mathcal{SL} .

We should mention here that such an analysis of weak associative laws of length ≤ 4 relative to the variety \mathcal{I} is still open.

The identities of associative type have also appeared in the literature. We mention several examples below, using \cdot for the binary operation instead of \rightarrow .

- The identity $x \cdot (y \cdot z) \approx (z \cdot x) \cdot y$ was considered in [30] by Suschkewitsch (see also [29, Theorem 11.5]).
- Abbot [1] uses the identity $x \cdot (y \cdot z) \approx y \cdot (x \cdot z)$ as one of the defining identities in his definition of implication algebras.
- The identities $x \cdot (y \cdot z) \approx z \cdot (y \cdot x)$, $x \cdot (y \cdot z) \approx y \cdot (x \cdot z)$, and $x \cdot (y \cdot z) \approx (z \cdot x) \cdot y$ were investigated for quasigroups by Hossuzú in [13].
- The identity $x \cdot (z \cdot y) \approx (x \cdot y) \cdot z$ is investigated by Pushkashu in [23].
- The identities $x \cdot (z \cdot y) \approx (x \cdot y) \cdot z$ and $x \cdot (y \cdot z) \approx z \cdot (y \cdot x)$ have appeared in [14] of Kazim and Naseeruddin.

JuanMa, I deleted the esxtra period at the end of the above line.

Ok, thanks!

The identities of mixed type have also been considered in the literature. A few are listed below:

- left distributivity: $x \cdot (y \cdot z) \approx (x \cdot y) \cdot (x \cdot z)$, appears, according to [19], already in the late 19th century publications of logicians Peirce and Schroeder (see [18] and [26], respectively).
- right distributive: $(z \cdot y) \cdot x \approx (z \cdot x) \cdot (y \cdot x)$ (see [27]),
- distributive if it is both left and right distributive (see [27]),
- medial: $(x \cdot y) \cdot (u \cdot v) \approx (x \cdot u) \cdot (y \cdot v)$ (see [27]),
- idempotent: $x \cdot x \approx x$ (see [27]),
- left involutory (or left symmetric): $x \cdot (x \cdot y) \approx y$ (see [27]),

Several identities of associative type have appeared in the literatiure on groupoids as well. For instance,

- $(x \cdot y) \cdot z \approx (z \cdot y) \cdot x$: Abel-Grassmann's groupoid (AG-groupoid, see [22]),
- $(x \cdot y) \cdot z \approx (z \cdot y) \cdot x$ and $x \cdot (y \cdot z) \approx y \cdot (x \cdot z)$ (AG**-groupoid),
- $x \cdot (z \cdot y) \approx (x \cdot y) \cdot z$: Hossuzu-Tarski identity (see [23]),
- $(x \cdot y) \cdot z \approx (z \cdot y) \cdot x$: Left almost semigroup (LA-semigroup) (see [23]),
- $x \cdot (y \cdot z) \approx z \cdot (y \cdot x)$: Right almost semigroup (RA-semigroup) (see [23]).

$$\begin{array}{c|cc} \rightarrow & 0 & 1 \\ \hline 0 & 0 & 0 \\ 1 & 0 & 0 \end{array} \qquad \begin{array}{c|cc} \rightarrow & 0 & 1 \\ \hline 0 & 0 & 1 \\ 1 & 1 & 1 \end{array} \qquad \begin{array}{c|cc} \rightarrow & 0 & 1 \\ \hline 0 & 1 & 1 \\ 1 & 0 & 1 \end{array}$$

Recall that $\mathcal{V}(\mathbf{2}_b) = \mathcal{BA}$. Recall also from [7, Corollary 10.4] that $\mathcal{V}(\mathbf{2}_s) = \mathcal{SL}$. The following lemma easily follows from the definition of \wedge given earlier in the introduction.

LEMMA 2.1. [25, 7.16] *Let \mathbf{A} be an \mathcal{I} -zroupoid. Then $\mathbf{A} \models x''' \rightarrow y \approx x' \rightarrow y$.*

LEMMA 2.2. [7, 3.4] *Let \mathbf{A} be an \mathcal{I} -zroupoid. Then \mathbf{A} satisfies:*

(a) $(x \rightarrow y) \rightarrow z \approx [(x \rightarrow y) \rightarrow z]''$,

(b) $(x \rightarrow y)' \approx (x'' \rightarrow y)'$.

LEMMA 2.3. [25, 8.15] *Let \mathbf{A} be an \mathcal{I} -zroupoid. Then the following are equivalent:*

1. $0' \rightarrow x \approx x$,
2. $x'' \approx x$,
3. $(x \rightarrow x')' \approx x$,
4. $x' \rightarrow x \approx x$.

Recall that $\mathcal{I}_{2,0}$ and \mathcal{MC} are the subvarieties of \mathcal{I} , defined, respectively, by the equations

$$x'' \approx x. \tag{I_{2,0}}$$

$$x \wedge y \approx y \wedge x. \tag{MC}$$

LEMMA 2.4. [25] *Let $\mathbf{A} \in \mathcal{I}_{2,0}$. Then*

1. $x' \rightarrow 0' \approx 0 \rightarrow x$,
2. $0 \rightarrow x' \approx x \rightarrow 0'$.

LEMMA 2.5. *Let $\mathbf{A} \in \mathcal{I}_{2,0}$. Then \mathbf{A} satisfies:*

- (a) $(x \rightarrow 0') \rightarrow y \approx (x \rightarrow y') \rightarrow y$,
- (b) $(y \rightarrow x) \rightarrow y \approx (0 \rightarrow x) \rightarrow y$,
- (c) $0 \rightarrow x \approx 0 \rightarrow (0 \rightarrow x)$,
- (d) $(0 \rightarrow x) \rightarrow (0 \rightarrow y) \approx x \rightarrow (0 \rightarrow y)$,
- (e) $x \rightarrow y \approx x \rightarrow (x \rightarrow y)$,

- (f) $0 \rightarrow (x \rightarrow y) \approx x \rightarrow (0 \rightarrow y)$,
 (g) $0 \rightarrow (x \rightarrow y')' \approx 0 \rightarrow (x' \rightarrow y)$,
 (h) $x \rightarrow (y \rightarrow x') \approx y \rightarrow x'$,

Proof. For the proofs of items (a), (b), (c), (f), (g), and (h) we refer the reader to [7]. The proofs of items (d) and (e) are in [5]. \square

THEOREM 2.6. [8]

Let $t_i(\bar{x}), i = 1, \dots, 6$ be terms and \mathcal{V} a subvariety of \mathcal{I} . If

$$\mathcal{V} \cap \mathcal{I}_{2,0} \models [t_1(\bar{x}) \rightarrow t_2(\bar{x})] \rightarrow t_3(\bar{x}) \approx [t_4(\bar{x}) \rightarrow t_5(\bar{x})] \rightarrow t_6(\bar{x}),$$

then

$$\mathcal{V} \models [t_1(\bar{x}) \rightarrow t_2(\bar{x})] \rightarrow t_3(\bar{x}) \approx [t_4(\bar{x}) \rightarrow t_5(\bar{x})] \rightarrow t_6(\bar{x}).$$

2.1 Identities of Associative Type

We now turn our attention to identities of associative type of length 3. Recall that such an identity will contain three distinct variables that occur in any order and that are grouped in one of the two (obvious) ways. The following identities play a crucial role in the sequel.

Let Σ denote the set consisting of the following 14 identities of associative type (of length 3 in the binary language $\langle \rightarrow \rangle$):

- | | |
|---|---|
| (A1) $x \rightarrow (y \rightarrow z) \approx (x \rightarrow y) \rightarrow z$
(Associative law) | (A8) $x \rightarrow (y \rightarrow z) \approx (z \rightarrow x) \rightarrow y$ |
| (A2) $x \rightarrow (y \rightarrow z) \approx x \rightarrow (z \rightarrow y)$ | (A9) $x \rightarrow (y \rightarrow z) \approx z \rightarrow (y \rightarrow x)$ |
| (A3) $x \rightarrow (y \rightarrow z) \approx (x \rightarrow z) \rightarrow y$ | (A10) $x \rightarrow (y \rightarrow z) \approx (z \rightarrow y) \rightarrow x$ |
| (A4) $x \rightarrow (y \rightarrow z) \approx y \rightarrow (x \rightarrow z)$ | (A11) $(x \rightarrow y) \rightarrow z \approx (x \rightarrow z) \rightarrow y$ |
| (A5) $x \rightarrow (y \rightarrow z) \approx (y \rightarrow x) \rightarrow z$ | (A12) $(x \rightarrow y) \rightarrow z \approx (y \rightarrow x) \rightarrow z$ |
| (A6) $x \rightarrow (y \rightarrow z) \approx y \rightarrow (z \rightarrow x)$ | (A13) $(x \rightarrow y) \rightarrow z \approx (y \rightarrow z) \rightarrow x$ |
| (A7) $x \rightarrow (y \rightarrow z) \approx (y \rightarrow z) \rightarrow x$ | (A14) $(x \rightarrow y) \rightarrow z \approx (z \rightarrow y) \rightarrow x$. |

We will denote by \mathcal{A}_i the subvariety of \mathcal{I} defined by the identity (Ai), for $1 \leq i \leq 14$. Such varieties will be referred to as subvarieties of \mathcal{I} of associative type.

The following proposition is crucial for the rest of the paper.

PROPOSITION 2.7. *Let \mathcal{G} be the variety of all groupoids of type $\{\rightarrow\}$ and Let \mathcal{V} denote the subvariety of \mathcal{G} defined by a single identity of associative type. Then $\mathcal{V} = \mathcal{A}_i$, for **some** $i \in \{1, 2, \dots, 14\}$.*

Proof. In an identity $p \approx q$ of associative type of length 3, p and q have exactly 3 (distinct) variables, say x, y, z , and these variables are grouped according to one of the two ways of bracketing mentioned above. Thus, there are six permutations of 3 variables which give rise to the following 12 terms:

$$\begin{array}{ll}
 1a : x \rightarrow (y \rightarrow z) & 1b: (x \rightarrow y) \rightarrow z \\
 2a : x \rightarrow (z \rightarrow y) & 2b: (x \rightarrow z) \rightarrow y \\
 3a : y \rightarrow (x \rightarrow z) & 3b: (y \rightarrow x) \rightarrow z \\
 4a : y \rightarrow (z \rightarrow x) & 4b: (y \rightarrow z) \rightarrow x \\
 5a : z \rightarrow (x \rightarrow y) & 5b: (z \rightarrow x) \rightarrow y \\
 6a : z \rightarrow (y \rightarrow x) & 6b: (z \rightarrow y) \rightarrow x.
 \end{array}$$

It is clear that these 12 terms, in turn, will lead to 66 identities in view of the symmetric property of equality. It is routine to verify that each of the 66 identities is equivalent to one of the 14 identities of Σ in the variety of groupoids. Then the proposition follows. \square

Our goal, in this paper, is to determine the distinct subvarieties of \mathcal{I} and to describe the poset of subvarieties of \mathcal{I} of associative type. It suffices to concentrate on the varieties defined by identities (A1)-(A14), in view of the above proposition.

3 Properties of subvarieties of \mathcal{I} of Associative type

In this section we present properties of several subvarieties of \mathcal{I} which will play a crucial role in our analysis of the identities of associative type relative to \mathcal{I} .

LEMMA 3.1. *Let $\mathbf{A} \in \mathcal{I}$ such that $\mathbf{A} \models x' \rightarrow y \approx x \rightarrow y'$, then $\mathbf{A} \models (x \rightarrow y) \rightarrow y' \approx x \rightarrow y'$.*

Proof. Let $a, b \in A$. Then $(a \rightarrow b) \rightarrow b' \stackrel{2.5(a) \& 2.6}{=} (a \rightarrow 0') \rightarrow b' \stackrel{hyp}{=} (a' \rightarrow 0) \rightarrow b' = a'' \rightarrow b' \stackrel{hyp}{=} a''' \rightarrow b \stackrel{2.1}{=} a' \rightarrow b \stackrel{hyp}{=} a \rightarrow b'$. \square

LEMMA 3.2. *Let $\mathbf{A} \in \mathcal{I}_{2,0}$ such that $\mathbf{A} \models (x \rightarrow y)' \approx x \rightarrow (0 \rightarrow y)$, then $\mathbf{A} \models x \rightarrow y' \approx x \rightarrow (0 \rightarrow y)$.*

Proof. Let $a, b \in A$. Then $a \rightarrow b' \stackrel{2.3(1)}{=} a \rightarrow (0' \rightarrow b)' \stackrel{hyp}{=} a \rightarrow [0' \rightarrow (0 \rightarrow b)] \stackrel{2.3(1)}{=} a \rightarrow (0 \rightarrow b)$. \square

LEMMA 3.3. *Let $\mathbf{A} \in \mathcal{I}_{2,0}$ such that $\mathbf{A} \models (x \rightarrow y)' \approx x \rightarrow (0 \rightarrow y)$. Then $\mathbf{A} \models [x \rightarrow (y \rightarrow z)']' \approx x \rightarrow (y \rightarrow (0 \rightarrow z))'$.*

Proof. Let $a, b, c \in A$. We have that $[a \rightarrow (b \rightarrow c)']' \stackrel{hyp}{=} a \rightarrow (0 \rightarrow (b \rightarrow c))'$
 $\stackrel{hyp}{=} a \rightarrow (0 \rightarrow (b \rightarrow (0 \rightarrow c))) \stackrel{2.5(f)}{=} a \rightarrow (b \rightarrow (0 \rightarrow (0 \rightarrow c))) \stackrel{hyp}{=} a \rightarrow [b \rightarrow (0 \rightarrow c)']$. \square

LEMMA 3.4. *Let $\mathbf{A} \in \mathcal{I}$ such that \mathbf{A} satisfies:*

$$(1) (x \rightarrow y)' \approx x \rightarrow (0 \rightarrow y),$$

$$(2) x' \rightarrow y \approx x \rightarrow y'.$$

Then, $\mathbf{A} \models 0 \rightarrow [x \rightarrow (y \rightarrow z)] \approx 0 \rightarrow [(x \rightarrow y) \rightarrow z]$.

Proof. Let $a, b, c \in A$. Then, $0 \rightarrow [(a \rightarrow b) \rightarrow c] \stackrel{(I)}{=} 0 \rightarrow [(c' \rightarrow a) \rightarrow (b \rightarrow c)']'$
 $\stackrel{(I)}{=} 0 \rightarrow \{[(b \rightarrow c)'' \rightarrow c'] \rightarrow [a \rightarrow (b \rightarrow c)']\}'' \stackrel{2.6}{=} 0 \rightarrow \{[(b \rightarrow c) \rightarrow c'] \rightarrow [a \rightarrow (b \rightarrow c)']\}''$
 $\stackrel{3.1}{=} 0 \rightarrow \{[b \rightarrow c'] \rightarrow [a \rightarrow (b \rightarrow c)']\}'' \stackrel{3.2 \& 2.6 \& hyp}{=} 0 \rightarrow \{[b \rightarrow (0 \rightarrow c)] \rightarrow [a \rightarrow (b \rightarrow c)']\}''$
 $\stackrel{3.3 \& hyp}{=} 0 \rightarrow \{[b \rightarrow (0 \rightarrow c)] \rightarrow [a \rightarrow [b \rightarrow (0 \rightarrow c)]]\}'' \stackrel{2.5(h) \& 2.6}{=} 0 \rightarrow \{a \rightarrow [b \rightarrow (0 \rightarrow c)']\}'' \stackrel{(1)}{=} 0 \rightarrow \{a \rightarrow [b \rightarrow c]''\}'' \stackrel{2.6}{=} 0 \rightarrow \{a \rightarrow [b \rightarrow c]\}''$
 $\stackrel{(2)}{=} 0' \rightarrow \{a \rightarrow [b \rightarrow c]\}' \stackrel{(2)}{=} 0'' \rightarrow \{a \rightarrow [b \rightarrow c]\} = 0 \rightarrow \{a \rightarrow [b \rightarrow c]\}$. \square

LEMMA 3.5. *Let $\mathbf{A} \in \mathcal{I}$ such that $\mathbf{A} \models (x \rightarrow y)' \approx (y \rightarrow x)'$. Then $\mathbf{A} \models (x \rightarrow y) \rightarrow z \approx (y \rightarrow x) \rightarrow z$.*

Proof. Let $a, b, c \in A$. Then, $(a \rightarrow b) \rightarrow c \stackrel{2.6}{=} (a \rightarrow b)'' \rightarrow c \stackrel{hyp}{=} (b \rightarrow a)'' \rightarrow c$
 $\stackrel{2.6}{=} (b \rightarrow a) \rightarrow c$. \square

DEFINITION 3.6. *Let $\mathbf{A} \in \mathcal{I}$. We say that \mathcal{A} is of type 1 if the following identities hold in \mathbf{A} :*

$$(E1) (x \rightarrow y)' \approx x \rightarrow (0 \rightarrow y),$$

$$(E2) x' \rightarrow y \approx x \rightarrow y',$$

$$(E3) 0 \rightarrow (x \rightarrow y) \approx 0 \rightarrow (y \rightarrow x),$$

$$(E4) x \rightarrow (y \rightarrow z) \approx (p(x) \rightarrow p(y)) \rightarrow p(z), \text{ where } p \text{ is some permutation of } \{x, y, z\}.$$

THEOREM 3.7. *If $\mathbf{A} \in \mathcal{I}$ is of type 1 then $\mathbf{A} \models (A_j)$ for all $1 \leq j \leq 14$.*

Proof. Let $\mathbf{A} \in \mathcal{I}$ be of type 1, and $a, b, c \in A$. In view of equations (E1), (E2) and Lemma 3.4 we have that

$$\mathbf{A} \models 0 \rightarrow [x \rightarrow (y \rightarrow z)] \approx 0 \rightarrow [(x \rightarrow y) \rightarrow z]. \quad (3.1)$$

Then we can consider the following cases.

- Assume that $j = 1$. Then $a \rightarrow (b \rightarrow c) \stackrel{(E4)}{\equiv} (p(a) \rightarrow p(b)) \rightarrow p(c) \stackrel{2.2}{\equiv} [(p(a) \rightarrow p(b)) \rightarrow p(c)]'' \stackrel{(E1)}{\equiv} [(p(a) \rightarrow p(b)) \rightarrow (0 \rightarrow p(c))]' \stackrel{2.5(f) \& 2.6}{\equiv} [0 \rightarrow [(p(a) \rightarrow p(b)) \rightarrow p(c)]]' \stackrel{(E3) \& (3.1)}{\equiv} [0 \rightarrow [(a \rightarrow b) \rightarrow c]]' \stackrel{2.5(f) \& 2.6}{\equiv} [(a \rightarrow b) \rightarrow [0 \rightarrow c]]' \stackrel{(E1)}{\equiv} [(a \rightarrow b) \rightarrow c]'' \stackrel{2.2}{\equiv} (a \rightarrow b) \rightarrow c$.

The cases $j = 3, 5, 7, 8, 10$ are similar.

- Assume that $j = 2$. Then, in the same way as in the case of $j = 1$ we have that

$$\mathbf{A} \models x \rightarrow (y \rightarrow z) \approx [0 \rightarrow [(p(x) \rightarrow p(y)) \rightarrow p(z)]]'. \quad (3.2)$$

Then, $a \rightarrow (b \rightarrow c) \stackrel{(3.2)}{\equiv} [0 \rightarrow [(p(a) \rightarrow p(b)) \rightarrow p(c)]]' \stackrel{(E3) \& (3.1)}{\equiv} [0 \rightarrow [(p(a) \rightarrow p(c)) \rightarrow p(b)]]' \stackrel{2.5(f) \& 2.6}{\equiv} [(p(a) \rightarrow p(c)) \rightarrow [0 \rightarrow p(b)]]' \stackrel{(E1)}{\equiv} [(p(a) \rightarrow p(c)) \rightarrow p(b)]'' \stackrel{2.2(a)}{\equiv} (p(a) \rightarrow p(c)) \rightarrow p(b) \stackrel{(E4)}{\equiv} (a \rightarrow c) \rightarrow b$.

The cases $j = 4, 6, 9$ are similar.

- Assume that $j = 11$. $(a \rightarrow b) \rightarrow c \stackrel{2.2(a)}{\equiv} [(a \rightarrow b) \rightarrow c]'' \stackrel{(E1)}{\equiv} [(a \rightarrow b) \rightarrow (0 \rightarrow c)]' \stackrel{2.5(f) \& 2.6}{\equiv} [0 \rightarrow [(a \rightarrow b) \rightarrow c]]' \stackrel{(E3) \& (3.1)}{\equiv} [0 \rightarrow [(a \rightarrow c) \rightarrow b]]' \stackrel{2.5(f) \& 2.6}{\equiv} [(a \rightarrow c) \rightarrow [0 \rightarrow b]]' \stackrel{(E1)}{\equiv} [(a \rightarrow c) \rightarrow b]'' \stackrel{2.2}{\equiv} (a \rightarrow c) \rightarrow b$.

The cases $j = 12, 13, 14$ are similar.

□

To prove that the variety \mathcal{A}_j is of type 1, with $j \in \{3, 5, 7, 8, 10\}$, we need the following lemmas.

LEMMA 3.8. *If $\mathbf{A} \in \mathcal{A}_5$ then \mathbf{A} satisfies*

(a) $x' \rightarrow y \approx x \rightarrow y'$

(b) $(x \rightarrow y)' \approx 0 \rightarrow (x \rightarrow y)$

(c) $x \rightarrow (0 \rightarrow y) \approx 0 \rightarrow (x \rightarrow y)$

Proof. Let $a, b \in A$. Then

(a) $a \rightarrow b' = a \rightarrow (b \rightarrow 0) \stackrel{(A5)}{\equiv} (b \rightarrow a) \rightarrow 0 \stackrel{2.6}{\equiv} (b \rightarrow a'') \rightarrow 0 = (b \rightarrow (a' \rightarrow 0)) \rightarrow 0 \stackrel{(A5)}{\equiv} [(a' \rightarrow b) \rightarrow 0] \rightarrow 0 = (a' \rightarrow b)'' \stackrel{2.6}{\equiv} a' \rightarrow b$.

(b) Observe that $(a \rightarrow b)' = (a \rightarrow b) \rightarrow 0 \stackrel{(A5)}{\equiv} b \rightarrow (a \rightarrow 0) = b \rightarrow a' \stackrel{3.8(a)}{\equiv} b' \rightarrow a = (b \rightarrow 0) \rightarrow a \stackrel{(A5)}{\equiv} 0 \rightarrow (b \rightarrow a)$.

(c) Notice that $0 \rightarrow (a \rightarrow b) \stackrel{(A5)}{\equiv} (a \rightarrow 0) \rightarrow b \stackrel{2.6}{\equiv} (a'' \rightarrow 0) \rightarrow b \stackrel{3.8(a)}{\equiv} (a' \rightarrow 0') \rightarrow b \stackrel{2.4(2) \& 2.6}{\equiv} (0 \rightarrow a) \rightarrow b \stackrel{(A5)}{\equiv} a \rightarrow (0 \rightarrow b)$.

□

LEMMA 3.9. *If $\mathbf{A} \in \mathcal{A}_8$ then \mathbf{A} satisfies:*

(a) $x \rightarrow y' \approx x' \rightarrow y'$,

(b) $x \rightarrow y' \approx 0 \rightarrow (y' \rightarrow x)$.

Proof. Let $a, b \in A$. Then $a \rightarrow b' = a \rightarrow (b \rightarrow 0) \stackrel{(A8)}{=} (0 \rightarrow a) \rightarrow b \stackrel{(I)}{=} [(b' \rightarrow 0) \rightarrow (a \rightarrow b)'] = [(b' \rightarrow 0) \rightarrow (a \rightarrow b)'] \rightarrow 0 \stackrel{(A8)}{=} (a \rightarrow b)' \rightarrow (0 \rightarrow (b' \rightarrow 0)) \stackrel{(A8)}{=} (a \rightarrow b)' \rightarrow ((0 \rightarrow 0) \rightarrow b') = (a \rightarrow b)' \rightarrow (0' \rightarrow b') \stackrel{2.3(1) \& 2.6}{=} (a \rightarrow b)' \rightarrow b' = ((a \rightarrow b) \rightarrow 0) \rightarrow b' \stackrel{(A8)}{=} 0 \rightarrow (b' \rightarrow (a \rightarrow b)) \stackrel{(A8)}{=} 0 \rightarrow ((b \rightarrow b') \rightarrow a) \stackrel{2.6}{=} 0 \rightarrow ((b'' \rightarrow b') \rightarrow a) \stackrel{2.3(4) \& 2.6}{=} 0 \rightarrow (b' \rightarrow a)$, implying that \mathbf{A} satisfies the identity (b). Next, $0 \rightarrow (b' \rightarrow a) \stackrel{(A8)}{=} (a \rightarrow 0) \rightarrow b' = a' \rightarrow b'$, thus \mathbf{A} satisfies the identity (a). □

LEMMA 3.10. *If $\mathbf{A} \in \mathcal{A}_{10}$ then \mathbf{A} satisfies:*

(a) $[0 \rightarrow (x \rightarrow y)]' \approx x \rightarrow y'$,

(b) $(y \rightarrow x)'' \approx x \rightarrow y'$,

(c) $(x \rightarrow y)' \approx x \rightarrow y'$.

Proof. Let $a, b \in A$.

(a) We have that

$$\mathbf{A} \models [0 \rightarrow (x \rightarrow y)]' \approx x \rightarrow y',$$

since

$$\begin{aligned} a \rightarrow b' &= a \rightarrow (b \rightarrow 0) \\ &= (0 \rightarrow b) \rightarrow a && \text{by (A10)} \\ &= [(a' \rightarrow 0) \rightarrow (b \rightarrow a)'] && \text{by (I)} \\ &= [((a \rightarrow 0) \rightarrow 0) \rightarrow (b \rightarrow a)'] \\ &= [(0 \rightarrow (0 \rightarrow a)) \rightarrow (b \rightarrow a)'] && \text{by (A10)} \\ &= [(0 \rightarrow a) \rightarrow (b \rightarrow a)'] && \text{by 2.5 (c) and 2.6} \\ &= [(0 \rightarrow a) \rightarrow [(b \rightarrow a) \rightarrow 0]]' \\ &= [(0 \rightarrow a) \rightarrow [0 \rightarrow (a \rightarrow b)]]' && \text{by (A10)} \\ &= [(0 \rightarrow a) \rightarrow [(0 \rightarrow a) \rightarrow (0 \rightarrow b)]]' && \text{by 2.5 (f) and (d) and by 2.6} \\ &= [(0 \rightarrow a) \rightarrow (0 \rightarrow b)]' && \text{by 2.5 (e) and 2.6} \\ &= [0 \rightarrow (a \rightarrow b)]' && \text{by 2.5 (d), (f) and by 2.6.} \end{aligned}$$

(b) Observe that $a \rightarrow b' \stackrel{3.10(a)}{=} [0 \rightarrow (a \rightarrow b)]' \stackrel{(A10)}{=} [(b \rightarrow a) \rightarrow 0]' = (b \rightarrow a)''$. Hence, $\mathbf{A} \models$ (b).

- (c) Since $a \rightarrow b' \stackrel{3.10(a)}{=} [0 \rightarrow (a \rightarrow b)]' \stackrel{2.5(c) \& 2.6}{=} [0 \rightarrow (0 \rightarrow (a \rightarrow b))]'$ $\stackrel{(A10)}{=} [((a \rightarrow b) \rightarrow 0) \rightarrow 0]' = (a \rightarrow b)''' = (a \rightarrow b)'$, we conclude that $\mathbf{A} \models (c)$.

□

LEMMA 3.11. *If $\mathbf{A} \in \mathcal{A}_3 \cup \mathcal{A}_5 \cup \mathcal{A}_7 \cup \mathcal{A}_8 \cup \mathcal{A}_{10}$ then \mathbf{A} satisfies*

- (1) $(x \rightarrow y)' \approx x \rightarrow (0 \rightarrow y)$,
- (2) $x' \rightarrow y \approx x \rightarrow y'$ and
- (3) $0 \rightarrow (x \rightarrow y) \approx 0 \rightarrow (y \rightarrow x)$.

Proof. Let $a, b \in A$.

- Suppose $\mathbf{A} \in \mathcal{A}_3$. Then $(a \rightarrow b)' = (a \rightarrow b) \rightarrow 0 \stackrel{(A3)}{=} a \rightarrow (0 \rightarrow b)$, implying $\mathbf{A} \models (1)$.

Observe that $a' \rightarrow b = (a \rightarrow 0) \rightarrow b \stackrel{(A3)}{=} a \rightarrow (b \rightarrow 0) = a \rightarrow b'$. So, (2) holds in \mathbf{A} .

Also, $0 \rightarrow (a \rightarrow b) \stackrel{(A3)}{=} (0 \rightarrow b) \rightarrow a \stackrel{(I)}{=} [(a' \rightarrow 0) \rightarrow (b \rightarrow a)']' = [a'' \rightarrow (b \rightarrow a)']' \stackrel{2.6}{=} [a \rightarrow (b \rightarrow a)']' \stackrel{(3.11)}{=} [a' \rightarrow (b \rightarrow a)]' \stackrel{2.5(h) \& 2.6}{=} (b \rightarrow a)' \stackrel{2.3(1) \& 2.6}{=} 0' \rightarrow (b \rightarrow a)' \stackrel{3.11(2)}{=} 0'' \rightarrow (b \rightarrow a) = 0 \rightarrow (b \rightarrow a)$, proving that (3) holds in \mathbf{A} .

- Assume that $\mathbf{A} \in \mathcal{A}_5$. Then $(a \rightarrow b)' \stackrel{3.8(b)}{=} 0 \rightarrow (b \rightarrow a) = 0'' \rightarrow (b \rightarrow a) \stackrel{3.8(a)}{=} 0' \rightarrow (b \rightarrow a)' \stackrel{2.3(1) \& 2.6}{=} (b \rightarrow a)' = (b \rightarrow a) \rightarrow 0 \stackrel{(A5)}{=} a \rightarrow (b \rightarrow 0) = a \rightarrow b' \stackrel{3.8(a)}{=} a' \rightarrow b \stackrel{2.3(1) \& 2.6}{=} (0' \rightarrow a') \rightarrow b \stackrel{3.8(a)}{=} (0 \rightarrow a'') \rightarrow b \stackrel{2.6}{=} (0 \rightarrow a) \rightarrow b \stackrel{(A5)}{=} a \rightarrow (0 \rightarrow b)$, proving that (1) is true in \mathbf{A} . (2) is immediate from Lemma 3.8 (a). Next, $0 \rightarrow (a \rightarrow b) \stackrel{3.8(b)}{=} (a \rightarrow b)' = (a \rightarrow b) \rightarrow 0 \stackrel{(A5)}{=} b \rightarrow (a \rightarrow 0) = b \rightarrow a' \stackrel{3.8(a)}{=} b' \rightarrow a \stackrel{2.6}{=} (b' \rightarrow a)'' \stackrel{3.8(a)}{=} (b \rightarrow a)'' \stackrel{3.8(b)}{=} [0 \rightarrow (b \rightarrow a)']' \stackrel{3.8(c)}{=} [b \rightarrow (0 \rightarrow a)']' \stackrel{3.8(a)}{=} [b \rightarrow (0' \rightarrow a)']' \stackrel{2.3(1) \& 2.6}{=} (b \rightarrow a)' \stackrel{3.8(b)}{=} 0 \rightarrow (b \rightarrow a)$, proving (3) holds in \mathbf{A} .

- Assume that $\mathbf{A} \in \mathcal{A}_7$. Then $(a \rightarrow b)' \stackrel{2.6}{=} (a \rightarrow b)''' = [(a \rightarrow b) \rightarrow 0]'' \stackrel{(A7)}{=} [0 \rightarrow (a \rightarrow b)]'' \stackrel{2.5(f) \& 2.6}{=} [a \rightarrow (0 \rightarrow b)]'' \stackrel{(A7)}{=} [(0 \rightarrow b) \rightarrow a]'' \stackrel{2.6}{=} (0 \rightarrow b) \rightarrow a \stackrel{(A7)}{=} a \rightarrow (0 \rightarrow b)$, proving that \mathbf{A} satisfies (1).

Next, $a' \rightarrow b \stackrel{2.6}{=} [a' \rightarrow b]'' = [(a' \rightarrow b) \rightarrow 0]' \stackrel{(A7)}{=} [0 \rightarrow (a' \rightarrow b)]' \stackrel{2.6}{=} [0 \rightarrow (a' \rightarrow b'')] \stackrel{2.5(g) \& 2.6}{=} [0 \rightarrow (a \rightarrow b)']' = [0 \rightarrow ((a \rightarrow b') \rightarrow 0)]' \stackrel{(A7)}{=} [0 \rightarrow (0 \rightarrow (a \rightarrow b'))]' \stackrel{2.5(c) \& 2.6}{=} [0 \rightarrow (a \rightarrow b)']' = [0 \rightarrow (a \rightarrow b')] \rightarrow 0$

$\stackrel{(A7)}{=} [(a \rightarrow b') \rightarrow 0] \rightarrow 0 = (a \rightarrow b')'' = (a \rightarrow (b \rightarrow 0))'' \stackrel{(A7)}{=} ((b \rightarrow 0) \rightarrow a)''$
 $\stackrel{2.6}{=} (b \rightarrow 0) \rightarrow a \stackrel{(A7)}{=} a \rightarrow (b \rightarrow 0) = a \rightarrow b'$, proving that (2) is true in \mathbf{A} .
 Finally, observe that $0 \rightarrow (a \rightarrow b) \stackrel{(A7)}{=} (a \rightarrow b) \rightarrow 0 \stackrel{2.6}{=} (a'' \rightarrow b) \rightarrow 0 =$
 $[(a' \rightarrow 0) \rightarrow b] \rightarrow 0 \stackrel{(A7)}{=} [b \rightarrow (a' \rightarrow 0)] \rightarrow 0 = [b \rightarrow a''] \rightarrow 0 \stackrel{2.6}{=} [b \rightarrow a] \rightarrow 0$
 $\stackrel{(A7)}{=} 0 \rightarrow (b \rightarrow a)$, proving that (3) holds in \mathbf{A} .

- Let $\mathbf{A} \in \mathcal{A}_8$. First, we will prove (2) hold in \mathbf{A} . Now, $a \rightarrow b' \stackrel{3.9(b)}{=} 0 \rightarrow$
 $(b' \rightarrow a) \stackrel{2.6}{=} 0 \rightarrow (b' \rightarrow a'') \stackrel{3.9(a)}{=} 0 \rightarrow (b'' \rightarrow a'') \stackrel{(A8)}{=} (a'' \rightarrow 0) \rightarrow b''$
 $\stackrel{2.6}{=} (a'' \rightarrow 0) \rightarrow b = a''' \rightarrow b \stackrel{2.1}{=} a' \rightarrow b$, proving (2).

Notice that $a \rightarrow (0 \rightarrow b) \stackrel{(A8)}{=} (b \rightarrow a) \rightarrow 0 = (b \rightarrow a)' \stackrel{2.6}{=} (b \rightarrow a)'''$
 $= (b \rightarrow a)'' \rightarrow 0 \stackrel{(2)}{=} (b \rightarrow a)' \rightarrow 0' \stackrel{2.3(1) \& 2.6}{=} [0' \rightarrow (b \rightarrow a)'] \rightarrow 0' \stackrel{(2)}{=}$
 $[0'' \rightarrow (b \rightarrow a)] \rightarrow 0' = [0 \rightarrow (b \rightarrow a)] \rightarrow 0' \stackrel{2.5(f) \& 2.6}{=} [b \rightarrow (0 \rightarrow a)] \rightarrow 0'$
 $\stackrel{(A8)}{=} [(a \rightarrow b) \rightarrow 0] \rightarrow 0' = (a \rightarrow b)' \rightarrow 0' \stackrel{(2)}{=} (a \rightarrow b) \rightarrow 0'' = (a \rightarrow b) \rightarrow 0$
 $= (a \rightarrow b)'$, proving (1) holds in \mathbf{A} .

The identity

$$0 \rightarrow (x \rightarrow y) \approx (x \rightarrow y)' \quad (3.3)$$

holds in \mathbf{A} , since $0 \rightarrow (a \rightarrow b) = 0'' \rightarrow (a \rightarrow b) \stackrel{(2)}{=} 0' \rightarrow (a \rightarrow b)' \stackrel{2.3(1) \& 2.6}{=}$
 $(a \rightarrow b)'$.

Then $0 \rightarrow (a \rightarrow b) \stackrel{(3.3)}{=} (a \rightarrow b)' = (a \rightarrow b) \rightarrow 0 \stackrel{(A8)}{=} b \rightarrow (0 \rightarrow a) \stackrel{(1)}{=} (b \rightarrow a)'$
 $\stackrel{(3.3)}{=} 0 \rightarrow (b \rightarrow a)$, proving (3) is true in \mathbf{A} .

- Assume that $\mathbf{A} \in \mathcal{A}_{10}$. Hence

$$\begin{aligned} a' \rightarrow b &= a' \rightarrow b'' && \text{by 2.6} \\ &= (a' \rightarrow b')' && \text{by 3.10 (c)} \\ &= (a' \rightarrow b') \rightarrow 0 \\ &= 0 \rightarrow (b' \rightarrow a') && \text{by (A10)} \\ &= 0 \rightarrow (b' \rightarrow (a \rightarrow 0)) \\ &= 0 \rightarrow ((0 \rightarrow a) \rightarrow b') && \text{by (A10)} \\ &= (b' \rightarrow (0 \rightarrow a)) \rightarrow 0 && \text{by (A10)} \\ &= (b' \rightarrow (0 \rightarrow a))' \\ &= ((b \rightarrow 0) \rightarrow (0 \rightarrow a))' \\ &= [(0 \rightarrow a) \rightarrow (0 \rightarrow b)]' && \text{by (A10)} \\ &= [0 \rightarrow (a \rightarrow b)]' && \text{by 2.5 (f) \& (d) and by 2.6} \\ &= [(b \rightarrow a) \rightarrow 0]' && \text{by (A10)} \\ &= (b \rightarrow a)'' \\ &= a \rightarrow b' && \text{by 3.10 (b),} \end{aligned}$$

proving (2) holds in \mathbf{A} .

Consider $a \rightarrow (0 \rightarrow b) \stackrel{(A10)}{=} (b \rightarrow 0) \rightarrow a \stackrel{(I)}{=} [(a' \rightarrow b) \rightarrow (0 \rightarrow a)]' \stackrel{3.10(c)}{=} [(a' \rightarrow b) \rightarrow (0 \rightarrow a)']' \stackrel{(2)}{=} [(a' \rightarrow b) \rightarrow (0' \rightarrow a)]' \stackrel{2.3(1) \& 2.6}{=} [(a' \rightarrow b) \rightarrow a]' \stackrel{(2)}{=} [(a \rightarrow b') \rightarrow a]' \stackrel{2.5(b) \& 2.6}{=} [(0 \rightarrow b') \rightarrow a]' \stackrel{(A10)}{=} [a \rightarrow (b' \rightarrow 0)]' = [a \rightarrow b'']' \stackrel{2.6}{=} (a \rightarrow b)'$, proving (1).

To finish off the proof, $0 \rightarrow (a \rightarrow b) \stackrel{(A10)}{=} (b \rightarrow a) \rightarrow 0 \stackrel{3.10(c)}{=} b \rightarrow a' \stackrel{(2)}{=} b' \rightarrow a = (b \rightarrow 0) \rightarrow a \stackrel{(A10)}{=} a \rightarrow (0 \rightarrow b) \stackrel{(1)}{=} (a \rightarrow b)' = (a \rightarrow b) \rightarrow 0 \stackrel{(A10)}{=} 0 \rightarrow (b \rightarrow a)$.

□

THEOREM 3.12. $\mathcal{A}_3 = \mathcal{A}_5 = \mathcal{A}_7 = \mathcal{A}_8 = \mathcal{A}_{10}$.

Proof. Let $\mathbf{A} \in \mathcal{A}_3 \cup \mathcal{A}_5 \cup \mathcal{A}_7 \cup \mathcal{A}_8 \cup \mathcal{A}_{10}$. By Lemma 3.11 we have that \mathbf{A} is of type 1. Then, using Theorem 3.7, $\mathbf{A} \in \mathcal{A}_j$ for all $j \in \{3, 5, 7, 8, 10\}$. □

LEMMA 3.13. *If $\mathbf{A} \in \mathcal{A}_{13}$ then \mathbf{A} satisfies*

(a) $(x \rightarrow y)' \approx (0 \rightarrow x) \rightarrow y$

(b) $(x \rightarrow y)' \approx x' \rightarrow y'$

(c) $(x \rightarrow y)' \approx (0 \rightarrow y) \rightarrow x'$

(d) $(x \rightarrow y)' \approx (x \rightarrow y)''$

(e) $(x \rightarrow y)' \approx (y \rightarrow x)'$

Proof. Let us consider $a, b \in A$.

(a) $(a \rightarrow b)' = (a \rightarrow b) \rightarrow 0 \stackrel{(A13)}{=} (b \rightarrow 0) \rightarrow a \stackrel{(A13)}{=} (0 \rightarrow a) \rightarrow b$. Hence $\mathbf{A} \models$ (a).

(b) Observe that $(a \rightarrow b)' \stackrel{2.6}{=} 0' \rightarrow (a \rightarrow b)' = (0 \rightarrow 0) \rightarrow (a \rightarrow b)' \stackrel{(A13)}{=} [0 \rightarrow (a \rightarrow b)'] \rightarrow 0 \stackrel{(I20) \& 2.6}{=} [0 \rightarrow (a \rightarrow b'')] \rightarrow 0 \stackrel{2.5(g) \& 2.6}{=} [0 \rightarrow (a' \rightarrow b)'] \rightarrow 0 \stackrel{(A13)}{=} [(a' \rightarrow b') \rightarrow 0] \rightarrow 0 = (a' \rightarrow b')'' \stackrel{2.6}{=} a' \rightarrow b'$

(c) Observe $(a \rightarrow b)' \stackrel{(a)}{=} (0 \rightarrow a) \rightarrow b \stackrel{2.6}{=} (0 \rightarrow a)'' \rightarrow b \stackrel{(b)}{=} (0' \rightarrow a) \rightarrow b \stackrel{2.6}{=} a'' \rightarrow b \stackrel{(A13)}{=} (0 \rightarrow b) \rightarrow a'$.

(d) Note that $(a \rightarrow b)' \stackrel{(c)}{=} (0 \rightarrow b) \rightarrow a' \stackrel{(I)}{=} [(a'' \rightarrow 0) \rightarrow (b \rightarrow a)']' = [a''' \rightarrow (b \rightarrow a)']' \stackrel{2.1}{=} [a' \rightarrow (b \rightarrow a)']' \stackrel{(b)}{=} [a' \rightarrow (b' \rightarrow a'')] \stackrel{2.6}{=} [a' \rightarrow (b' \rightarrow a)]' \stackrel{2.5(h) \& 2.6}{=} (b' \rightarrow a)' \stackrel{(A13)}{=} ((0 \rightarrow a) \rightarrow b)' \stackrel{(A13)}{=} ((a \rightarrow b) \rightarrow 0)' = (a \rightarrow b)''$.

(e) We have $(b \rightarrow a)' \stackrel{(c)}{\underline{=}} (0 \rightarrow a) \rightarrow b' \stackrel{2.6}{\underline{=}} [(0 \rightarrow a) \rightarrow b]'' \stackrel{(b)}{\underline{=}} [(0 \rightarrow a)' \rightarrow b]''$
 $\stackrel{(d)}{\underline{=}} [(0 \rightarrow a)'' \rightarrow b]'' \stackrel{2.6}{\underline{=}} [(0 \rightarrow a) \rightarrow b]'' \stackrel{(d)}{\underline{=}} [(0 \rightarrow a) \rightarrow b]'' \stackrel{2.6}{\underline{=}} (0 \rightarrow a) \rightarrow b$
 $\stackrel{(a)}{\underline{=}} (a \rightarrow b)'.$

□

THEOREM 3.14. $\mathcal{A}_{11} = \mathcal{A}_{12} = \mathcal{A}_{13}.$

Proof. Let us consider $\mathbf{A} \in \mathcal{A}_{11}$ and $a, b, c \in A$. Hence $(a \rightarrow b) \rightarrow c \stackrel{2.3(1) \& 2.6}{\underline{=}}$
 $((0' \rightarrow a) \rightarrow b) \rightarrow c \stackrel{(A11)}{\underline{=}} ((0' \rightarrow b) \rightarrow a) \rightarrow c \stackrel{2.3(1) \& 2.6}{\underline{=}} (b \rightarrow a) \rightarrow c$. Hence,
 $\mathbf{A} \in \mathcal{A}_{12}$, implying $\mathcal{A}_{11} \subseteq \mathcal{A}_{12}$.

Now assume that $\mathbf{A} \in \mathcal{A}_{12}$ and $a, b, c \in A$. Then $(a \rightarrow b) \rightarrow c \stackrel{(I)}{\underline{=}} [(c' \rightarrow a) \rightarrow$
 $(b \rightarrow c)]' \stackrel{(I)}{\underline{=}} \{[(b \rightarrow c)'' \rightarrow c'] \rightarrow [a \rightarrow (b \rightarrow c)]'\}''' \stackrel{2.6}{\underline{=}} \{[(b \rightarrow c) \rightarrow c'] \rightarrow [a \rightarrow$
 $(b \rightarrow c)]'\}''' \stackrel{(A12)}{\underline{=}} \{[c' \rightarrow (b \rightarrow c)] \rightarrow [a \rightarrow (b \rightarrow c)]'\}''' \stackrel{2.5(h) \& 2.6}{\underline{=}} \{(b \rightarrow c) \rightarrow$
 $[a \rightarrow (b \rightarrow c)]'\}''' \stackrel{(A12)}{\underline{=}} \{[a \rightarrow (b \rightarrow c)]' \rightarrow (b \rightarrow c)\}'' = \{[[a \rightarrow (b \rightarrow c)]' \rightarrow$
 $0] \rightarrow (b \rightarrow c)\}'' \stackrel{(A12)}{\underline{=}} \{[0 \rightarrow [a \rightarrow (b \rightarrow c)]] \rightarrow (b \rightarrow c)\}'' \stackrel{2.5(b) \& 2.6}{\underline{=}} \{(b \rightarrow c) \rightarrow$
 $[a \rightarrow (b \rightarrow c)]\}'' \stackrel{2.5(h) \& 2.6}{\underline{=}} \{[a \rightarrow (b \rightarrow c)]' \rightarrow (b \rightarrow c)\}'' \stackrel{2.5(a) \& 2.6}{\underline{=}}$
 $\{[a \rightarrow 0'] \rightarrow (b \rightarrow c)\}'' \stackrel{(A12)}{\underline{=}} \{[0' \rightarrow a] \rightarrow (b \rightarrow c)\}'' \stackrel{2.3(1) \& 2.6}{\underline{=}} \{a \rightarrow (b \rightarrow c)\}''$
 $\stackrel{(A12)}{\underline{=}} \{(b \rightarrow c) \rightarrow a\}'' \stackrel{2.6}{\underline{=}} (b \rightarrow c) \rightarrow a$, which implies that $\mathcal{A}_{12} \subseteq \mathcal{A}_{13}$.

If $\mathbf{A} \in \mathcal{A}_{13}$ and $a, b, c \in A$, then $(a \rightarrow b) \rightarrow c \stackrel{(A13)}{\underline{=}} (b \rightarrow c) \rightarrow a \stackrel{(A13)}{\underline{=}} (c \rightarrow$
 $a) \rightarrow b \stackrel{2.6}{\underline{=}} (c \rightarrow a)'' \rightarrow b \stackrel{3.13(e)}{\underline{=}} (a \rightarrow c)'' \rightarrow b \stackrel{2.6}{\underline{=}} (a \rightarrow c) \rightarrow b$, concluding that
 $\mathcal{A}_{13} \subseteq \mathcal{A}_{11}$. □

4 Main Theorem

In this section we will prove our main theorem. But first we need one more lemma.

LEMMA 4.1. *If $\mathbf{A} \in \mathcal{A}_2 \cup \mathcal{A}_6 \cup \mathcal{A}_9$ then $\mathbf{A} \in \mathcal{A}_{11}$.*

Proof. We will see that $\mathbf{A} \models (x \rightarrow y)' \approx (y \rightarrow x)'$.

Let $a, b \in A$.

- If $\mathbf{A} \in \mathcal{A}_2$, $(a \rightarrow b) \rightarrow 0 \stackrel{2.3(1) \& 2.6}{\underline{=}} (0' \rightarrow (a \rightarrow b)) \rightarrow 0 \stackrel{(A2)}{\underline{=}} (0' \rightarrow (b \rightarrow a)) \rightarrow$
 $0 \stackrel{2.3(1) \& 2.6}{\underline{=}} (b \rightarrow a) \rightarrow 0$.
- If $\mathbf{A} \in \mathcal{A}_6$, $(a \rightarrow b) \rightarrow 0 \stackrel{2.3(1) \& 2.6}{\underline{=}} (0' \rightarrow (a \rightarrow b)) \rightarrow 0 \stackrel{(A6)}{\underline{=}} (a \rightarrow (b \rightarrow 0')) \rightarrow$
 $0 \stackrel{(A6)}{\underline{=}} (b \rightarrow (0' \rightarrow a)) \rightarrow 0 \stackrel{2.3(1) \& 2.6}{\underline{=}} (b \rightarrow a) \rightarrow 0$.

- If $\mathbf{A} \in \mathcal{A}_9$, then

$$\begin{aligned}
 (a \rightarrow b)' &= (a \rightarrow b) \rightarrow 0 \\
 &= (a \rightarrow (0' \rightarrow b)) \rightarrow 0 && \text{by 2.3 (1) and 2.6} \\
 &= (b \rightarrow (0' \rightarrow a)) \rightarrow 0 && \text{by (A9)} \\
 &= (b \rightarrow a) \rightarrow 0 && \text{by 2.3 (1) and 2.6} \\
 &= (b \rightarrow a)'
 \end{aligned}$$

Now, apply Lemma 3.5, to get $\mathbf{A} \in \mathcal{A}_{12}$. Therefore, using Theorem 3.14, we conclude $\mathbf{A} \in \mathcal{A}_{11}$. \square

We are now ready to present the main theorem of this paper.

THEOREM 4.2. *We have*

- (a) *The following are the 8 subvarieties of \mathcal{I} of associative type that are distinct from each other.*

$$\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3, \mathcal{A}_4, \mathcal{A}_6, \mathcal{A}_9, \mathcal{A}_{11} \text{ and } \mathcal{A}_{14}.$$

- (b) *They satisfy the following relationships:*

1. $\mathcal{SL} \subset \mathcal{A}_3 \subset \mathcal{A}_4$,
2. $\mathcal{BA} \subset \mathcal{A}_4 \subset \mathcal{I}$,
3. $\mathcal{A}_3 \subset \mathcal{A}_1 \subset \mathcal{I}$,
4. $\mathcal{A}_3 \subset \mathcal{A}_2 \subset \mathcal{A}_{11}$, $\mathcal{A}_3 \subset \mathcal{A}_6 \subset \mathcal{A}_{11}$ and $\mathcal{A}_3 \subset \mathcal{A}_9 \subset \mathcal{A}_{11}$,
5. $\mathcal{A}_{11} \subset \mathcal{A}_{14} \subset \mathcal{I}$.

Proof. Observe that, in view of Theorem 3.12 and Theorem 3.14 we can conclude that each of the 14 subvarieties of associative type of \mathcal{I} is equal to one of the following varieties:

$$\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3, \mathcal{A}_4, \mathcal{A}_6, \mathcal{A}_9, \mathcal{A}_{11}, \mathcal{A}_{14}.$$

We first wish to prove (b). Notice that by Lemma 3.11 we have that $\mathbf{A} \in \mathcal{A}_3$ is of type 1. Then, using Theorem 3.7, $\mathbf{A} \in \mathcal{A}_j$ for all $1 \leq j \leq 14$. Hence

$$\mathcal{A}_3 \subseteq \mathcal{A}_j \text{ for all } 1 \leq j \leq 14. \quad (4.1)$$

1. Recall that $\mathcal{SL} = \mathcal{C} \cap I_{1,0}$. Then, we get $\mathcal{C} \subseteq \mathcal{A}_1$ and $\mathcal{I}_{1,0} \subseteq \mathcal{A}_1$ by [7, Theorem 8.2] and [7, Theorem 9.3], respectively, implying $\mathcal{SL} \subseteq \mathcal{A}_3$, and $\mathcal{A}_3 \subseteq \mathcal{A}_4$ by (4.1).

The algebras $\mathbf{2}_z$ and $\mathbf{2}_b$ show that $\mathcal{SL} \neq \mathcal{A}_3$ and $\mathcal{A}_3 \neq \mathcal{A}_4$, respectively.

2. In view of [10] we have that $\mathcal{BA} \subset \mathcal{S}$. By [9, Lemma 3.1], $\mathcal{S} \models x \rightarrow (y \rightarrow z) \approx y \rightarrow (x \rightarrow z)$. Thus, $\mathcal{BA} \subseteq \mathcal{A}_4$.

The algebra $\mathbf{2}_s$ shows that $\mathcal{BA} \neq \mathcal{A}_4$ and the following algebra shows that $\mathcal{A}_4 \neq \mathcal{I}$, respectively.

\rightarrow :	0	1	2
0	0	0	0
1	2	0	0
2	0	0	0

3. The algebra $\mathbf{2}_b$ shows that $\mathcal{A}_1 \neq \mathcal{I}$ and the following algebra witnesses that $\mathcal{A}_3 \neq \mathcal{A}_1$.

\rightarrow :	0	1	2
0	0	1	2
1	1	1	2
2	2	1	2

4. Using (4.1) and Lemma 4.1 we can conclude that $\mathcal{A}_3 \subseteq \mathcal{A}_2 \subseteq \mathcal{A}_{11}$, $\mathcal{A}_3 \subseteq \mathcal{A}_6 \subseteq \mathcal{A}_{11}$ and $\mathcal{A}_3 \subseteq \mathcal{A}_9 \subseteq \mathcal{A}_{11}$.

The following algebras show that $\mathcal{A}_3 \neq \mathcal{A}_2$ and $\mathcal{A}_2 \neq \mathcal{A}_{11}$, respectively.

\rightarrow :	0	1	2
0	0	0	0
1	2	0	2
2	0	0	0

\rightarrow :	0	1	2
0	0	0	0
1	2	0	0
2	0	0	0

The following algebras show that $\mathcal{A}_3 \neq \mathcal{A}_6$ and $\mathcal{A}_6 \neq \mathcal{A}_{11}$, respectively.

\rightarrow :	0	1	2	3
0	0	0	0	0
1	0	2	3	0
2	0	0	0	0
3	0	0	0	0

\rightarrow :	0	1	2
0	0	0	0
1	2	0	0
2	0	0	0

The following algebras show that $\mathcal{A}_3 \neq \mathcal{A}_9$ and $\mathcal{A}_9 \neq \mathcal{A}_{11}$, respectively.

→:	0	1	2	3
0	0	0	0	0
1	0	2	3	0
2	0	0	0	0
3	0	0	0	0

→:	0	1	2
0	0	0	0
1	2	0	0
2	0	0	0

5. Let $\mathbf{A} \in \mathcal{A}_{11}$ and $a, b, c \in A$. By Theorem 3.14, $\mathcal{A}_{11} = \mathcal{A}_{12} = \mathcal{A}_{13}$. Hence $(a \rightarrow b) \rightarrow c \stackrel{(A13)}{=} (b \rightarrow c) \rightarrow a \stackrel{(A12)}{=} (c \rightarrow b) \rightarrow a$. Therefore $\mathcal{A}_{11} \subseteq \mathcal{A}_{14}$.


The algebra $\mathbf{2}_b$ shows that $\mathcal{A}_{14} \neq \mathcal{I}$ and the following algebra shows that

	→:	0	1	2	3
	0	0	1	2	3
$\mathcal{A}_{11} \neq \mathcal{A}_{14}$.	1	2	3	2	3
	2	1	1	3	3
	3	3	3	3	3

The proof of the theorem is now complete since (a) is an immediate consequence of (b). □

Big space?

The Hasse diagram of the poset of subvarieties of \mathcal{I} of associative type, together with \mathcal{SL} and \mathcal{BA} , is:



5 Identities of Associative type in Symmetric Implication Zroupoids

Let $\mathbf{A} \in \mathcal{I}$. \mathbf{A} is involutive if $\mathbf{A} \in \mathcal{I}_{2,0}$. \mathbf{A} is meet-commutative if $\mathbf{A} \in \mathcal{MC}$. \mathbf{A} is symmetric if \mathbf{A} is both involutive and meet-commutative. Let \mathcal{S} denote the variety of symmetric \mathcal{I} -zroupoids. In other words, $\mathcal{S} = \mathcal{I}_{2,0} \cap \mathcal{MC}$. The variety \mathcal{S} was investigated in [7], [9] and [10] and has some interesting properties.

In this section we give an application of the main theorem, Theorem 4.2, to describe the poset of the subvarieties of the variety \mathcal{S} .

LEMMA 5.1. [9, Lemma 3.1 (a)] *Let $\mathbf{A} \in \mathcal{S}$. Then \mathbf{A} satisfies $x \rightarrow (y \rightarrow z) \approx y \rightarrow (x \rightarrow z)$.*

LEMMA 5.2. [9, Lemma 2.1] $\mathcal{MC} \cap \mathcal{I}_{1,0} \subseteq \mathcal{C} \cap \mathcal{I}_{1,0} = \mathcal{SL}$.

LEMMA 5.3. [9, Lemma 3.2] *Let $\mathbf{A} \in \mathcal{S}$ such that $\mathbf{A} \models x \rightarrow x \approx x$. Then $\mathbf{A} \models x' \approx x$.*

LEMMA 5.4. $\mathcal{A}_{11} \cap \mathcal{S} = \mathcal{SL}$.

Proof. Let $\mathbf{A} \in \mathcal{A}_{11} \cap \mathcal{S}$ and $a \in A$. Since $\mathcal{S} \subseteq \mathcal{I}_{2,0}$, we have

$$\begin{aligned} a &= a' \rightarrow a && \text{by Lemma 2.3 (4)} \\ &= (0' \rightarrow a') \rightarrow a && \text{by Lemma 2.3 (1)} \\ &= (0' \rightarrow a) \rightarrow a' && \text{by (A11)} \\ &= a \rightarrow a' && \text{by Lemma 2.3 (1)} \\ &= a'' \rightarrow a' \\ &= a' && \text{by Lemma 2.3 (4)}. \end{aligned}$$

Therefore, $\mathbf{A} \models x \approx x'$. Then, by Lemma 5.2, $\mathbf{A} \in \mathcal{SL}$. \square

LEMMA 5.5. $\mathcal{A}_1 \cap \mathcal{S} \subseteq \mathcal{SL}$.

Proof. Let $\mathbf{A} \in \mathcal{A}_1 \cap \mathcal{S}$ and $a \in A$. Then

$$\begin{aligned} a &= 0' \rightarrow a && \text{by Lemma 2.3 (1)} \\ &= (0 \rightarrow 0) \rightarrow a \\ &= 0 \rightarrow (0 \rightarrow a) && \text{by (A1)} \\ &= 0 \rightarrow a && \text{by Lemma 2.5 (c)}. \end{aligned}$$

Consequently,

$$\mathbf{A} \models x \approx 0 \rightarrow x. \quad (5.1)$$

Therefore,

$$\begin{aligned} a &= a' \rightarrow a && \text{by Lemma 2.3 (4)} \\ &= (a \rightarrow 0) \rightarrow a \\ &= a \rightarrow (0 \rightarrow a) && \text{by (A1)} \\ &= a \rightarrow a && \text{by equation (5.1)} \end{aligned}$$

Thus, by Lemma 5.3, $\mathbf{A} \models x' \approx x$. Using Lemma 5.2 we can conclude the proof. \square

We will denote by \mathcal{S}_i the variety $\mathcal{A}_i \cap \mathcal{S}$ with $1 \leq i \leq 14$.

We will now give an application of the main theorem, Theorem 4.2, of Section ??.

PROPOSITION 5.6. *Each of the 14 subvarieties of associative type of \mathcal{S} is equal to one of the following varieties:*

$$\mathcal{SL}, \mathcal{S}_{14}, \mathcal{S}.$$

Proof. From Theorem 3.12 and Theorem 3.14 we know that each of the 14 subvarieties of associative type of \mathcal{I} is equal to one of the following varieties:

$$\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3, \mathcal{A}_4, \mathcal{A}_6, \mathcal{A}_9, \mathcal{A}_{11}, \mathcal{A}_{14}.$$

Using Theorem 4.2, Lemma 5.4 and Lemma 5.5 we have that

$$\mathcal{S}\mathcal{L} \subseteq \mathcal{S}_3 \subseteq \mathcal{S}_2 \subseteq \mathcal{S}_{11} \subseteq \mathcal{S}\mathcal{L},$$

$$\mathcal{S}\mathcal{L} \subseteq \mathcal{S}_6 \subseteq \mathcal{S}_{11} \subseteq \mathcal{S}\mathcal{L},$$

$$\mathcal{S}\mathcal{L} \subseteq \mathcal{S}_9 \subseteq \mathcal{S}_{11} \subseteq \mathcal{S}\mathcal{L}$$

and

$$\mathcal{S}\mathcal{L} \subseteq \mathcal{S}_1 \subseteq \mathcal{S}\mathcal{L}$$

By Lemma 5.1, $\mathcal{A}_4 = \mathcal{S}$, $\mathcal{S}_0, \mathcal{S}_4 = \mathcal{S}$. □

We are now ready to present the main theorem of this section.

THEOREM 5.7. *We have*

- (a) *The following are the 3 subvarieties of \mathcal{S} of associative type that are distinct from each other.*

$$\mathcal{S}\mathcal{L}, \mathcal{S}_{14}, \mathcal{S}.$$

- (b) *They satisfy the following relationships*

1. $\mathcal{S}\mathcal{L} \subset \mathcal{S}_{14} \subset \mathcal{S}$,
2. $\mathcal{B}\mathcal{A} \not\subset \mathcal{S}_{14}$.

Proof. We first prove (b).

1. By Theorem 4.2, $\mathcal{S}\mathcal{L} \subseteq \mathcal{S}_{14}$.

The following algebras show that $\mathcal{S}\mathcal{L} \neq \mathcal{S}_{14}$ and $\mathcal{S}_{14} \neq \mathcal{S}$, respectively.


$\rightarrow:$	0	1	2	3
0	0	1	2	3
1	2	3	2	3
2	1	1	3	3
3	3	3	3	3

$\rightarrow:$	0	1
0	1	1
1	0	1

2. Since $\mathbf{2}_b \not\models (S_{14})$, it follows that $\mathcal{B}\mathcal{A} \not\subset \mathcal{S}_{14}$.

The proof of the theorem is now complete since (a) is an immediate consequence of Proposition 5.6 and (b). □

The Hasse diagram of the poset of subvarieties of \mathcal{S} of associative type, together with \mathcal{BA} , is:



Acknowledgment: The first author wants to thank the institutional support of CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas).

Compliance with Ethical Standards:

Conflict of Interest: The first author declares that he has no conflict of interest. The second author declares that he has no conflict of interest.

Ethical approval: This article does not contain any studies with human participants or animals performed by any of the authors.

Funding: The work of Juan M. Cornejo was supported by CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas) and Universidad Nacional del Sur. Hanamantagouda P. Sankappanavar did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- [1] J.C. Abbott, *Semi-boolean algebra*, *Matematicki Vesnik* **19**, (1967), 177 – 198.

-
- [2] **R. Balbes, P.H. Dwinger**, *Distributive lattices*, Univ. of Missouri Press, Columbia, 1974.
- [3] **B.A. Bernstein**, *A set of four postulates for Boolean algebras in terms of the implicative operation*, Trans. Amer. Math. Soc. **36** (1934), 876 – 884.
- [4] **S. Burris, H.P. Sankappanavar**, *A Course in universal algebra*, Graduate texts in Mathematics 78, Springer-Verlag, New York, 1981. The free, corrected version (2012) is available online as a PDF file at <http://www.math.uwaterloo.ca/~snburris>.
- [5] **J.M. Cornejo, H.P. Sankappanavar**, *Order in implication zroupoids*. Stud Logica, **104** (2016), no. 3, 417 – 453.
- [6] **J.M. Cornejo, H.P. Sankappanavar**, *Semisimple varieties of implication zroupoids*, Soft Comput. **20** (2016), 3139 – 3151.
- [7] **J.M. Cornejo, H.P. Sankappanavar**, *On implicator groupoids*, Algebra univers. **77** (2017), no. 2, 125 – 146.
- [8] **J.M. Cornejo, H.P. Sankappanavar**, *On derived algebras and subvarieties of implication zroupoids*, Soft Comput. **21** (2017), 6963 – 6982.
- [9] **J.M. Cornejo, H.P. Sankappanavar**, *Symmetric implication zroupoids and the identities of Bol-Moufang type*, Soft Comput. **2017** (published online: 25 October 2017), 1-15. DOI 10.1007/s00500-017-2869-z.
- [10] **J.M. Cornejo, H.P. Sankappanavar**, *Symmetric implication zroupoids and the weak associative identities* (2017). (submitted).
- [11] **J.M. Cornejo, H.P. Sankappanavar**, *Varieties of implication zroupoids*, (2018). (In preparation).
- [12] **F. Fenyves**, *Extra loops II. On loops with identities of Bol-Moufang type*, Publ. Math. Debrecen **16** (1969), 187 – 192 .
- [13] **M. Hosszu**, *Some functional equations related with the associative law*, Publ. Math. Debrecen **3**, (1954), 205 – 214.
- [14] **M.A. Kazim, M. Naseeruddin**, *On almost-semigroups*, The Alig. Bull Math **2**, (1972), 1 – 7.
- [15] **K. Kunen**, *Quasigroups, loops, and associative laws* J. Algebra **185**, (1996), 194 – 204.
- [16] **W. McCune**, *Prover9 and Mace4* <http://www.cs.unm.edu/mccune/prover9/> (2005 – 2010).
- [17] **D.C. Murdoch**, *Structure of abelian quasi-groups*, Trans. Amer. Math. Soc. **49**, (1941), 392 – 409.
- [18] **C.S. Peirce**, *On the algebra of logic*, Amer. J. Math. **3** (1880), 15 – 57.
- [19] **H.O. Pfugfelder**, *Historical notes on loop theory*, Comment. Math. Univ. Carolin. **41**, (2000), no. 2, 359 – 370.
- [20] **J.D. Phillips, P. Vojtechovsky**, *The varieties of loops of Bol-Moufang type*, Algebra univers. **54** (2005), 259 – 271.
- [21] **J.D. Phillips JD, P. Vojtechovsky P**, *The varieties of quasigroups of Bol-Moufang type: an equational reasoning approach*, J. Algebra **293** (2005), 17 – 33.

-
- [22] **P.V. Protic, N. Stevanovic**, *Abel-Grassmann bands*, *Quasigroups and Related Systems* **11** (2004), 95 – 101.
- [23] **D.I. Pushkashu**, *Para-associative groupoids*, *Quasigroups and Related Systems* **18** (2010), 187 – 194.
- [24] **H. Rasiowa**, *An algebraic approach to non-classical logics*, North-Holland Publishing Co., Amsterdam. *Studies in Logic and the Foundations of Mathematics*, **78**, 1974.
- [25] **H.P. Sankappanavar**, *De Morgan algebras: new perspectives and applications*, *Sci. Math. Jpn.* **75** (2012), no. 1, 21 – 50.
- [26] **E. Schroeder**, *Über algorithmen und Calculi*, *Arch. der Math. und Phys.* **5** (1887), 225 – 278.
- [27] **D. Stanovsky**, *A guide to self-distributive quasigroups, or latin quandles*, *Quasigroups and Related Systems* **23** (2015), 91 – 128.
- [28] **S.K. Stein**, *Left-distributive quasigroups*, *Proc. Amer. Math. Soc.* **10** (1959), 577 – 578.
- [29] **S.K. Stein**, *On the foundations of quasigroups*, *Trans Amer. Math Soc.* **85** (1957), 228 – 256.
- [30] **A. Suschkewitsch**, *On a generalization of the associative law*, *Trans. Amer. Math. Soc.* **31** (1929), no. 1, 204 – 214.

JUAN M. CORNEJO
DEPARTAMENTO DE MATEMÁTICA
UNIVERSIDAD NACIONAL DEL SUR
ALEM 1253, BAHÍA BLANCA, ARGENTINA
INMABB - CONICET
E-MAIL: JMCORNEJO@UNS.EDU.AR

HANAMANTAGOUDA P. SANKAPPANAVAR
DEPARTMENT OF MATHEMATICS
STATE UNIVERSITY OF NEW YORK
NEW PALTZ, NEW YORK 12561
U.S.A.
E-MAIL: SANKAPPH@NEWPALTZ.EDU