

Title	Secondary Aortoenteric Erosion Followed by Recurrent Lower Extremity Abscesses
Author(s) Alternative	Ono, S; Samejima, Y; Watada, S; Kakefuda, T
Journal	Annals of vascular surgery, 42(): 302.e1-302.e5
URL	http://hdl.handle.net/10130/5048
Right	
Description	

Secondary aortoenteric erosion followed by recurrent lower extremity abscesses

Shigeshi Ono,¹ Yoshitomo Samejima,¹ Susumu Watada,¹ and Toshihiro Kakefuda¹

¹ Department of Vascular Surgery, Kawasaki Municipal Hospital, Kanagawa, Japan.

Correspondence to: Shigeshi Ono, MD, PhD, Department of Vascular Surgery,

Kawasaki Municipal Hospital, 12-1 Shikawadori, Kawasaki-ku, Kawasaki, Kanagawa,

Japan; telephone: 81-44-233-5521; fax: 81-44-245-9600;

E-mail: shigeshi.ono81@gmail.com

ABSTRACT:

A secondary aortoenteric fistula (SAEF) is a relatively rare complication of aortoiliac reconstructive surgery, often involving relatively fixed duodenal 3rd portion and a vascular anastomosis. We observed a 52-year-old man with a recurrent right lower leg abscess following the erosion between ipsilateral bifurcated prosthetic graft limb and non-fixed jejunum. In situ graft-sparing surgical treatment with aggressive debridement was successfully performed. A SAEF may occur even at a non-vascular anastomosis site, or in non-fixed small bowel, and may become a source of a septic embolus. A high index of suspicion for SAEF is required for early diagnosis and treatment of this life-threatening complication.

A secondary aortoenteric fistula (SAEF) is one of the rare complications of abdominal aortoiliac reconstructive surgery, with a reported incidence of 0.36%-3% (1-7). The clinical and radiological manifestations of SAEF are still unspecified, despite improved knowledge regarding the pathology. However, early diagnosis is essential to manage this life-threatening complication, with high operative mortality rates of 40%-50% (8). A fistula may occur when a prosthetic graft erodes into the bowel (a continuous mechanical contact may be necessary for the incidence). The findings from our case highlight the importance of considering the possibility of SAEF in any patient with a history of aortoiliac reconstructive surgery; even appendicitis may cause adhesions between the free bowel and the paraprosthetic site. The patient consented to the publication of this report.

CASES REPORT

A 52-year-old man was admitted to our hospital with a presentation of fever and swelling of right lower leg. Two-and-a-half years prior to admission, he had undergone an emergent bifurcated graft

replacement (Dacron, 18 x 9 mm) for a ruptured abdominal aortic aneurysm (rAAA). Retroperitonization had been performed at the operation. He had been intermittently plagued with high fever and abdominal pain for one year prior to admission. Six-months prior to admission, he developed an abscess in the right lower leg, accompanied by fever. Broad-spectrum antibiotics were administered, followed by open drainage. The blood and pus cultures were negative for growth at that time. The abscess cured promptly, though its etiology was not identified. At the time, he had one episode of hematochezia, although, no specific lesion was detected on colonoscopy.

On the day of admission, his blood pressure was 154/87 mmHg; pulse rate was 67 beats/min; and his temperature was 38.3°C. His abdomen was soft and flat, and his right leg was swollen and reddish. The laboratory findings were as follows: white blood cell count: 13,130/mm³, hemoglobin: 12.7 g/dl, and C-reactive protein: 6.93 mg/dL. Enhanced CT scanning revealed air density mass around the prosthetic bifurcated graft, adjacent to the jejunum (Fig. 1A), a phlegmonous appendicitis, and a recurrence of the right lower leg abscess (Fig.1B); however, the blood and

the abscess cultures revealed no growth.

He underwent an open laparotomy through a midline incision. On exploration, the tip of the appendix and the jejunum were found clearly adhered to the swollen retroperitoneum (Fig.2A), however, no other adhesion was apparent. The appendix was slightly swollen without perforation, and the jejunum was eroded by the underlying prosthetic graft limb (Fig.2B). The graft limb was bile-stained, and slightly oozed blood (Fig. 2C). The bile stain was localized on the right limb, and it was completely separated from the vascular suture line. There was only a little pus around the graft.

The bile stained graft limb was resected with a safety margin, which was 3 cm long in total, and anastomosed in an end-to-end fashion. Aggressive debridement of the surrounding tissue was performed. Irrigation with 10,000 mL of normal saline was performed, and no irrigation tube was placed. The jejunum was partially resected and anastomosed in an end-to-end fashion, while an appendectomy was performed at the same time. Using vascularized omentum, the graft was covered and the retroperitoneal space was packed to control infection.

The culture from the resected graft grew *Escherichia coli* and *Enterobacter cloacae*, however, the blood culture remained negative for growth. The mucosa of the resected appendix revealed chronic inflammation. The patient's recovery was uneventful. Based on the cultured organisms, CPMX was administered intravenously for 3 weeks, followed by oral LVFX for 8 months. Blood analysis was performed every 3-month, and an enhanced CT was performed every year, with additional CT scanning if there were any signs of inflammation or infection, and no evidence of re-infection was observed over 5 years after the surgery.

DISCUSSION

A secondary aortoenteric fistula (SAEF) is a rare complication of abdominal aortoiliac reconstructive surgery, which may occur after a few days or up to 27 years after the primary surgery (8-10). The operative mortality is high, reported as 40%-50% (2, 7, 8); thus early diagnosis and aggressive treatment are essential to improve the outcomes. This critical condition often presents with unspecific symptoms, such as fever, abdominal pain, and anorexia (8), which makes it difficult to diagnose

correctly and promptly.

Two types of SAEF have been described (11, 12). Type-1 fistula is a true aortoenteric or graft-enteric fistula, developing between the aortic suture line and the bowel. “Herald bleeding”, which is a minor bleeding preceding major hemorrhage from bowel tract, has been observed and reported in many cases (2, 8). Sepsis and abdominal pain are relatively rare with this type of fistula (11, 12).

Type-2 fistula is a paraprosthetic-enteric erosion, with no communication between the bowel and the graft, as in our case. It is known to account for 15%-20% of SAEF (11, 12). In this type of fistula, the bleeding occurs from the edges of the eroded bowel because of the mechanical pulsations of the aortic graft. Sepsis is more frequently associated with this type of fistula (57%) (11, 12). As a clinical presentation, septic emboli in the lower extremities, septic arthritis, multicentric osteomyelitis and hypertrophic osteoarthropathy have been reported, with an incidence rate of 2%-8% (2, 8, 13-16). An infectious clot may form in an eroding prosthetic graft, and become the source of an embolus.

The etiology of SAEFs still remains unclear, but two possible mechanisms have been reported. One is the constant pulsating contact of the graft on the bowel, and the other is an already existing adhesion due to an infection or inflammation (8, 17, 18). However, SAEFs occur even after endovascular aneurysm repair regardless of the absence of aortic suture line and prosthetic graft outside the aortic lumen, which could make it difficult to understand the exact pathogenesis (19). Most type-1 SAEFs have been reported to involve the relatively fixed duodenal 3rd portion and a vascular suture line of the anastomosis (20). In our case, the suture line and the duodenum were not associated with the erosion. The erosion was located between right graft limb and the non-fixed jejunum. We considered the possibility that the chronic appendicitis spread retroperitoneum and caused the adhesions between the jejunum and the lesion, resulting in the origin of constant pulsating contact. It may be possible that the primary surgery for rAAA caused the adhesions between the jejunum and the retroperitoneum, although, the retroperitonization was performed as far as possible at the time of operation. However, that still does not explain why the tip of the appendix penetrated the lesion or

the mucosal inflammation seen in the resected appendix. Moreover, chronic appendicitis would be consistent with the patient's intermittent abdominal pain. Appendicitis involved in SAEF has been reported to be as rare as 3% (8).

Infection control is one of the most important factors in the management of this life-threatening complication. There is no consensus on the optimal duration of antibiotic therapy after the treatment of infected aortic lesions. Parenteral antibiotics are commonly administered for 2 to 8 weeks after surgery; whether lifelong oral antibiotics are needed, is still debated (21-23).

There are several reported operative strategies to treat this severe condition, which are often complex procedures, e.g., graft excision with extra anatomical reconstruction, in situ reconstruction with autogenous vein or cryopreserved allograft or rifampin-soaked grafts, and simultaneous or two-stage surgeries (2, 8, 20, 24, 25). Although extra anatomical reconstruction has improved results, the mortality and aortic stump blowout rate still remain high (26). Replacement of the grafts with biological conduits will most likely remain the best treatment option,

although, there are no universal guidelines on managing prosthetic graft infections (27, 28). Moreover, cryopreserved allograft may not be available for such urgent operations, and it is expensive. Conversely, in situ graft-preserving strategies have also been reported (29-33). Previous reports have described either a prolonged period of irrigation with povidone-iodine, or intermittent packing with povidone-iodine-soaked sponges (32, 33). These techniques require stage-surgical procedures to cover the graft, and povidone-iodine irrigation is limited by its toxicity (29). We resected the bile-stained graft limb with an adequate safety margin and preserved the original graft because we thought the infection was completely localized, and the length of the graft limb was long enough to allow suturing in an end-to-end fashion without a need of even an autogenous vein graft. A single irrigation with normal saline was administered to reduce the residual bacteria count, which was absolutely not toxic and did not require staged therapy to cover the graft.

Furthermore, the omentum has been used for a variety of surgical procedures (34). It can prevent or control aortic graft infection (35), and has been described as an ideal biological drain (30). Our case had a rich

and vascularized greater omentum, and the infection was localized. Our strategy may be a feasible alternative to conventional replacement surgery for such limited cases.

CONCLUSION

The aim of this case report is to emphasize the importance of early diagnosis and treatment of sepsis or lower extremity abscesses in patients who have a history of aortic reconstructive surgery. A SAEF may occur in the prosthetic graft limb even away from the anastomosis, and could be a source of septic embolism. In situ graft-sparing surgical treatment with aggressive debridement may be feasible in the treatment of localized type-2 SAEF.

DISCLOSURE

We all have no conflict of interest to declare.

FIGURE CAPTIONS

Fig.1A The arrowhead shows an air density mass around the prosthetic bifurcated graft limb.

Fig.1B Arrowheads showing the right lower leg abscesses.

Fig.2A The tip of appendix and the jejunum found clearly adhered to swollen retroperitoneum.

Fig.2B The jejunum eroded by the underlying prosthetic graft limb. The arrow shows the jejunal mucosa, and the arrowhead shows the underlying prosthetic graft.

Fig.2C The graft limb is bile-stained and slightly oozing blood. The bile stain is localized, and completely separated from the vascular suture line.

REFEERENCES

1. Baril DT, Carroccio A, Ellozy SH, Palchik E, Sachdev U, Jacobs TS, et al. Evolving strategies for the treatment of aortoenteric fistulas. *Journal of vascular surgery*. 2006 Aug;44(2):250-7. PubMed PMID: 16890849.
2. Batt M, Jean-Baptiste E, O'Connor S, Saint-Lebes B, Feugier P, Patra P, et al. Early and late results of contemporary management of 37 secondary aortoenteric fistulae. *European journal of vascular and endovascular surgery : the official journal of the European Society for Vascular Surgery*. 2011 Jun;41(6):748-57. PubMed PMID: 21414817.
3. Champion MC, Sullivan SN, Coles JC, Goldbach M, Watson WC. Aortoenteric fistula. Incidence, presentation recognition, and management. *Annals of surgery*. 1982 Mar;195(3):314-7. PubMed PMID: 7059240. Pubmed Central PMCID: 1352637.
4. Dorigo W, Pulli R, Azas L, Pratesi G, Innocenti AA, Pratesi C. Early and long-term results of conventional surgical treatment of secondary aorto-enteric fistula. *European journal of vascular and endovascular surgery : the official journal of the European Society for Vascular Surgery*. 2003 Nov;26(5):512-8. PubMed PMID: 14532879.
5. Hallett JW, Jr., Marshall DM, Petterson TM, Gray DT, Bower TC, Cherry KJ, Jr., et al. Graft-related complications after abdominal aortic aneurysm repair: reassurance from a 36-year population-based experience. *Journal of vascular surgery*. 1997 Feb;25(2):277-84; discussion 85-6. PubMed PMID: 9052562.
6. Kieffer E, Sabatier J, Plissonnier D, Knosalla C. Prosthetic graft infection after descending thoracic/ thoracoabdominal aortic aneurysmectomy: management with in situ arterial allografts. *Journal of vascular surgery*. 2001 Apr;33(4):671-8. PubMed PMID: 11296316.
7. Swain TW, 3rd, Calligaro KD, Dougherty MD. Management of infected aortic prosthetic grafts. *Vascular and endovascular surgery*. 2004 Jan-Feb;38(1):75-82. PubMed PMID: 14760481.
8. Bergqvist D, Bjorck M. Secondary arterioenteric fistulation--a systematic literature analysis. *European journal of vascular and endovascular surgery : the official journal of the European Society for Vascular Surgery*. 2009 Jan;37(1):31-42. PubMed PMID: 19004648.

9. Shindo S, Tada Y, Sato O, Idezuki Y, Nobori M, Tanaka N. A case of an aortocolic fistula occurring 27 years after aorto-femoral bypass surgery, treated successfully by surgical management. *Surgery today*. 1993;23(11):993-7. PubMed PMID: 8292869.
10. Bastounis E, Papalambros E, Meringas V, Maltezos C, Diamantis T, Balas P. Secondary aortoduodenal fistulae. *The Journal of cardiovascular surgery*. 1997 Oct;38(5):457-64. PubMed PMID: 9358802.
11. Dachs RJ, Berman J. Aortoenteric fistula. *American family physician*. 1992 Jun;45(6):2610-6. PubMed PMID: 1595511.
12. Mohammadzade MA, Akbar MH. Secondary aortoenteric fistula. *MedGenMed : Medscape general medicine*. 2007;9(3):25. PubMed PMID: 18092031. Pubmed Central PMCID: 2100090.
13. Martin A, Copeman PW. Aorto-jejunal fistula from rupture of Teflon graft, with septic emboli in the skin. *British medical journal*. 1967 Apr 15;2(5545):155-6. PubMed PMID: 5297743. Pubmed Central PMCID: 1841650.
14. O'Mara C, Imbembo AL. Paraprosthetic-enteric fistula. *Surgery*. 1977 May;81(5):556-66. PubMed PMID: 300510.
15. Gordon SL, Nicholas GG, Carter SL, Mira AJ. Aorto-enteric fistula presenting as multicentric osteomyelitis. *Clinical orthopaedics and related research*. 1978 Mar-Apr(131):255-8. PubMed PMID: 657631.
16. Dalinka MK, Reginato AJ, Berkowitz HD, Turner ML, Freundlich B, Steinberg M. Hypertrophic osteoarthropathy as indication of aortic graft infection and aortoenteric fistula. *Archives of surgery*. 1982 Oct;117(10):1355-9. PubMed PMID: 6751288.
17. Barman AA, Kerr P. Primary and secondary aortoenteric fistula and thoracic aortic aneurysm. *New York state journal of medicine*. 1992 Apr;92(4):156-8. PubMed PMID: 1594148.
18. Chang MW, Chan YL, Hsieh HC, Chang SS. Secondary aortoduodenal fistula. *Chang Gung medical journal*. 2002 Sep;25(9):626-30. PubMed PMID: 12479626.
19. Lyden SP, Tanquilut EM, Gavin TJ, Adams JE. Aortoduodenal fistula after abdominal aortic stent graft presenting with extremity abscesses. *Vascular*. 2005 Sep-Oct;13(5):305-8. PubMed PMID:

16288707.

20. Schoell T, Manceau G, Chiche L, Gaudric J, Gibert H, Tresallet C, et al. Surgery for secondary aorto-enteric fistula or erosion (SAEFE) complicating aortic graft replacement: a retrospective analysis of 32 patients with particular focus on digestive management. *World journal of surgery*. 2015 Jan;39(1):283-91. PubMed PMID: 25189446.

21. Kan CD, Lee HL, Yang YJ. Outcome after endovascular stent graft treatment for mycotic aortic aneurysm: a systematic review. *Journal of vascular surgery*. 2007 Nov;46(5):906-12. PubMed PMID: 17905558.

22. Luo CY, Ko WC, Kan CD, Lin PY, Yang YJ. In situ reconstruction of septic aortic pseudoaneurysm due to Salmonella or Streptococcus microbial aortitis: long-term follow-up. *Journal of vascular surgery*. 2003 Nov;38(5):975-82. PubMed PMID: 14603203.

23. Moneta GL, Taylor LM, Jr., Yeager RA, Edwards JM, Nicoloff AD, McConnell DB, et al. Surgical treatment of infected aortic aneurysm. *American journal of surgery*. 1998 May;175(5):396-9. PubMed PMID: 9600286.

24. Luo CY, Lai CH, Wen JS, Lin BW. Secondary aortocolic fistula: case report and review of the literature. *Annals of vascular surgery*. 2010 Feb;24(2):256 e5-12. PubMed PMID: 19892515.

25. Harlander-Locke MP, Harmon LK, Lawrence PF, Oderich GS, McCready RA, Morasch MD, et al. The use of cryopreserved aortoiliac allograft for aortic reconstruction in the United States. *Journal of vascular surgery*. 2014 Mar;59(3):669-74. PubMed PMID: 24239113.

26. Kuestner LM, Reilly LM, Jicha DL, Ehrenfeld WK, Goldstone J, Stoney RJ. Secondary aortoenteric fistula: contemporary outcome with use of extraanatomic bypass and infected graft excision. *Journal of vascular surgery*. 1995 Feb;21(2):184-95; discussion 95-6. PubMed PMID: 7853593.

27. LeMaire SA, Coselli JS. Options for managing infected ascending aortic grafts. *The Journal of thoracic and cardiovascular surgery*. 2007 Oct;134(4):839-43. PubMed PMID: 17903492.

28. Teebken OE, Bisdas T, Assadian O, Ricco JB.

Recommendations for reporting treatment of aortic graft infections. European journal of vascular and endovascular surgery : the official journal of the European Society for Vascular Surgery. 2012 Feb;43(2):174-81. PubMed PMID: 22155121.

29. Umminger J, Krueger H, Beckmann E, Kaufeld T, Fleissner F, Haverich A, et al. Management of early graft infections in the ascending aorta and aortic arch: a comparison between graft replacement and graft preservation techniques. European journal of cardio-thoracic surgery : official journal of the European Association for Cardio-thoracic Surgery. 2016 May 12. PubMed PMID: 27174548.

30. Tossios P, Karatzopoulos A, Tsagakis K, Sapolidis K, Grosomanidis V, Kalogera A, et al. Treatment of infected thoracic aortic prosthetic grafts with the in situ preservation strategy: a review of its history, surgical technique, and results. Heart, lung & circulation. 2014 Jan;23(1):24-31. PubMed PMID: 24103706.

31. Akowuah E, Narayan P, Angelini G, Bryan AJ. Management of prosthetic graft infection after surgery of the thoracic aorta: removal of the prosthetic graft is not necessary. The Journal of thoracic and cardiovascular surgery. 2007 Oct;134(4):1051-2. PubMed PMID: 17903532.

32. Mathes DW, Yaremchuk MJ, Isselbacher EM, Madsen JC. Successful in situ treatment of an infected ascending aortic graft. The Annals of thoracic surgery. 2000 Oct;70(4):1410-2. PubMed PMID: 11081915.

33. Nakajima N, Masuda M, Ichinose M, Ando M. A new method for the treatment of graft infection in the thoracic aorta: in situ preservation. The Annals of thoracic surgery. 1999 Jun;67(6):1994-6; discussion 7-8. PubMed PMID: 10391356.

34. Miller DW, Jr., Johnson DD. Omental pedicle graft in the management of infected ascending aortic prostheses. The Annals of thoracic surgery. 1987 Dec;44(6):614-7. PubMed PMID: 3318738.

35. Goldsmith HS, De los Santos R, Beattie EJ, Jr. Experimental protection of vascular prosthesis by omentum. Archives of surgery. 1968 Dec;97(6):872-8. PubMed PMID: 4882426.