

Neurofibromatosis type-1 with retinal microvascular corkscrew tortuosity

Adrian J Battiston¹, Lauren A Dalvin^{1,2}, Carol L Shields¹

Key words: Corkscrew vessels, neurofibromatosis type-1, retina, vascular tortuosity

A 40-year-old Caucasian woman presented with the history of neurofibromatosis type-1 (NF-1), diagnosed at age 5 years, with cutaneous neurofibromas on the face, back, and chest, and arachnoid cyst. Best-corrected visual acuity was 20/40 in the right eye and 20/20 in the left eye. Anterior segment examination revealed multifocal iris Lisch nodules in both eyes. Fundoscopic examination in the right eye revealed [Fig. 1a] 3 areas of retinal microvascular tortuosity, including superonasal to the optic disc (white arrows, 6.0 mm by 3.0 mm), in the papillomacular bundle (yellow arrows 1.5 mm by 1.0 mm), and inferior to the macula (blue arrows, 2.0 by 2.0 mm). In the left eye, (b) there was 1 area of microvascular tortuosity in the papillomacular bundle (yellow arrows, 2.0 × 2.0 mm). Retinal microvascular tortuosity was confirmed on fundus autofluorescence imaging in the right (c) and left (d) eyes




Figure 1: Fundoscopic examination in the right eye revealed (a) 3 areas of retinal microvascular tortuosity, including superonasal to the optic disc (white arrows, 6.0 mm by 3.0 mm), in the papillomacular bundle (yellow arrows 1.5 mm by 1.0 mm), and inferior to the macula (blue arrows, 2.0 by 2.0 mm). In the left eye, (b) there was 1 area of microvascular tortuosity in the papillomacular bundle (yellow arrows, 2.0 × 2.0 mm). Retinal microvascular tortuosity was confirmed on fundus autofluorescence imaging in the right (c) and left (d) eyes

Access this article online	
Quick Response Code:	Website: www.ijo.in
	DOI: 10.4103/ijo.IJO_1796_19

¹Ocular Oncology Service, Wills Eye Hospital, Thomas Jefferson University, 840 Walnut Street, 14th Floor, Philadelphia, PA, ²Department of Ophthalmology, Mayo Clinic, Rochester, MN, USA

Correspondence to: Dr. Carol L Shields, Ocular Oncology Service, 840 Walnut Street, Suite 1440, Philadelphia - 19107, PA, USA. E-mail: carolshields@gmail.com

Received: 27-Sep-2019

Revision: 01-Nov-2019

Accepted: 06-Nov-2019

Published: 22-Nov-2019

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Battiston AJ, Dalvin LA, Shields CL. Neurofibromatosis type-1 with retinal microvascular corkscrew tortuosity. *Indian J Ophthalmol* 2019;67:2054-5.




Figure 2: In the right eye, areas of retinal microvascular tortuosity were visible (A) superonasal to the optic disc (white arrows) and (B) inferior to the macula (blue arrows). (C) In the left eye, retinal microvascular tortuosity was observed in the papillomacular bundle (yellow arrows)

left eye [Fig. 1b], there was 1 area of microvascular tortuosity in the papillomacular bundle (yellow arrows, 2.0 × 2.0 mm). Retinal microvascular tortuosity was confirmed on fundus autofluorescence imaging in the right [Fig. 1c] and left [Fig. 1d] eyes.

By fluorescein angiography, areas of retinal microvascular tortuosity were visible [Fig. 2] in each eye, without evidence of leakage, staining, or neovascularization.

Underlying the microvascular abnormalities were multiple hypopigmented choroidal lesions, suggestive of hamartoma, nevus, or neurofibroma. On OCT (vertical cut), these corresponded to [Fig. 3] areas of increased choroidal density in the outer choroid (white arrow). Based on the benign retinal and choroidal findings, observation was advised.

Retinal microvascular findings in NF-1 are likely congenital, representing defective migration of vasomotor nerve cells from the neural crest.^[1-3] These benign abnormalities remain stable over time without retinal ischemia, exudation, hemorrhage, vascular leakage, or visual acuity loss.^[1-4] An analysis of 34 NF-1 patients




Figure 3: Underlying the microvascular abnormalities were multiple hypopigmented choroidal lesions, suggestive of hamartoma, nevus, or neurofibroma. On OCT (vertical cut), these corresponded to [Figure 3] areas of increased choroidal density in the outer choroid (white arrow)

demonstrated retinal microvasculature abnormalities (6/34) and choroidal nodules (23/34),^[4] while a larger cohort found no correlation of microvascular changes (18/294) with choroidal nodules (186/294) ($P = 0.48$).^[3] Our case exemplifies the subtle nature of microvascular abnormalities with underlying choroidal nodules in NF-1.

Support provided in part by the Eye Tumor Research Foundation, Philadelphia, PA (CLS), an unrestricted grant from Research to Prevent Blindness, Inc (LAD), and the Heed Ophthalmic Foundation (LAD). The funders had no role in the design and conduct of the study, in the collection, analysis, and interpretation of the data, and in the preparation, review, or approval of the manuscript. Carol L. Shields, M.D. has had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. The authors have no conflicts of interest. All authors have read and approved the manuscript.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient has given his consent for his images and other clinical information to be reported in the journal. The patient understands that his name and initials will not be published and due efforts will be made to conceal his identity, but anonymity cannot be guaranteed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- Abdolrahimzadeh S, Felli L, Piraino DC, Mollo R, Calvieri S, Recupero SM. Retinal microvascular abnormalities overlying choroidal nodules in neurofibromatosis type 1. *BMC Ophthalmol* 2014;14:146.
- Muci-Mendoza R, Ramella M, Fuenmayor-Rivera D. Corkscrew retinal vessels in neurofibromatosis type 1: Report of 12 cases. *Br J Ophthalmol* 2002;86:282-4.
- Parrozzani R, Pilotto E, Clementi M, Frizziero L, Leonardi F, Convento E, *et al.* Retinal vascular abnormalities in a large cohort of patients affected by neurofibromatosis type 1: A study using optical coherence tomography angiography. *Retina* 2018;38:585-93.
- Cassiman C, Casteels I, Stalmans P, Legius E, Jacob J. Optical coherence tomography angiography of retinal microvascular changes overlying choroidal nodules in neurofibromatosis type 1. *Case Rep Ophthalmol* 2017;8:214-20.