



Plant eco-physiological response patterns to summer drought, elevated CO₂ and warming in a natural temperate heath ecosystem

Ecological Society of America, 93rd ESA annual meeting August 3-8, Milwaukee, Wisconsin

Albert, Kristian; N. Mikkelsen, Teis; Ro-Poulsen, Helge; Michelsen, Anders

Published in:

93rd ESA Annual Meeting Sunday, August 3 -Friday, August 8, 2008. The Midwest Airlines Center - Milwaukee, Wisconsin

Publication date:

2008

Document version

Publisher's PDF, also known as Version of record

Citation for published version (APA):

Albert, K., N. Mikkelsen, T., Ro-Poulsen, H., & Michelsen, A. (2008). Plant eco-physiological response patterns to summer drought, elevated CO₂ and warming in a natural temperate heath ecosystem: Ecological Society of America, 93rd ESA annual meeting August 3-8, Milwaukee, Wisconsin. In *93rd ESA Annual Meeting Sunday, August 3 -Friday, August 8, 2008. The Midwest Airlines Center - Milwaukee, Wisconsin*

Plant eco-physiological response patterns to summer drought, elevated CO₂ and warming in a semi-natural temperate heath ecosystem

Kristian Rost Albert¹, **Teis N. Mikkelsen**², **Helge Ro-Poulsen**¹ and **Anders Michelsen**¹, (1)Department of Biology, University of Copenhagen, Øster Farimagsgade 2D, 1353-Copenhagen K, Denmark, (2)Biosystems Department, Risø National Laboratory for Sustainable Energy, Technical University of Denmark, BIO-309, P.O. Box 49, Frederiksborgvej 399, Roskilde, DK-4000, Denmark

Background/Question/Methods

Plant eco-physiological responses to multiple environmental changes are being studied in a temperate semi-natural heath ecosystem according to a realistic Danish climatic scenario anno 2075. Since direction of ecosystem responses can not be deduced from single factor experiments alone this necessitates the multi-factor approach. The environmental treatments are elevated level of CO₂ (FACE 510 ppm, [CO₂]), passive nighttime warming (IR-reflective curtains, [T]) and summer drought (rain activated curtains, [D]) and started in Oct. 2005. Leaf level carbon input and water consumption through photosynthesis were measured by gas exchange techniques on Common Heather (*Calluna vulgaris*) and Hair Grass (*Deschampsia flexuosa*). Parallel measurements of chlorophyll-flourescence, xylem water potential, leaf carbon and nitrogen content were conducted.

Results/Conclusions

In summer 2007, [D] decreased plant available soil water led to significantly lower plant water potential compared to controls, whereas [CO₂] and [T] had no effect. The plant water potential was 2-3 times lower in the deep rooted woody shrub *C. vulgaris* compared to the grass *D. flexuosa*. No visual symptoms of drought stress were seen for *C. vulgaris*, but part of *D. flexuosa* had senescent leaves. For both species no effects on maximal carboxylation velocity, V_{cmax} , and maximum rate of electron transport, J_{max} , were seen, but for *D. flexuosa* [D] increased day time respiration, R_{d} , was found. Despite small impact on photosynthetic parameters V_{cmax} and J_{max} we observed a significantly decreased maximal photosynthesis in [D] for both species. Under field conditions the transpiration rates were significantly lowered by [D], but [CO₂] and [T] had no effect. Interactions between [D*CO₂] and [T*D*CO₂] showed lower transpiration rates than expected whereas [T*CO₂] were higher for *C. vulgaris*. Parallel responses were seen for values of stomatal conductance. These responses indicate the strong impact of [D] linking low water availability to

decreased plant water potential and water consumption via transpiration. Further, the [D] also decreased the net photosynthesis for both species, while [CO₂] had the opposite effect. In combination this led to a significantly higher Water Use Efficiency in [CO₂]. The increased carbon uptake increased the leaf C/N ratio through a marginal, but significant, C% decrease and a stronger N% decrease in [CO₂]. These issues of the contrasting response patterns could not be deduced from single factor studies alone and adds to the importance of long term multifactor studies.
