



**Borassus akeassii Bayton, Ouédraogo & Guinko sp. nov.**

Sacande, Moctar

*Published in:*  
Seed Leaflet

*Publication date:*  
2007

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Sacande, M. (2007). *Borassus akeassii* Bayton, Ouédraogo & Guinko sp. nov. *Seed Leaflet*, (121).



# SEED LEAFLET

No. 121 September 2007



## *Borassus akeassii* Bayton, Ouédraogo & Guinko sp. nov.

*Borassus akeassii* sp. nov. (Arecaceae) is described as a new species from western Burkina Faso, West Africa. It has been confused with the widely distributed African species *Borassus aethiopum* and more recently with the Asian *Borassus flabellifer*.

### Taxonomy and nomenclature

**Family:** Arecaceae

**Vernacular/common names:** African fan palm, ron palm, elephant palm, deleb palm, black rhun palm. Local names: sebe (Bambara) and koanga (Mooré, Burkina).

**Related species of interest:** *Borassus aethiopum*, whose petioles are yellow and the fruits are yellow-orange when ripe.

### Distribution and habitat

*Borassus akeassii* is indigenous and restricted to West Africa; being found in semi-arid and sub-humid zones in Benin, Burkina Faso, Cote d'Ivoire, Mali, Niger and Nigeria. It is also recorded in Senegal and the Central African Republic. The range of *B. akeassii* may be much wider than suggested, owing to its misidentification as *B. aethiopum*. It is found in Sudanian and Sudano-Guinean savannahs with 800–1100 mm annual precipitation, often in close proximity to human settlements. In Burkina Faso, it is the common palm in the western provinces. Vast areas of land are populated with *B. akeassii*, with food and cash crops growing beneath the palms. *Borassus akeassii* shows a preference for well-drained soils and is fire-resistant.

The species has developed a root foraging strategy, which enables it to exploit sparse nutrient-rich patches. The palm is slow growing and very long lived, to over 100 years old. Little is known about its ecology outside the sphere of human influence and there is a lack of accurate information regarding the population size and geographical range of the species. There is a risk of over-exploitation, although in Burkina Faso the species is abundant, but where it is also a fully protected species. Populations of *B. akeassii* are threatened by over-exploitation in Senegal and Guinea.

### Uses

*Borassus akeassii* provides a large number of useful products and is exploited in several ways, though wine production is the most significant of these. The



*Borassus akeassii* crown with hanging mature green fruits. Photo: M Sacande.

apical bud is hollowed out and the resulting sap is collected and fermented. The wine is made for sale (enhancing local earnings), to drink at traditional ceremonies, as well as forming a part of the balanced diet in many rural areas, and used for aphrodisiac preparations. Excessive wine tapping kills the plant. Both the fruit pulp and seeds are edible. Endosperm from the seeds of immature fruits is harvested and consumed. Almost all parts of *B. akeassii* are used, producing food, oils, timber, dyes, fibre, wine, and raw materials (leaves) for mats and baskets. The dark brown, strong, coarsely fibrous wood is a highly prized timber. The wood is very resistant to termites and fungi, and is used in carpentry, construction, bridges and gutters, and also for household articles. The roots serve for the treatment of stomach parasites, sore throats and asthma, as well as being used for a mouthwash

### Botanical description

*Borassus akeassii* is distinguished by its glaucous, green leaves with weakly armed petioles and a characteristic pattern of lamina venation. The flowers of the pistillate inflorescence are arranged in three spirals. It is a dioecious, pleonanthic (does not die after flowering) palm, whose straight dark grey trunk attains 40–50 cm in diameter and 15 m tall, with prominent leaf scars. Mature trees have between 10 and 30 living leaves in crown, though leaf harvesting is common. The petioles are rounded abaxially and flattened; 90–160 cm long, 3–7.4 cm wide at the junction with the lamina, vivid green in colour, with sharp edges or minute spines 0.2–0.6 cm long.

The male inflorescence is 0.8 to 1.2 m long, with 3-6 partial inflorescences that are ca. 50 cm long. The female inflorescence is usually unbranched, with larger flowers of 3 cm in diameter. Females typically bear 50-100 fruits in a fruiting season.

### Fruit and seed description

**Fruit:** The ovoid to ellipsoid fruits are about 15 cm long and 12 cm wide, though variation owing to the number of pyrenes in the fruit; mesocarp yellow and fragrant, with longitudinal fibres in the pulp. They have a pointed apex and are greenish when ripe. The fibrous fruit pulp is yellow to white and slightly oily. There are 1-3 pyrenes of 7–9 x 5–8 cm per fruit, and there is one seed in each pyrene, with hard white endosperm, seed almost filling endocarp.

**Seed:** The seed handling unit is the pyrene which can attain up to 10 cm in diameter and weigh about 100 g each. The morphological seeds are bilobed, pointed and basally attached, with an apical embryo of about 4 x 3 mm, and a homogeneous and centrally hollow endosperm

### Flowering and fruiting habit

The reproductive period of the plant, before death, is short, an average of 22 years. Flowering usually occurs in the second part of the rainy season. Seeds are dispersed by mammals, such as chimpanzees.

### Processing and handling

Fruits are usually harvested in November till May. The pericarp is then removed by hand-slashing to release the pyrenes.

### Storage and viability

Seeds have a short viability and should be sown as soon as they are removed from the fruit pulp.



Germinating *B. akeassii* seed. Photo: M. Sacande.

### Sowing and germination

Seeds do not require a pre-treatment. Seeds germinate after about 4 weeks. Germination is remote-tubular. The seedlings produce a very long taproot, which can be 1 m deep, while the leaf is only 1 cm high, therefore if possible they should be sown directly at the location where they are required. Growth is very slow.

### Selected readings

**Arbonnier, M. 2004.** Trees, shrubs and lianas of West African dry zones. CIRAD, Montpellier; Museum national d'histoire naturelle, Paris.

**Barot, S. & Gignoux, J. 1999.** Population Structure and Life Cycle of *Borassus aethiopum* Mart. Evidence of Early Senescence in a Palm Tree. *Biotropica*, 31 (3), pp. 439-448.

**Bayton, P.R., Ouedraogo, A. & Guinko, S. 2006.** The genus *Borassus* (Arecaceae) in West Africa, with a description of a new species from Burkina Faso. *Botanical Journal of the Linnean Society*, (150), pp. 419–427.

THIS NOTE WAS PREPARED IN COLLABORATION WITH CENTRE NATIONAL DE SEMENCES FORESTIÈRES BURKINA FASO

Author: Moctar Sacande

Millennium Seed Bank project	Phone: +44-1444 894100
Wakehurst Place, Ardingly	Fax: +44-1444 894110
West Sussex	Email: msbsci@kew.org
RH17 6TN, UK	Website: www.kew.org/msbp

Seedleaflets are a series of species wise extension leaflets for tropical forest species with special emphasis on seed technology. Leaflets are compiled from existing literature and research available at the time of writing. In order to currently improve recommendations, FLD encourage feedback from users and researchers who have experience with the species. Comments, corrections, improvements and amendments will be incorporated into future edited leaflets. Please write your comments to: SL-International@life.ku.dk