



Expression and Binding Properties of CBM45 from *Solanum tuberosum*

Baumann, M. J.; Andersen, J.M.; Glaring, Mikkel Andreas; Nakai, H.; Hachem, M. Abou; Blennow, Andreas; Svensson, B.

Publication date:
2008

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Baumann, M. J., Andersen, J. M., Glaring, M. A., Nakai, H., Hachem, M. A., Blennow, A., & Svensson, B. (2008). *Expression and Binding Properties of CBM45 from Solanum tuberosum*. Abstract from XXVI Carbohydrate Symposium, Oslo, Norway.



8th

Carbohydrate Bioengineering
Meeting

PROGRAM AND ABSTRACTS

May 10-13, 2009 Ischia, Naples, Italy

Expression and binding properties of CBM45 from *Solanum tuberosum*

**M.I. Baumann¹, J.M. Andersen¹, M.A. Glaring², H. Nakai¹, M. Abou Hachem¹,
A. Blennow², B. Svensson¹**

¹Enzyme and Protein Chemistry, Department of Systems Biology, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

²VKR Research Centre Pro-Active Plants, Department of Plant Biology and Biotechnology, Faculty of Life Sciences, University of Copenhagen, 1870 Frederiksberg, Denmark

Starch binding domains (SBDs) have been found in CBM families 20, 21, 25, 26, 34, 41, 45, 48 and 53 ¹, http://www.cazy.org/fam/acc_CBM.html. Recently a new SBD family, CBM45 was discovered. ², this family only contains sequences of the plant kingdom. CBM45 occur as an N-terminal tandem domain in only two classes of intracellular enzymes, plastidial amylases (EC 3.2.1.1) and glucan water dikinases (GWDs, EC 2.7.9.4). GWD1 is expressed in photosynthetic organisms and phosphorylates transient and storage starch in the plastids stimulating starch mobilisation. Plastidial α -amylases are presumably involved in the starch turnover, even though the exact role is currently unclear.

In order to probe the domain borders multiple single CBM45 constructs of GWD1 from *Solanum tuberosum* (potato) were made and expressed as a cleavable His-fusion protein in *E. coli*. Protein integrity was tested after removal of the His-tag by differential scanning calorimetry (DSC).

Even though CBM45s do not contain cysteines the internal CBM, CBM45-2, is unexpectedly thermostable (T_m of 84°C). Binding properties of the CBM45 were tested by isothermal calorimetry (ITC) and surface plasmon resonance (Biacore) using soluble low molecular weight starch mimetic motifs. The rather low affinity as compared to most other starch binding domains suggests reversible or regulated binding of CBM45-containing enzymes. Similar constructs were fused with YFP to probe the ability of the isolated CBMs *in vivo*. Analyses of tobacco leaves transiently expressing CBM45-YFP-fusions by confocal microscopy suggest *in vivo* binding of the CBM45 to transient starch granules.

This work has been supported by an H. C. Ørsted Postdoctoral fellowship from DTU (MJB), the Carlsberg Foundation, a Novo Student Scholarship (JMA) and a grant of the Danish Research Council for Technology and Production (MAG).

References

1. M. Machovic and S. Janecek, *Cell. Mol. Life Sci.*, **63**, 2710 (2006)
2. R. Mikkelsen, K. Suszkiewicz, A. Blennow, *Biochemistry*, **14**, 4674 (2006)