



Plant community responses to climate change

Nielsen, Jane Kongstad

Publication date:
2012

Document version
Publisher's PDF, also known as Version of record


Citation for published version (APA):
Nielsen, J. K. (2012). *Plant community responses to climate change*. Forest & Landscape, University of Copenhagen. Forest and landscape research, No. 48/2012



Plant community responses to climate change

FOREST & LANDSCAPE RESEARCH

48 / 2012



Jane Kongstad





FOREST & LANDSCAPE

Plant community responses to climate change

Jane Kongstad



Forest & Landscape Research is issued by Forest & Landscape Denmark which is a national centre for research, education and advisory services within the fields of forest and forest products, landscape architecture and landscape management, urban planning and urban design.

The journal accepts Ph.D. theses, D.Sc. theses, and other major research reports of scientific standard concerning forest, park, landscape, and planning research written in Danish or English.

The content of the journal undergoes a scientific peer-review process.

Forest & Landscape Research is to be considered the continuation of Forskningsserien - The Research Series (ISSN: 1398-3423).

Editorial board:

Niels Elers Koch (editor-in-chief), director, professor, Forest & Landscape Denmark

Frank Søndergaard Jensen (associate editor), senior researcher, Forest & Landscape Denmark

Tilde Tvedt (associate editor), senior consultant, Forest & Landscape Denmark

J. Bo Larsen, professor, Forest & Landscape Denmark

Jørgen Primdahl, professor, Forest & Landscape Denmark

Erik Dahl Kjær, professor, Forest & Landscape Denmark

Title: Plant community responses to climate change

Series-title, no.: Forest & Landscape Research, No. 48-2012

Author: Jane Kongstad

Citation: Kongstad, J. (2012): Plant community responses to climate change. Forest & Landscape Research No. 48-2012. Forest & Landscape Denmark, Frederiksberg. 155 pp.

ISBN: 978-87-7903-575-1 (paper)
978-87-7903-576-8 (internet)

ISSN: 1601-6734

Printed by: Prinfo Aalborg, DK

Number printed: 30

Order: Single issues are available from Forest & Landscape Denmark - see last page. Also published at www.sl.life.ku.dk.



ENGLISH SUMMARY

Climate change is expected to affect terrestrial ecosystems across the globe with increased atmospheric CO₂ concentration, higher temperatures and changes in precipitation patterns. These environmental factors are drivers of many important ecosystem processes, and changes in ecosystem function are therefore expected in the future. The aim of this PhD-thesis was to examine the effects of climate change on aboveground plant growth, plant composition and plant phenology in Danish heathland ecosystems. Two sites were investigated in large-scale field experiments: 1) the CLIMAITE site, 'Brandbjerg' and 2) the INCREASE site at Mols. Field manipulations lasted years and included: Warming, summer drought and (CLIMAITE only) elevated CO₂ concentrations. The treatments were applied individually and in all possible combinations. Further, at Brandbjerg, but outside the treatment plots, a study was performed on the effects nitrogen and phosphorus addition on phenology, chemistry and growth of the dominant grass *Deschampsia flexuosa* (Wavy Hairgrass).

In general, the aboveground vegetation responded less than expected to changing climatic conditions; even though *Calluna vulgaris* (Heather) increased in biomass over the study period, the biomass was not affected by the manipulations, indicating that *C. vulgaris*, has a strong resistance to changes in climate. Also, the grass biomass (primarily *D. flexuosa*) was not affected and was relatively constant over the period. I argue that the resilience of *D. flexuosa* towards the climatic treatments came from the plants ability to let the tissue die back, and then quickly recover once conditions again became favourable. That gave the plant a high resilience to changes in climatic factors. *Calluna vulgaris*, on the other hand, showed a resistance to changes by constantly maintaining the growth during the whole season, probably because of its evergreen status. Together, the two different strategies made the heathland ecosystem more resilient to the climatic treatments than expected. We also found that the amount of flowering culms of *D. flexuosa* increased in response to increased CO₂, whereas the seed germination success decreased. The bryophyte biomass and the nitrogen content decreased in response to nitrogen addition. Even such apparently minor changes might, given time, affect the plant composition and thereby possibly also the major ecosystem processes. Further, we observed changes in the aboveground plant composition in response to the climate manipulations at the Mols site, where *C. vulgaris* was regenerating after a disturbance. Here a decrease in biomass of the pioneer stage was seen, when subjected to the drought treatment compared to warmed and control treatments. I therefore conclude, that the stage of the *C. vulgaris* population as well as the magnitude and frequency of disturbances determine the effects of future climate change on the plant community in heathland ecosystems.

DANSK RESUMÉ

Fremtidens klimaforandringer vil ændre forholdene for vores økosystemer. Højere atmosfæriske CO₂-koncentrationer, varmere klima og ændrede nedbørsmønstre er alle parametre, der vil påvirke biologiske processer og dermed påvirke vores økosystemer på flere niveauer. Formålet med denne ph.d.-afhandling er at belyse effekterne af klimaforandringer på plantevækst, artssammensætning og fænologi i danske hedeøkosystemer. To lokaliteter blev undersøgt med storskala feltstudier: 1) CLIMAITE-lokaliteten, Brandbjerg og INCREASE-lokaliteten på Mols. Behandlinger i eksperimentet var: forhøjet temperatur, forlænget sommertørke samt (kun på CLIMATE lokaliteten) forhøjet atmosfærisk CO₂-koncentration. Alle behandlinger blev undersøgt enkeltvis og i samtlige mulige kombinationer. Desuden blev der på Brandbjerg, men uden for behandlingsfelterne, udført et gødskningforsøg, der skulle undersøge effekten af øget nitrogen- og fosfortilførsel på fænologi, blad-kemi og vækst af *Deschampsia flexuosa* (Bølget Bunke).

Generelt blev den overjordiske vegetation påvirket i mindre grad af klimabehandlingerne end forventet. Selvom *Calluna vulgaris* (Alm. Hedelyng) forøgede sin biomasse hen over perioden, var der ingen effekt af behandlingerne. Dette indikerer, at *C. vulgaris* var mere modstandsdygtigt over for ændringer i klimaet end forventet. Græsbiomassen af *D. flexuosa* blev heller ikke påvirket af klimabehandlingerne, og biomassen var forholdsvis ens alle årene. Jeg argumenterer for, at *D. flexuosas* modstandsdygtighed skyldes dens evne til at visne ned under en tørkeperiode og hurtigt skyde igen så snart vilkårene igen blev favorable. Denne evne til at visne ned og genskyde gjorde arten overlevelsedygtig under de ændrede klimaforhold. Lyngen visnede ikke på samme måde ned. Til gengæld kunne arten, formodentligt på grund af at den er stedsegrøn, opretholde en konstant, lav vækst gennem hele sæsonen, uanset klimabehandling. Tilsammen gjorde disse to plantestrategier heden som økosystem langt mere modstandsdygtigt over for forandringer i klimaet end forventet. Antallet af blomsterstande for *D. flexuosa* steg under forhøjet CO₂-koncentration, mens spiringsevnen faldt. Nitrogenkoncentrationer i bryofytterne og bryofyt-biomassen faldt ved nitrogentilførsel. Selv sådan tilsyneladende små forandringer kan, med tiden, påvirke plantesamfundet og dermed muligvis også de væsentlige biologiske processer i økosystemet. Desuden viste eksperimentet på Mols, at regenerationen af *C. vulgaris*, efter et lyngbladbille angreb, blev påvirket af klimabehandlingerne. Her havde unge skud af *C. vulgaris* sværere ved at reetablere sig i felter, der havde været udsat for tørke, end i felter der enten var kontrolfelter eller varmfelter. Jeg konkluderer derfor, at både livs-stadiet af *C. vulgaris*-populationen samt omfanget og frekvensen af forstyrrelser er med til at bestemme effekten af fremtidens klimaforandringer på plantesamfundet i et hedeøkosystem.

PREFACE

This Ph.D. thesis is submitted to the Faculty of Life Sciences, University of Copenhagen, Denmark. The Ph.D. was a part of the Danish climate change experiment “CLIMAITE” (www.climate.dk) and the work was conducted at the Institute of Forest & Landscape Denmark. From September 2007 to March 2008 I visited the Division of Plant Sciences at the University of Tasmania.

The study was financed by the project CLIMAITE (CLIMATE change effects on biological processes In Terrestrial Ecosystems) funded by The Vilum Kann Rasmussen Foundation, the EU infrastructure INCREASE and by the research school REFOLANA.

I would like to thank several people, who have all contributed to make this thesis possible. First of all I would like to thank my supervisor Inger Kappel Schmidt, Forest & Landscape Denmark, for support during the whole period both in the field and during the writing process, but also for giving me the chance to teach half a year at the Forest College in Nødebo. Thanks also to Claus Beier for supervision and encouragement during the writing of this thesis.

I would also like to thank the CLIMAITE group for good meetings with high scientific outcome; especially I would like to thank the Ph.D. group for lot of good inspiring meetings, talks and fun. A group like that is unique and I felt very lucky to be part of it.

Thanks to the people at division 3 (now 34) at Forest & Landscape Denmark for six good years and of course a special thanks to my office mates; Marie, Jesper and Shimon and my fellow Ph.D. colleagues Andy and Lisbeth -a lot of things have been easier due to your help and support; practical as well as mental.

Special thanks goes to Mark Hovenden, his wife and two children for your hospitality when Lasse and I were half a year in Tasmania. We had a wonderful time at the university as well as outside.

I would like to thank the Forest College for my half year there. Thanks Trine and Michael for a good time and for a lot of inspiration.

Finally, I am very grateful for having a husband like Lasse who is now an expert on heathland ecosystems. Besides support with the thesis, he has been a fantastic listener and inspirer.

TABLE OF CONTENTS

ENGLISH SUMMARY	3
DANSK RESUMÉ.....	4
PREFACE.....	5
TABLE OF CONTENTS.....	6
THE AIM OF THE PROJECT	8
BACKGROUND	9
CLIMATE CHANGE SCENARIOS.....	9
DRY HEATHLANDS.....	9
CLIMATE CHANGE EFFECTS ON ABOVEGROUND VEGETATION	11
CHANGES IN SPECIES COMPOSITION	13
HYPOTHESES	14
METHODS.....	15
STUDY SITE	15
EXPERIMENTAL SETUP	16
VEGETATION ANALYSES	20
RESULTS AND DISCUSSION	23
WHY CLIMATE CHANGE EXPERIMENTS?.....	23
EFFECTS OF CLIMATE CHANGE ON VEGETATION AT CLIMAITE	23
RESILIENCE TO DISTURBANCES	35
CARBON SEQUESTRATION	36
HEATHLANDS IN A FUTURE CLIMATE.....	37
LAND-USE AND FRAGMENTATION	37
DISTRIBUTION REGIMES	38
NITROGEN DEPOSITION	39
CONCLUSION	40
FURTHER RESEARCH.....	41
REFERENCES.....	43
LIST OF PAPERS	53

THE AIM OF THE PROJECT

During the writing of these papers, one question remained in focus again and again. **Will changes in climate alter plant communities over time?** This question may seem simple and easy to answer, but as the following synthesis will hopefully show, simple questions are not always easy to answer – especially not when working with natural ecosystems.

Changes in the climate will affect most biological processes and thereby influence both natural, semi-natural and cultural ecosystems. The challenge of coping with climate changes will increase in the future, and knowledge about the magnitude of the effects on ecosystems is therefore needed.

The aim of the study was to examine the effects of climate changes on aboveground plant growth, plant composition and plant phenology in a Danish heathland.

The thesis consists of data from two projects; the Danish Climate Centre of Excellence, CLIMAITE and the EU infrastructure INCREASE both investigating ecosystem responses to climate in large scale field experiments. Further, four scientific papers are included, all concerned with the relationship between plant growth and climate change. The first paper investigates the effect of climate change on biomass production within a heathland ecosystem. The second paper is about growth and nutrient allocation within the plants in response to nutrient addition. It also deals with the interaction between climate and nutrient addition on the dominant grass species *Deschampsia flexuosa*. The third paper report changes in flower phenology brought about by climate change, based on studies of flowering and seed production of the grass *D. flexuosa*. Finally, the fourth paper is about regeneration of *Calluna vulgaris* after a heather beetle attack under warmer and drier climate conditions.

As stated earlier, my thesis is part of the large climate change experiment CLIMAITE and therefore a part of a network of research covering most biological patterns and processes within the investigated heathland; belowground processes, gas exchanges, plant physiological parameters and plant responses both below- and aboveground. We have all been working at the same study site, with the common key words “climate change” and “heathland ecosystem”. This thesis focuses on the aboveground vegetation, but due to the unique opportunity of knowledge sharing, it is possible to discuss my findings within the findings of others PhD.-students and researchers at the site, and I thereby seek to understand and describe the plant responses in an ecosystem perspective. Finally, the results found in this thesis are discussed in relation to other threats identified for heathland ecosystems including nitrogen deposition, lack of management and land-use changes.

BACKGROUND

Climate change scenarios

Global atmospheric concentrations of greenhouse gasses have increased over the last century, primarily as a result of human activity. The carbon dioxide concentration has increased from a pre-industrial level of 270 ppm to 380 ppm in 2005 (Christensen et al. 2007). This increase mainly owes to fossil fuel burning and land use changes and the CO₂ concentration is expected to increase even further depending on the magnitude of future CO₂ emissions.

Several models have been developed to estimate the magnitude of future changes. Although the models show somewhat different output CO₂ concentrations, there is nevertheless a consensus that the increase in CO₂ will have consequences for other climatic parameters including increased temperature and changes in precipitation patterns with more heavy rain events and longer drought periods. Wind directions and wind speeds are also expected to change with more storms and hurricanes as a result. Most models agree that changes in climate will not be distributed evenly around the world; some areas will be flooded and others may experience severe drought.

In Denmark the temperature is expected to increase with yearly average temperatures 2-3 °C higher in year 2100 compared to 1990. Night temperatures are expected to increase more than day temperatures and winter temperatures will increase more than summer temperatures (Danish Meteorological Institute [http:// www.dmi.dk](http://www.dmi.dk)) (fig 1). Also, precipitation patterns are expected to change in the future. In Denmark, winter precipitation is expected to increase by 20-40 %. In summer, on the other hand, a reduction of 85-90% of current precipitation is expected. Together with a higher frequency of heavy rain falls, the reduction is expected to result in longer drought periods during the growing season (Danish Meteorological Institute [http:// www.dmi.dk](http://www.dmi.dk)) (fig. 2).

Dry Heathlands

All studies within this thesis took place in dry inland heathlands. Heathlands are associated with dwarf shrub dominance, in Denmark mainly the evergreen heather; *Calluna vulgaris*. The ecosystem is characterised by a low level of plant-available nutrients and there is often a strong competition for nutrients between plants, fungi and bacteria (Jonasson et al. 1996). Further it is, at least periodically, limited by water. The diversity of higher plants is low, and the vegetation is adapted to cope with stressful conditions. The system depends on continuous management, since nutrients have to be removed to maintain the low levels.




Figure 1 Predicted changes in temperature ($^{\circ}\text{C}$) in winter (top) and summer (bottom) in the period 2071-2100 compared to the period 1961-1990 in Denmark (from www.DMI.dk).




Figure 2 Predicted changes (%) in winter (left) and summer (right) precipitation in the period 2071-2100 compared to the period 1961-1990 (from www.DMI.dk).

The need for management has increased progressively with the industrialization since the nutrient load into the system has increased markedly during this period. The increased nutrient load together with the lack of management has brought about changes in the plant species composition of heathlands. First of all by invasion of grasses, but also by invasion of trees and bushes, all resulting in higher decomposition rates, changes in soil structure and exclusion of previously present species (Terry et al. 2004). Climate changes could enhance these changes further by speeding up the processes.



Deschampsia flexuosa

Calluna vulgaris

Climate change effects on aboveground vegetation

Increased CO₂ directly stimulate plant growth due to a higher CO₂ assimilation rate (de Graaff et al. 2006) and it has also been shown to increase the number of flowers or seeds per plant (Jablonski et al. 2002; Thurig et al. 2003). Further, increased carbon assimilation may lead to higher rhizodeposition (Zak et al. 1993) and thereby stimulate belowground activity and the mineralization rate. However, nutrient and water limitation has been shown to quench the CO₂-induced biomass increase, resulting in a more limited response in natural ecosystems compared to agricultural systems (Leakey et al. 2009). Studies on natural ecosystems that have reported increased biomass production under elevated CO₂, concluded that this increase mainly was due to an indirect effect on the hydrological cycle, because elevated CO₂ decreases the stomatal conductance, leading to improved water use efficiency (WUE) (Ainsworth and Long 2005).

Increased CO₂ has, on the other hand, been shown to increase C/N and C/P ratios in litter, which may decrease the mineralization rate (van Heerwaarden et al. 2005; Hovenden et al. 2008). A decreased mineralization rate could result in an increased immobilization of N and P, and thus a reduction in the plant-available N/P pools. However, since CO₂ is expected to

increase the plant biomass production and thereby also the litter production this could compensate for the lower litter quality.

Since *C. vulgaris* is abundant from the North of Norway down to Spain and Portugal, a direct effect of warming should probably not be expected within the short time scale of this study. However, warming may influence the abundance of *C. vulgaris* indirectly, for instance by influencing the number of heather beetles, since they are stimulated by a warm and dry spring (Penuelas et al. 2004). Further, warming has been demonstrated to stimulate N-mineralization and thereby increase N-availability (Aerts et al. 2006; Emmett et al. 2004; Rustad et al. 2001). Thus, warming may therefore increase the aboveground biomass of especially the grass, since it is a weaker competitor for nutrients compared to the heather under nutrient deficient conditions.

Lately, monitoring of the Danish heathlands has shown that the cover of the grass *D. flexuosa* is now relatively high in many of the Danish heathlands (Andersen et al. 2005). Grass invasion on heathlands has been linked to the relatively high levels of N-deposition (Barker et al. 2004; Terry et al. 2004). If future warming increase the N-availability at the site, the grass could increase in cover at the expense of the slow growing *C. vulgaris*. The study site is at the 55°53' N, and at this latitude, warming also prolongs the growing season (Cleland et al. 2006; Mikkelsen et al. 2008) and thereby increases the biomass production in spring and autumn. Also, spring phenology, such as leaf appearance and flowering time, has been shown to shift forward in response to a warmer climate, due to an earlier start of the growth season (Hovenden et al. 2008; Menzel et al. 2006).

Drought has been shown to decrease plant biomass due to a slow down of most biological processes including photosynthesis (Gordon et al. 1999; Penuelas et al. 2007) mineralization, nutrient cycling and biomass production (Emmett et al. 2004; Schmidt et al. 2004; Larsen et al. 2011). The fraction of biomass found as litter has been shown to increase in response to severe drought (Kongstad et al. 2011), whereas actual litter production may decrease due to lower biomass production (Penuelas et al. 2004). Further, it has been reported that repeated droughts can change soil compactness and water holding capacity even in wet ecosystems (Sowerby et al. 2008). In semi-natural and nutrient poor ecosystems such as heathlands, such alterations can change the plant production and as a result also change plant-plant interactions (Damgaard 1999; Penuelas et al. 2007). However, heathland ecosystems are characterised by a low level of plant-available nutrients and the ecosystem is, at least periodically, limited by water. The vegetation is therefore adapted to cope with stressful conditions, and studies with heathlands similar to ours have recently shown an unexpected resistance towards changes in climate conditions including prolonged drought periods (Grime et al. 2008; Hudson and Henry 2010).

The climate change factors will interact. Recent studies on plant growth, phenology and plant composition have shown that changes in the main climate drivers, individually as well as interactions between increased CO₂ and precipitation and temperature, have a significant impact on ecosystem functions (Shaw et al. 2002; Beier 2004; Norby and Luo 2004; Luo et al. 2008). A higher CO₂-assimilation rate theoretically increases WUE and CO₂ enrichment was thus expected to counterbalance any drought effects by stimulating plant growth and thus also plant nutrient demands (Ainsworth and Long 2005). Drought may, as described above, decrease the biomass production in mid-summer and thereby counteract the increased biomass production expected from increased CO₂ concentration and from warming. Further, since this ecosystem periodically was limited by water, warming might add to the effect of drought, during drought periods in mid-summer (Beierkuhnlein et al. 2011; Penuelas et al. 2007).

Changes in species composition

The distribution of the North Western heathlands is highly linked to the distribution of the key species *C. vulgaris*. Heathlands are also native habitats for the grass *D. flexuosa* and the two species typically co-occur with high cover of the grass after disturbances such as fire or clearances resulting in nutrient releases. Lately, *D. flexuosa* has increased in both cover and area of distribution in Denmark (Nielsen et al. 2011). The increase is assumed to be associated with previously high levels of nitrogen deposition and to the lack of management in nutrient poor areas. *D. flexuosa* and *C. vulgaris* belong to two different functional groups, graminoid and dwarf shrub respectively, and have highly different life strategies, with the grass being faster growing, more productive and having a lower nutrient turn-over time. *Deschampsia flexuosa* has AM- mycorrhiza which facilitates the uptake of especially phosphorus (Smith and Read 2008). *Calluna vulgaris* is, on the other hand, evergreen with a long leaf life span, it has a high content of complex compounds in tissue that make the decomposition rate of the litter low. It has the ericoid mycorrhiza that provides nutrient uptake in organic form (Cornelissen et al. 2001). *Calluna vulgaris* has the so-called ‘s-strategy’ (Grime 2002); it is adapted to nutrient and water limitation, but on the other hand fails to compete when changes lead to a more productive ecosystem (Friedrich et al. 2011).

If warming increases mineralisation and increased CO₂ concentrations increases root exudation, altered climate conditions may induce changes in the plant community similar to nitrogen deposition. For instance, changes in climate conditions may, as seen in the case of increased nitrogen availability, speed up the life cycle of *C. vulgaris*. Invasion by grasses often occur in openings of the *C. vulgaris* canopy, where *D. flexuosa* capture the water and

quickly create a close litter layer that prevents *C. vulgaris* seedlings to establish (Aerts 1993). Changes in climate therefore may result in an even faster spreading of *D. flexuosa* into heathlands.

Grasses have been shown to respond more quickly to changes compared to heather (Aerts 1995; Chapin and Shaver 1996; Arft et al. 1999; Michelsen et al. 1999; Graglia et al. 2001) and the positive effect on biomass production in response to elevated CO₂ may therefore be more pronounced for the grass than for the heather. On the other hand, heather may be more tolerant than the grass towards drought and would suffer less from the higher C/N ratio in litter, due to the ericoid mycorrhiza that provides nutrient uptake in organic form (Cornelissen et al. 2001).

Hypotheses

Based on the above-mentioned I hypothesized that:

- Elevated CO₂ and warming would stimulate aboveground biomass production, but drought would decrease biomass.
- Drought would increase the fraction of litter, but due to the lower production of biomass, the litter pool would decrease in response to drought.
- The amount of flowering culms would increase under elevated CO₂, whereas drought would lead to higher mortality before reaching the state of flowering.
- The main climate drivers were expected to interact. Drought was expected to counterbalance the stimulation of both biomass and flowering culms by elevated CO₂.
- During drought episodes, warming was expected to enhance the intensity of the drought.
- Within the time scale of this thesis, I further expected the grass, *D. flexuosa* to be more responsive to the climate drivers and the nitrogen addition than the heather, *C. vulgaris* due to differences in life growth forms.

METHODS

Study site

CLIMAITE

The experimental site for paper I-III is situated at Brandbjerg, Denmark (55°53'N; 11°58'E) on a hilly, nutrient poor, sandy moraine from the Weichsel glaciation surrounded by areas of elevated sea bed. The site is a heathland/grassland community co-dominated by the perennial grass *Deschampsia flexuosa* and the evergreen dwarf shrub *Calluna vulgaris*. The soil consists of 70 % sand, 20 % coarse sand, 6 % silt and 2 % clay. The pH_{CaCl₂} is around 4.2 in the organic layer and 3.5 in the upper 10 cm of the mineral soil. The vegetation comprises 17 species of vascular plants, 10 mosses and 9 lichens. Plant species are listed in table 1. Pretreatment analyses in 2004 showed an aboveground biomass of approximately 700 g DW m⁻², where *C. vulgaris* accounted for 40 %, *D. flexuosa* 32 %, mosses 26 % and herbs and other grasses only comprised 2 %. At the same time *D. flexuosa* covered 76%, *C. vulgaris* 41 %, mosses 7%, herbs 1% and lichens < 1%. The vegetation can be classified as high light demanding, with relatively little need for nutrients (Ellenberg 1991). The annual bulk nitrogen deposition in 2006-2007 was ~13 kg N ha⁻¹ year⁻¹ measured at the location (Larsen et al. 2011). The study site was fenced to exclude larger herbivores.

Table 1 Species list and cover of species (%) in ambient plots in the study period of 5 years at the CLIMAITE site.

	2004	2006	2007	2008	2009	2010
<i>C. vulgaris</i>	41	55	67	58	63	65
<i>D. flexuosa</i>	76	56	66	42	42	38
<i>F. ovina</i>	0.7	0.1	3	1	2	0
<i>S. decumbens</i>	2	2	2	0.3	0	0.3
<i>A. stricta</i>	0.6	0	0	0	0.3	0
<i>C. arenaria</i>	3	2	1	0.3	0	0
<i>R. acetosella</i>	0.3	0	0	0	0	0
<i>H. cupressiforme</i>	1	2	3	2	2	-
<i>P. schreberi</i>	5	8	6	13	13	-
<i>D. scoparium</i>	0.6	0.3	0.3	1	1	-
<i>Brachythecium sp</i>	0	0	0.3	0	0	-

MOLS

The experimental site for paper IV is located at Mols, Denmark (56°23' N, 10°57'E) and is generally similar to the Brandbjerg site. It is part of the inter-European research projects VULCAN “Vulnerability assessment of shrubland ecosystems in Europe under climatic changes” (Beier 2004) and the European network of large scale climate change experimental sites IN-

CREASE (www.increase-infrastructure.eu). The site is a semi-natural heathland/grassland ecosystem subjected to low-intensity grazing until 1992 and with no further management activities prior to the start of the experiment in 1999. The soil at Mols is a sandy podzol with a shallow organic layer. Also here is the vegetation co-dominated by the evergreen ericaceous shrub *C. vulgaris* (45 %) and the perennial grass *D. flexuosa* (45 %) with low abundances of other grasses, herbs and mosses associated with acidic heathland/grassland. When the experiment was initiated in 1999 the aboveground biomass was approx. 1050 g DW m⁻².

Experimental setup

CLIMAITE - A CLIMATE CHANGE EXPERIMENT

The CLIMAITE experiment was set up to study the **climate change effects on biological processes in terrestrial ecosystems**. It was initiated in 2004 and the experimental treatment was initiated one year later. The manipulations were chosen to match the climate scenarios for Denmark in the year 2075; Increased CO₂ concentration, warmer climate, and changes in precipitation patterns. However, we had one important exception: precipitation is predicted to change with prolonged summer drought and increased winter precipitation. The CLIMAITE experiment focused on the summer drought only, because eventual responses would be difficult to interpret in a combined summer removal and winter addition scenario. Fig. 3 shows a schematic view of the plots. The treatments were: Untreated control (A), CO₂-enriched with a target concentration of 510 ppm (CO₂), increased temperature of 1°C (T) and prolonged drought period of 4-6 weeks during spring/summer (D). Drought treatment was alleviated if soil water content was about 5% in the top 20 cm soil (Table 2). The treatments were applied alone and in all possible combinations: temperature×drought (TD), temperature×CO₂ (TCO₂), drought×CO₂ (DCO₂), and temperature×drought×CO₂ (TDCO₂) replicated in 6 blocks. In each block, the 4 treatments with or without CO₂ were grouped in an octagon in a split-plot design. Each plot was 9.1 m². CO₂ was enriched by FACE (free air carbon enrichment) as described by others (Miglietta et al. 2001). Passive night-time warming increased the air temperature 1 °C by covering the vegetation from dusk to dawn with reflective curtains all year around except during rain events. In the drought period curtains automatically covered the vegetation during rain events. The curtains were activated by a rain sensor and retracted again as soon as the rain stopped. The drought treatment removed 95% of the precipitation during drought period (Table 2). The rest of the year, the curtains were inactive and the plots exposed to control conditions. Soil water contents were measured by TDR probes, air temperature and precipitation were




Figure 3 Schematic presentation of a block at the CLIMAITE site with two separate octagons, ambient CO_2 and receiving CO_2 respectively, together hosting all eight treatment combinations with CO_2 (CO_2), warming (T) drought (D) and untreated control (A) (Redrawn from Mikkelsen (2008))

measured by two weather stations located at the site (Table 2). For further technical details see (Mikkelsen et al. 2008).

Within each plot, two permanently marked vegetation subplots were placed with the size of 0.5 x 0.5 m. All CLIMAITE data collected in this thesis took place in these subplots. The vegetation plots remained untouched over the whole study period, and only non-destructive analyses were performed here.

FERTILIZER EXPERIMENT

To examine the nutrient status of the heathland and to investigate the plant responses to increased nitrogen and phosphorus addition, a fertilizer experiment was setup outside the CLIMAITE plots. Fertiliser was applied in plots of 1.2×1.2 m as three levels of nitrogen, 0, 25 (N) and 75 (NN) $\text{kg N ha}^{-1}\text{yr}^{-1}$ and two levels of phosphorus, 0 and 10 (P) $\text{kg P ha}^{-1}\text{yr}^{-1}$ as well as two combinations of nitrogen and phosphorus (NP) and (NNP) in a 6×6 factorial block design with six replicates. The nutrients were dissolved in 2 l distilled water and added monthly from April to June 2005 as NH_4Cl and $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$. The plots without nitrogen and phosphorus addition were control (C) plots, to which 2 l of distilled water were added.

THE MOLS EXPERIMENT

The field scale climate treatments were initiated in 1999 after one pre-treatment year in order to identify variability between plots (Beier et al. 2004). Three replicated blocks with field-scale night-time warming and extended summer drought treatments and an un-treated control were installed in 1998 and treatments were initiated spring 1999. A light-weight scaffold-

Table 2. Soil minimum and maximum temperatures for April and July are measured in the depth of 5 cm. Precipitation (mm) is an average of two weather stations in 2 m height at the site. Water excluded by the drought in mm and in % of annual precipitation is based on data from the two weather stations. SWC in volume % measured by TDR probes in 0-20 cm depth. See Mikkelsen et al. (2008) for technical details. Data from 2010 was not available from the site and was therefore acquired from the Danish Meteorological Institute measured in the region (www.DMI.dk).

	A	D	I	TD	CO ₂	DCO ₂	ICO ₂	IDCO ₂
2006								
<i>Drought period: 3/7- 21/7 and 26/7 - 8/8</i>								
Min temp Apr (°C)	4.5	4.5	5.0	4.8	4.6	4.6	5.1	5.0
Max temp Apr (°C)	7.1	7.1	8.0	7.9	7.3	7.0	7.1	7.2
Min temp July (°C)	15.4	15.2	16.2	15.7	15.4	15.2	15.7	15.6
Max temp July (°C)	20.7	20.6	21.7	22.1	21.0	21.0	20.2	20.8
Precipitation (mm yr ⁻¹)	629	577	629	577	629	577	629	577
Precipitation excluded (mm)	-	5 and 47	-	5 and 47	-	5 and 47	-	5 and 47
Precipitation excluded (% of annual)	-	1 and 7	-	1 and 7	-	1 and 7	-	1 and 7
Annual mean SWC in depth (Vol%)	15.73	14.54	14.55	13.57	15.19	15.65	14.44	13.99
2007								
<i>Drought period: 21/5- 22/6</i>								
Min temp Apr (°C)	6.4	6.4	7.3	7.0	6.4	6.6	7.1	7.1
Max temp Apr (°C)	9.8	9.6	10.7	10.8	10.3	9.8	10.1	10.2
Min temp July (°C)	15.2	14.8	15.3	15.1	15.2	15.0	15.3	15.1
Max temp July (°C)	17.5	17.7	19.4	17.8	18.2	17.9	17.4	17.5
Precipitation (mm yr ⁻¹)	852	758	852	758	852	758	852	758
Precipitation excluded (mm)	-	94	-	94	-	94	-	94
Precipitation excluded (% of annual)	-	11	-	11	-	11	-	11
Annual mean SWC in depth (Vol%)	17.31	15.63	15.96	14.88	16.37	16.81	15.72	14.96

2008												
<i>Drought period 5/5- 27/5 and 16/9- 2/10</i>												
Min temp Apr (°C)	5.4	5.4	6.1	5.9	5.5	5.6	6.1	6.1	6.1	6.1	6.1	6.1
Max temp Apr (°C)	8.0	7.8	8.8	8.8	8.3	8.0	8.2	8.2	8.2	8.2	8.2	8.5
Min temp July (°C)	14.2	14.1	14.9	14.6	14.2	14.1	14.4	14.4	14.4	14.4	14.4	14.4
Max temp July (°C)	18.0	17.7	18.8	19.2	18.0	17.6	17.4	17.4	17.4	17.4	17.4	17.7
Precipitation (mm yr ⁻¹)	632	597	632	597	632	597	632	632	632	632	632	597
Precipitation excluded (mm)	-	12 and 23	-	12 and 23	-	12 and 23	-	12 and 23	-	12 and 23	-	12 and 23
Precipitation excluded (% of annual)	-	2 and 4	-	2 and 4	-	2 and 4	-	2 and 4	-	2 and 4	-	2 and 4
Annual mean SWC in depth (Vol%)	16.56	14.83	14.44	13.87	15.12	15.77	14.28	14.28	14.28	14.28	14.28	14.16
2009												
<i>Drought period: 18/5 -25/5 and 25/6 – 13/7</i>												
Min temp Apr (°C)	5.5	5.6	6.3	6.0	5.4	5.5	6.0	6.0	6.0	6.0	6.0	6.0
Max temp Apr (°C)	8.8	8.6	9.7	10.0	8.5	8.4	8.49	8.49	8.49	8.49	8.49	8.8
Min temp July (°C)	14.8	14.6	15.3	14.9	14.6	14.4	14.7	14.7	14.7	14.7	14.7	14.7
Max temp July (°C)	17.8	17.5	19.2	18.9	17.4	17.3	17.0	17.0	17.0	17.0	17.0	17.3
Precipitation (mm yr ⁻¹)	668	625	668	625	668	625	668	668	668	668	668	625
Precipitation excluded (mm)	-	18 and 25	-	18 and 25	-	18 and 25	-	18 and 25	-	18 and 25	-	18 and 25
Precipitation excluded (% of annual)	-	3 and 4	-	3 and 4	-	3 and 4	-	3 and 4	-	3 and 4	-	3 and 4
Annual mean SWC in depth (Vol%)	15.47	13.86	13.71	12.90	14.02	14.41	13.87	13.87	13.87	13.87	13.87	13.29
2010												
<i>Drought period: 4/5- 3/6</i>												
Min temp Apr (°C)	4.4	4.6	5.1	4.9	4.6	4.7	5.1	5.1	5.1	5.1	5.1	4.9
Max temp Apr (°C)	7.6	7.8	9.2	8.3	7.4	7.3	7.4	7.4	7.4	7.4	7.4	8.0
Min temp July (°C)	14.5	14.5	15.3	15.1	14.6	14.5	14.8	14.8	14.8	14.8	14.8	14.9
Max temp July (°C)	19.1	19.9	22.3	20.9	18.7	19.2	19.0	19.0	19.0	19.0	19.0	19.9
Precipitation (mm yr ⁻¹)	844	771	844	771	844	771	844	844	844	844	844	771
Precipitation excluded (mm)	-	73	-	73	-	73	-	73	-	73	-	73
Precipitation excluded (% of annual)	-	9	-	9	-	9	-	9	-	9	-	9
Annual mean SWC in depth (Vol%)	17.43	14.82	15.55	14.61	15.38	17.40	16.29	16.29	16.29	16.29	16.29	16.25

ing was placed on each 4 x 5 m plot to carry the roof. The drought plots were subjected to a 1-2 months drought period in the spring/summer from 1999 to 2010 by covering the vegetation with transparent PVC roofs but only during rain events to avoid influences on wind, temperature and light conditions. As at the CLIMAITE the site the roofs were retracted again as soon as the rain stopped. The curtains removed up to 95 % of incoming rain during the drought which equals 20 % (11-29%) of the annual precipitation. The temperature treatment was designed to mimic an increased minimum temperature (night-time warming) rather than a general temperature increase. The warming plots were covered with a reflective aluminium curtain approximately 20 cm above the vegetation. The curtains reflect the major part of the infrared (IR) radiation (Beier et al. 2004). The curtains were controlled by a light sensor and automatically drawn over the vegetation to reduce the loss of IR radiation at dusk, and at sunrise the curtains were retracted to leave the plots open during the day. Further, a rain sensor over-ruled the night-time warming and the curtains were retracted during rain events to avoid major impact on the hydrological cycle. The curtains increased the mean temperature by 0.4°C in the air and by 1.2°C in the soil. The moderate increase in mean temperature increased the growing degree days by 112 % and decreased the number of days with frost by 44%. For further information on the field site and the experimental design, see (Beier et al. 2004).

Vegetation analyses

All the applied methods are described in detail in each paper, but in the following section I outline some of the considerations regarding the use of some of the different methods. Not all the methods that were applied during the work presented in this thesis are included, and those that are included are not described in detail – only some considerations and the rationale for choosing the methods that we did.

PIN-POINT METHOD

To measure plant cover, vegetation height and compactness we used the non-destructive pin-point analysis (Jonasson 1988), as described in more detail in paper I and paper IV. The pin-point method allows conversion of the data into biomass estimates. When using such a model there will always be advantages and disadvantages. The advantages are first of all that the method is a non-destructive method. This was a necessity in this study, since the aim was to follow the vegetation responses to climate change over a period of 1-5 years and the plots were far too small to do actual harvesting. Secondly, the method is relatively time-saving in the field, and to cover all 48 plots at

the site took about 2-3 weeks. However, data processing also takes time, and all together time will also be a disadvantage of the pin-point method since it only allows 1-3 analyses within a year, and therefore can only provide snapshots. Further 2-3 weeks is too long time when recording phenological patterns such as leaf appearance and flowering. The most important disadvantage, however, is the uncertainty of the model; the estimates are based on measurements from outside the CLIMAITE plots, and the model does not take into account differences between treatments that are not related to changes in cover or density. This could be thicker leaves, longer or thicker stems etc. Further, the method is useful in this ecosystem with only a few dominant species. It would be difficult to use in an ecosystem with a higher plant biodiversity. However, I do consider this method not only to be the best available option, but also overall a good tool for consistently estimating plant biomasses in this heathland ecosystem, with only a few dominant species.

PLANT PHENOLOGY

All measurements and observations related to plant phenology are time consuming. Counting flowers, observing flowering stages, collecting seeds and germination experiments all take time. Further, there is a discrepancy between the fact that flowering observations optimally should be performed every day or every second day and of the size of our experimental site (8 treatments, 6 replicates and 2 sub-plots), which resulted in a labour intensive period during flowering. Additionally, the phenological pattern turned out to be very different from one year to the next, and this made it difficult to plan before the flowering started. These difficulties caused some compromises, including the fact that flowering in 2008 should ideally have been followed more closely, since the poor temporal resolution in the end blurred any differences in flowering time between the treatments. Also, the study of flowering development was only done in one season, and the same was true for seed weight and seed germination studies. Thus, despite the shortcomings of the data, I do believe we managed to cover at least some of the phenological patterns of *D. flexuosa*. As a supplement to the pin-point method I, each year (2007-2009) in September, harvested 3-5 *C. vulgaris* shoots in a 10 cm buffer zone around the permanent vegetation plots. I fractionated the material into leaves, flowers and wood and weighted the fractions to identify any changes in allocation of resources into flowering-, leaf- or woody tissue.

NDVI

NDVI (Normalized Difference Vegetation Index) measurements were done using a spectrometer with four sensors for simultaneous measurements of both intrinsic and reflected light within the PAR and NIR bands. The method was applied because we expected to identify phenological patterns such as leaf appearance, flowering time, and biomass peaks. The method is simple

and a good supplement to the time consuming observation method mentioned above. The NDVI turned out to be most efficient during spring when green grass vegetation did not fully cover the area. At that time, I could measure clear differences in leaf appearance (*D. flexuosa*) between warmed and non-warmed plots (Fig 4). Later during the season I observed a point of saturation and the method could not handle the dense vegetation cover. In order to apply the method to determine flowering time and leaf senescence, the method therefore has to be developed or automated cameras for daily image collection could be applied. Further, the phenology measurements have to be performed very often and access to both the instrument and to the field site should be easy.




Figure 4 Difference in NDVI (Normalized Difference Vegetations Index) in warmed plots (1 °C) compared to ambient plots at the CLIMATE site. Measured in spring 2007. *= $p < 0.05$.



Vegetation plot

Pin-point frame

NDVI measurements

RESULTS AND DISCUSSION

Why climate change experiments?

Models are a useful tool in ecology; they can predict changes in future climate conditions and predict future responses in ecosystems due to changes in climate. But all models need input (data), and that is (one reason) why we need experiments. Experiments, besides providing knowledge about ecosystem processes and functions, feed models (Beier 2004). Experiments, both in laboratories and in the field, with effects of single climate factors on ecosystems have been carried out for the last decades. However, interactions between the main drivers of climate change are not necessarily simply additive (Shaw et al. 2002). Interactions can also be synergistic or antagonistic (Larsen et al. 2011), implying that conclusions based on single factor experiments could be insufficient, when trying to predict responses to climate change. Interactions are very difficult, if not impossible, to predict to some degree of certainty unless we have multi-factorial experiments to demonstrate any synergies or antagonisms. Multifactorial experiments have of course many disadvantages; they are often costly and time consuming, and in a global context they will always be case studies (Norby and Luo 2004). However, multi-factorial experiments are crucial in our attempt to understand ecosystem responses to a changing climate. They are needed to unravel the complexity of interactions between the main climate components, and to understand the responses on species level as well as on ecosystems level. Further, they are much needed if we are to identify site- and inter-annual variations, when attempting to extrapolate and upscale. To meet these requirements we first of all need experiments around the world and further we need the experiments to last for more than a couple of years.

Effects of climate change on vegetation at CLIMAITE

In general the aboveground vegetation responded less than I had expected; even though *C. vulgaris* increased in biomass over the study period, the biomass was not affected by the manipulations indicating that *C. vulgaris*, have a strong resistance to changes in climate determined by its evergreen status. Also, the grass biomass was not affected and was relatively constant over the period. The lack of response for *D. flexuosa* to the climatic treatments came from the plants ability to let the tissue dieback and then quickly recover once conditions again become favourable. It therefore showed a high resilience to changes in climate factors. Together the two strategies made the heathland ecosystem more resistant to the climatic treatments than I had expected.

In the following sections I describe the effects found in this thesis, and consider them in context with some of the many other findings from the study site.

ENHANCED CO₂

The photosynthetic rate measured at the site increased for both species in response to enhanced CO₂ concentration (Albert et al. 2011a; 2011c). However, this increase in carbon assimilation was not converted into above-ground biomass (paper I), since I saw no treatment response in biomass for neither *C. vulgaris* nor *D. flexuosa* (fig 5) except for an increased biomass production (green biomass and litter) in 2008 for *D. flexuosa*.

The question was therefore; where was the extra carbon then allocated to? Some of it was apparently allocated into grass reproduction since I saw an increase in the amount of flowering culms in response to increased CO₂ (paper III) (fig 6). But will this increase in flowering result in changes in the species composition? Since flowering is important for spreading and maintenance of the population, I consider an increased reproduction success as an indicator of future growth. However, we saw a negative effect on the germination success, indicating a lower seed quality under elevated CO₂ concentrations which has also been shown in other plants by (Andalo et al. 1996). It has been hypothesised that seed nitrogen content rather than seed weight determines germination success (Hara and Toriyama 1998; Miyagi et al. 2007; Hikosaka et al. 2011) and since there is abundant evidence that nitrogen concentrations of seeds decreases due CO₂ fertilisation (Jablonski et al. 2002; Thurig et al. 2003) this may be the reason for the lower germination in this study. Further the large inter annual variations in flowering, as described below, must also be taken into account before making conclusion on future spreading from seeds.

Allocation to reproduction also increased for *C. vulgaris*, the fraction of biomass from flowers was higher under elevated CO₂ compared to ambient CO₂. This was mainly at the expense of a decrease in the fraction of leaf biomass at that time. However, this pattern was only seen in one year (fig 7). *D. flexuosa* may also increase in cover by vegetative growth, and overall I therefore concluded that changes to the flowering pattern alone are not enough to positively identify changes in future interactions between *D. flexuosa* and *C. vulgaris*.

Apart from reproductive tissue, the excess carbon was also allocated belowground since increases in root biomass (Arndal et al submitted), microbial biomass (von Oheimb et al. 2009) and enchytraeids biomass was reported from the site in response to higher CO₂ concentration (Maraldo and Holmstrup 2009a; Andresen et al. 2010b). This indicated that carbon from




Figure 5 A) Biomass of *D. flexuosa* (g m⁻², mean±SE), B) biomass of *C. vulgaris* (g m⁻², mean±SE) Ambient CO₂ (White) and elevated CO₂, 510 ppm (grey). Litter (hatched) and alive (unhatched) material. * =p<0.05.

the increased photosynthetic assimilation was transported from the leaves and excreted as root exudates to stimulate the microbial activity. It was thus quickly incorporated into belowground biomass or the food web. This was further supported by both an increased microbial activity seen under elevated CO₂ concentrations (Selsted et al. 2012) and by an isotope study done by Andresen et al. (2011) that showed traceable labeled C and N in the detritivores within a year from labeling in CO₂ plots. The large allocation of carbon to belowground processes may reflect nutrient limitation of the plants and may explain why I did not observe any responses in biomass to the CO₂ fumigation (paper I). If the increased microbial activity released nutrients, I would expect a stimulation of plant growth with time, but immobilization of nutrients by microbes may prevent such a release at least in the short-term (Michelsen et al. 1999).




Figure 6 Number of *D. flexuosa* flowering culms (mean±SE) from 2006 to 2010 at the CLIMAITE site. There was an overall CO₂ effect $p=0.036$; repeated measurement. Treatments were: A-ambient, D-summer drought, T-warming 1-2°C, CO₂-enriched CO₂ concentration of 510 ppm, and all treatment combinations.

Some nutrients used for flowering were lost to the litter pool after flowering (paper III). Since grass has a less complex structure and a faster decomposition rate compared to *C. vulgaris* (Aerts 1995), an increased production of flowering culms in *D. flexuosa*, and thus of litter, could increase the rate of nutrient turn-over even further. However, the C:N ratio of both leaf tissue and of roots was found to increase in response to CO₂ fumigation (Albert 2011a; Arndal (submitted)). This was also reflected in litter chemistry (Larsen et al. 2011), and with time, this may result in a decreased mineralization rate, increased nutrient immobilization by microbes and thus in a decreased amount of plant-available nutrients.

I would expect the increase in C:N ratio, as described by Albert et al. (2011a; 2011b) and Larsen (2011), to be caused by an increase in plant biomass in response to CO₂ and thus a dilution of N, nevertheless I did not see an increase in the aboveground biomass within this experiment. But Arndal (pers. comm.) found an increase in belowground biomass and it is therefore plausible that nitrogen was allocated from the above ground part to belowground. The results could perhaps also be explained by an increase in carbon-based secondary metabolites or low molecular mass sugar compounds that increased the biomass, but was undetectable with the pin-point method used for biomass estimates. Unfortunately, presently, we have no data to document whether or not this was indeed the case.

So, to answer the question whether the plant community will change over time: The results from the CO₂ manipulation suggested that the system was a lot more resilient towards changes than expected.

DROUGHT

Drought influenced the aboveground vegetation directly by causing a decrease in biomass during midsummer (paper I) (Fig 8) and by increasing the amount of flower abortions (paper III) (Fig 9). However, the drought effect on plant biomass only lasted for a short period of time; just two months after the drought period, we observed a full recovery of both grass and heather biomass. On the other hand, drought decreased the rate of most biological processes studied at the site, at least periodically. This included photosynthesis, litter decomposition and gross mineralization (Albert et al. 2011a; Albert et al. 2011c; Larsen et al. 2011). Also, the enchytraid biomass (Maraldo et al. 2010a) and the C:N ratio of plant tissue decreased (Albert et al. 2011b) in response to the drought treatment, and I therefore consider drought to be an important factor in changing the plant community in the future. Nevertheless, drought effects often last for a short period of time, and in many cases CO₂ counteracted the effects. For instance, the photosynthetic rate, enchytraeid biomass and the C:N ratio all responded oppositely, when drought was combined with elevated CO₂, indicating that short term drought periods may not be as important to these biological processes as previously




Figure 7 Weight fraction of *Calluna vulgaris* in year 2007-2009. Fractions were: wood (A), leaves (B) and flowers (C). Treatments were: A-ambient, D-summer drought, T-warming 1-2°C, CO₂-enriched CO₂ concentration of 510 ppm and all treatment combinations.

believed. However, results in this thesis indicate that the severity of the drought is very important. In order to avoid a major tissue dieback, the drought treatment was alleviated whenever the SWC fell below 5%, and we may therefore not see the full effect of the predicted future drought patterns. I expect the ecosystem would show long-lasting responses if future drought pushes heathland plants beyond their drought threshold and causes major tissue death. Further, as seen for both enchytraeid biomass and photosynthetic activity of *D. flexuosa*, the magnitude of the rewetting after drought also has a large impact on the organisms' opportunity to recover (Maraldo and Holmstrup 2009b; Albert et al. 2011c).

At Brandbjerg, where *C. vulgaris* was in the building/mature stage (Gimingham 1972), we observed an increase in *C. vulgaris* biomass over the study period (paper I) but no response to the drought treatment. At the Mols site, on the other hand, where *C. vulgaris* was regenerating after a disturbance, we observed a decrease in biomass of the pioneer stage, when subjected to drought treatment compared to warmed and control treatments (paper IV) (fig 10). This drought-induced decrease in *C. vulgaris* was associated with an increase in grass biomass. We consider this a key finding to answer some of the unanswered questions regarding the CLIMAITE site, Brandbjerg. With this finding from Mols in mind, I expect *C. vulgaris* to be much more susceptible to drought damage after an intense disturbance event such as heather beetle attacks, but potentially also frost damage and severe drought. This is important to keep in mind, since heather beetle attacks are expected to increase in the future due to higher N-deposition (Sheppard et al. 2008; Heil & Bobbink 1993). Also, frost damage events will properly be more frequent because of more frequent freeze-thaw cycles due to increased temperature and earlier snow melt (Walther et al. 2002). Furthermore, frost damage/winter desiccation will induce openings in the heather canopy and thus allow invasion of *D. flexuosa* (Bobbink et al. 1998).

TEMPERATURE

I did not observe any changes in aboveground vegetation in response to increased temperature. *Calluna vulgaris* has a natural distribution from the North of Norway down to Portugal, and *D. flexuosa* is common in most of Europe. Both species are therefore able to compete successfully within a large temperature amplitude which may explain the lack of response to warming. However, observations of an earlier leaf appearance (NDVI measurement) (fig 5) (paper I) and earlier appearance of inflorescences when subjected to increased temperatures (paper III) were reported in this study. Warming also resulted in higher photosynthetic rates at the site (Albert et al. 2011a), and together these changes are evidence of an earlier onset of the growing season in response to warming. Even so, these changes were neither transformed into increased biomass later during the season nor into more




Figure 8 A) Biomass of *D. flexuosa* (g m⁻², mean±SE), B) biomass of *C. vulgaris* (g m⁻², mean±SE). Ambient (White) and drought treatment (grey). Litter (hatched) and alive (unhatched) material. *=p<0.05.




Figure 9 Flowering in *Deschampsia flexuosa*. A) Flowers dead before flowering and fruit setting in 2008 and 2009. Treatments: A-ambient, D-summer drought for about 4-5 weeks, T-warming of 1-2°C, CO₂-enriched CO₂ concentration of 510 ppm and all combinations. Means±SE.

flowers. Thus, the changes should probably be seen only as a transient event, with limited ecological repercussions.

Warming was also hypothesized to accelerate most biological processes and thereby have an indirect effect on the plant community. An increase in the mineralization rate and N-turnover were reported by Andresen et al. (2010a) and Larsen et al. (2011), but within the time frame of this thesis I did not observe any secondary effects on the aboveground biomass. How, and whether, the plant community will change under a future warmer climate depends on the outcome of those interactions, but it seems fair to expect that changes in belowground processes will result in effects on the aboveground plant community – at least given a longer time scale than our five year study period.

INTERACTIONS

In general I did not see many statistically significant interactions between the climate parameters. This indicates that interactions did not influence the effects of single manipulations on the aboveground vegetation. However, the experiment has so far lasted only five years, and effects of interactions




Figure 10 Recruitment of new *Calluna vulgaris* plants after the heather beetle attack in 1999 registered as hits per pin. Treatments: Control, Summer drought for about 4-5 weeks and Warming of 1-2°C. Values and bars are means \pm SE.

between single factors may appear only after a longer period of climate manipulation. I had expected some of the single-factor effects to be either leveled out or enhanced by interactions between climatic factors similar to the way reported by Larsen et al. (2011). They reported most interactions to be antagonistic and therefore the system to be more robust when more than one factor was changed. For instance C/N ratio in mosses increased in response to drought and to CO₂, but not when these factors were combined.

SEASONAL VARIATIONS

We observed that several effects of the treatment were dependent on the seasonal variation in natural climate –mainly the amount and timing of precipitation within a year. Photosynthesis measured at the site was high in spring and autumn (Albert et al. 2011a), whereas a combination of tissue dieback in *D. flexuosa*, as shown in paper I and paper II, and a lower photosynthetic capacity resulted in lower carbon assimilation during midsummer (Albert et al 2011 submitted). However, this dieback and reduction in carbon assimilation did not reduce the overall biomass – quite the contrary: Albert et al. (2011b) showed a higher photosynthetic rate following the prolonged drought period, and I observed that grass biomass had already recovered one month after the severe drought event (Paper I). Seasonal variations in microbial biomass may be essential for plants, since a microbial dieback could release nutrients and thus temporarily increase the N-availability. The fast recovery could have been due to such a nutrient release during re-wetting following the drought period (van Meeteren et al. 2008).

There are many reports on nutrient immobilization by microbes (Jonasson et al. 1996; Schmidt et al. 1999). Andreasen et al. (2010a) concluded, based on a study from the same site, that microbes acquire glycine faster than plants, and therefore may be better competitors for nutrients. Immobilization of the nutrients and a nutrient limitation may therefore constitute at least some of the reason for the absent response of the plant biomass (paper I) to the climate change manipulations, despite the higher photosynthetic rate after re-wetting (Albert et al. 2011b). However, nutrients immobilized by microbes may periodically be released, and in general the microbial biomass is kept at the same level by the availability of carbon and by grazers. This was also seen at the site where enchytraeid biomass increased in response to elevated CO₂ depending on season, with the largest response during summer (Maraldo et al. 2010a). Further, Nielsen et al 2009 showed that there was a good correlation between nutrient availability and SWC at the site and this may reflect that microbial activity is controlled by the soil moisture (Nielsen et al. 2009). This indicates that the amount of nutrients that were available for the plants differed within the season, and that a potential microbe-plant interaction may explain the plant biomass to a higher degree than expected. This is of course important in order to understand the ecosystem, but it also

highlights the importance of incorporating such seasonal variations when planning studies and fieldwork. One sampling only provides a snapshot, and to understand the ecosystem processes, it is important not to make conclusion about ecosystem responses, based upon short time drought effects during e.g. midsummer.

INTER-ANNUAL VARIATIONS

Climate change is ongoing; in the period of 5 years where the study took place we observed large inter-annual variations in the climate. In addition to the climate manipulations within the two climate experiments, we can use this inter-annual variation in climate to describe and test the effects of climate change on the plant growth and phenology.

Especially the precipitation pattern showed inter-annual variations. There were, for instance, large differences between years in the amount of rain falling from September to November (Table 2), and since autumn SWC was positively correlated with the number of inflorescences, this could potentially explain the large inter-annual variation found in flowering (paper III).

The growing season in 2007 was very wet compared to the last 30 years average in Denmark. The following season in 2008 was, on the other hand, very dry compared to normal. This inter-annual variation in precipitation resulted in variations in the amount of precipitation that was excluded under the drought treatments (see table 2). Furthermore, we recorded months with precipitation far above average indicating heavy rain falls (July 2007 and June 2009).

In all years from 2006 to 2009 the temperature in the warmest month was 4-6 °C above the 30 year average for that month, and the coldest months were also mostly warmer than average, but with a larger variation, with some years close to average and others 5°C above average. By comparison, the warming curtains increased the temperature with 1°C in the soil surface all year round, and thus the increase caused by the temperature treatment was less than the natural variation between years. All this caused inter-annual variations in many biological processes that out-scaled those seen in response to the treatments. The large climatic inter-annual variations also resulted in different responses to the manipulations from year to year. For instance, we saw an increase in the number of *D. flexuosa* inflorescences in response to enhanced CO₂, but only in 2 out of 5 years. Maraldo et al. (2010b) also reported that the observed drought effect on enchytraeid biomass was dependent on the amount of precipitation within that year, i.e. the drought treatment had a more pronounced impact on enchytraeids during the year with a dry summer and autumn (2006) compared to the wet year (2007). Further, leaching and thus loss of nitrogen from the system was shown to vary between years, with $0.13 \pm 0.06 \text{ g N m}^{-2} \text{ yr}^{-2}$ in 2006 and $0.56 \pm 0.20 \text{ g N m}^{-2} \text{ yr}^{-2}$ in 2007 (Larsen et al. 2011). These findings signify the importance of

long term studies, and emphasise that conclusions based on only one or two seasons should be drawn with outmost caution.

Resilience to disturbances

Results from this thesis showed that the vulnerability of the heathland ecosystems depends on the phase of *C. vulgaris*. The ecosystem in the aggregating to mature phase was a lot more resilient towards changes in climate than expected. So to the question whether the plant community will change due to climate change, the answer may be 'yes, but not so much as we thought'. So, the next question followed: Why? In the following I will try to give some answers to this question.

1) This study showed that *D. flexuosa* has a high resilience to changes in climatic factors determined by its ability to let the tissue dieback and then quickly recover once conditions again become favourable, whereas *C. vulgaris* shows a strong resistance determined by its evergreen status, at least when it is established. Overall this led to a general lack of response for this heathland ecosystem to the climatic treatments. The plant biomass recovered rapidly after drought events, and was not affected by an earlier onset of the growing season (paper I). Thus the manipulation did not directly affect the plant biomass production. However, other studies at the site have shown that changes do occur; photosynthesis, mineralization, litter quality and flowering were all affected by the climate manipulations. Thus, it is possible that some of the observed changes in other ecosystem processes with time will affect the aboveground plant biomass, and this way the manipulations may still affect plant biomass, however in a more indirect way.

2) The manipulations caused smaller variations than the natural climate conditions. The plants are thus adapted to changes in climate exceeding the manipulations applied. Further, they are adapted to nutrient- and water limited conditions, and the capability of the plants to use both nutrients and water when available and then survive long periods without, may help to keep the ecosystem in balance.

3) In general, the drought treatment affected most biological processes, but it was also the climate driver for which effects were most dependent upon the inter-annual natural variation; i.e. the drought treatment was only statistically significant if supplementing a naturally occurring drought. In CLIMAITE it was decided not to push the plants beyond the drought threshold, in order to avoid a full dieback of any of the two species. This may explain the high resilience of the aboveground biomass. A stronger water limitation could possibly also have shed light on the differences between the two life form strategies in their responses to drought.

4) Results from this thesis indicate that the heather at Brandbjerg was in a state where it was very resistant towards changes. Although a number of es-

pecially soil parameters responded to the treatment, the plant community responded neither to direct nor to indirect effects. Together with results from the regeneration of *C. vulgaris* at the Mols site (paper IV), this leads us to conclude that both the state of the *C. vulgaris* population and the magnitude and frequency of disturbances determine the effects of future climate change on the plant community.

Carbon sequestration

Changes in climate can not only influence plant species but may also change the ecosystem structure and ecosystem functioning (Aerts et al. 2006; Beier et al. 2009). Terrestrial ecosystems play an important role in regulating the atmospheric carbon concentration. Based on this thesis, the plant composition at Brandbjerg did not change, meaning that aboveground vegetation did not contribute to changes in the carbon pools. However, we expect changes in the plant community on a longer time scale, either due to indirect effects slowly altering the relations between the two species or due to sudden major events negatively affecting the heather vegetation and providing the grass with an opportunity to increase in cover (Aerts et al. 1990; Heil & Diemont 1983; Heil & Bruggink 1987). If the vegetation changes from heather to grass, it would lead to a loss of organically bound carbon, simply because the standing biomass of a *C. vulgaris* vegetation is larger than that of a *D. flexuosa* vegetation (Kongstad et al. 2011). Since litter from *D. flexuosa* is more easily remineralized than *C. vulgaris* litter (Aerts 1995), a change towards grass-dominance, would probably also cause the system to release more CO₂, and thus increase the ‘source effect’ of the system compared to the ‘sink effect’. Whether coupled to changes in plant species composition or not, results from the site have indeed shown an increase in the release of CO₂ from the ecosystem due to increased microbial activity under full treatment combination (T×D×CO₂) compared to ambient plots (Selsted et al. 2012). If carbon sequestration of the ecosystem is to remain unchanged in the future, this increased loss would have to be mirrored by an equal increase in photosynthetic assimilation.

The future carbon sequestration will however depend on the level of future management. Since heathlands are semi natural ecosystems lack of management will lead to invasions by bushes and trees and thus increase the vegetation carbon pool and change the source- and sink processes.

HEATHLANDS IN A FUTURE CLIMATE

Several conditions have changed and will change heathland ecosystems in the future. In the following section I describe some of the expected interactions between changes in climate conditions and other threats affecting heathland ecosystems, including land-use, nitrogen deposition and management. It is an attempt to see the results from this thesis in a larger context and answer the question whether plant communities will change in the future. Since most of the threats were not studied within this thesis, this part relies mostly on the literature.

Land-use and fragmentation

Lately, heathland research has focused on climate changes in order to understand changes in functions and services of the ecosystem and to prepare for incorporation into management plans. But to venture into the future, we must first go back in time. Most heathlands are a result of human activity and land-use. Nutrients were removed from the heathlands by domesticated grassing animals or by harvesting of the heather for later use as winter fodder. The organic layer was used as fuel and the heather was burned to keep some areas in early succession phase and thereby ensure a high nutrient quality of the fodder. All together the ecosystem was kept in a nutrient poor state, where only species adapted to this condition would grow.

Heathlands no longer provide services for human existence. Many heathlands were taken into formal agricultural use in the middle of the last century, or were disturbed by infrastructural and urban expansion – a process that is still going on today.

This results in a decrease in the total heathland area but also in a high fragmentation of heathlands. Fragmentation decrease genetic variation of both heathland plants and animals within an area, due to a prevented or delayed spreading and pollination from other populations (Assmann and Janssen 1999; Vergeer et al. 2003). This could, with time, make the vegetation less resistant to changes, both due to less genetic variation and due to a decrease in biodiversity - both of which may decrease the stability of ecosystems. Climate change will alter growth conditions further, and may result in suppression of some species and the introduction of others. Alien species are often characterised by high spreading ability and new growth conditions may lead them to germinate in sensitive habitats such as heathlands (Penuelas et al. 2010). Biodiversity is also threatened within heathland ecosystems. Heathlands are characterized by a low plant biodiversity, dominated by dwarf shrubs co-existing with a few graminoids and herbs. Disturbance of growth conditions by climate change or land-use may initially increase the plant biodiversity due to an introduction of new species (Bobbink et al.

1998). However, with time heathland species will disappear and plant biodiversity will decrease, and land-use changes is predicted to be the most severe driver of changes in biodiversity in grassland ecosystems (Sala et al. 2000).

Distribution regimes

Plant distribution is among other things determined by species-specific physiological thresholds of temperature and water requirements (Woodward and Williams 1987). *Calluna vulgaris* has a wide distribution in Europe. It is a drought tolerant species with xeromorphic leaves (Aerts 1995) and low, but constant, photosynthesis even under dry conditions (Albert et al. 2011c). Higher temperature and changes in precipitation may alter the plant distribution and may accelerate the transformation of heathland/grassland into temperate forest.

Crawford (2008) predicted the biomes of *C. vulgaris* to move either northward if summer temperature increases more than winter temperature, or eastward if winter temperature increases more than summer temperature. Fosaa et al. (2004) found *C. vulgaris* to be positively correlated to increased temperature whereas *D. flexuosa* was found to be negatively correlated. Further, reports from Spain have shown that a replacement of *C. vulgaris* by oak (*Quercus* spp.) has already occurred due to a warmer climate and land use changes (Penuelas and Boada 2003).

These findings indicate that increased temperature may change the plant community at our site with time. *Calluna vulgaris* seedlings are, however, susceptible to drought and require maintenance of high soil moisture levels and high air humidity (Miller and Cummins 1987) and the species is therefore often found near the coastline because of the higher amount of precipitation here compared to continental climate. Results from paper IV indicate that a decrease in summer precipitation will have consequences for the spreading of *C. vulgaris* due to water limitation in the pioneer phase, and future changes in precipitation may therefore be more important for the future distribution regime of this species than increased temperature.

Besides changes in temperature and precipitation, competition with other present and new-coming species also has to be taken into account. Further, the genetic variation in the *C. vulgaris* population from this site may be very different from that of the *C. vulgaris* population in the boreal and subarctic region, where growing seasons are shorter and the climate is cooler, or from the Southern European populations where drought tolerance is a key element to survive. Changes due to climate change may therefore be different from site to site.

Nitrogen deposition

Changing climatic conditions may interact with effects of the present high nitrogen deposition. Some changes will enhance the effects of the increased nitrogen load whereas others will counteract.

Brandbjerg heathland was, as mentioned before, very resilient towards changes in climate and also towards nitrogen and phosphorus addition (paper II). There was no response in biomass when adding fertilizer, and already one year after the fertilizer addition, leaf nutrient concentrations were back to previous levels. This could be an effect of nutrient poor adapted plants being less efficient in translating an enhanced nutrient supply into a biomass increase – a strategy that could be important if the source is only temporarily available. In that case it would be advantageous to save the excess nutrients in storage organs or invest it in seed production (Falk et al. 2010). Further, other studies from the CLIMAITE experiment have indicated that the ecosystem seasonally was limited by water (Albert et al. 2012 in press; paper II) and since nutrient availability are correlated with SWC (Nielsen et al. 2009), the lack of response to nutrient addition may be caused by water limitation. However, the bryophytes responded negatively to nutrient addition by a decrease in biomass which may result in a decrease in biodiversity since 10 species were mosses compared to 17 vascular plants and 9 lichens (the latter were not included in this study).

Higher nitrogen availability has also been shown to increase *C. vulgaris* sensitivity towards drought (Gordon et al. 1999; Sheppard et al. 2008), and increased nitrogen deposition may therefore decrease the drought threshold and enhance the damage induced by future severe drought events. An increase in nitrogen availability either due to increased mineralization or increased nitrogen deposition may also accelerate the life cycle of *C. vulgaris* (Gimingham 1972). Since grasses are competitively superior in the early phase of heathland development, especially after exposure to drought as shown in paper IV, shortening of the *C. vulgaris* life cycle or more frequent heather beetle attacks or severe drought events may result in further dominance of grasses. Heather beetle attacks are expected to be more frequent in a milder climate, where both larval and adult survival may be higher (Power et al. 1998) and where warming may increase nutrient levels in the leaves (Penuelas et al. 2004). All this indicate a shift from heather to grass in the future. On the other hand, Albert et al. (2011a) and Larsen et al. (2011) reported an increased C:N ratio in leaf tissue in response to an enhanced CO₂ concentration and thereby a decrease in the nutrient quality of the leaves. That is opposed to the general effect of increased nitrogen deposition in heathlands which results in lower tissue C:N ratio and thus a better food quality for herbivores (Berdowski 1994). The increased C:N ratio in litter in response to increased CO₂ consequently counterbalance the N induced in-

crease in mineralization rate, and Larsen et al. (2011) concluded that the outcome would be a decrease in nitrogen cycling. Warming is also expected to increase the mineralisation rate and thereby enhance the nitrogen effect making more nutrients available for plants. However, the third environmental factor, precipitation, also plays a role in shaping the N-cycle: As seen in Larsen et al. (2011), leaching of nitrogen was dependent on the precipitation pattern – a higher frequency of heavy rain falls is expected to increase nitrogen leaching and thereby the nitrogen output from the site.

Finally, increased N deposition can have an impact on the carbon sequestration within the ecosystem (Wamelink et al. 2009). If plant growth is limited by nitrogen, increased nitrogen availability will increase growth and thereby the amount of carbon incorporated into plant biomass. However, De graff et al. (2006) concluded that increased C sequestration will only occur if nutrients are supplied. We assume the yearly nitrogen supply at Brandbjerg will not increase further, since it is used for military activities, but the current level may already be too high to maintain the current vegetation composition. Time will show whether increased nitrogen availability will change vegetation composition from dwarf shrub dominance with a high, all year around, standing carbon pool to grass dominance with higher productivity but lower standing biomass, and thus less carbon storage in aboveground vegetation.

CONCLUSION

In conclusion, and to answer the question: **Will changes in climate alter plant communities over time?**

This heathland ecosystem in the aggregating/mature phase was a lot more resilient towards changes in climate than expected. None of the applied manipulations resulted in long lasting changes in the aboveground biomass. The resilience was probably due to:

- *Calluna vulgaris* had a strong resistance to changes in climate determined by its evergreen status, maintaining the growth constantly over the season.
- The lack of response for *D. flexuosa* to the climatic treatments came from the plants ability to let the tissue dieback and then quickly recover once conditions again became favourable. It therefore showed a high resilience to changes in climate factors.

I conclude that this plant community was resilient to changes in climate – at least on a short time scale of a few years. On the other hand, the Mols study, where *C. vulgaris* was regenerating after a disturbance, showed a decrease in biomass of the pioneer stage, when subjected to drought treatment compared to warmed and control treatments. I therefore conclude that the vulnerability of the heathland ecosystems may depend on the growth phase of *C. vulgaris*, indicating that, in the long run, we may see effects at Brandbjerg as well.

The increase in flowering culms in response to elevated CO₂, the higher fraction of flower abortions in response to drought and the decrease in moss biomass in response to nitrogen addition indicate that changes do occur. This, together with findings from other studies at the Brandbjerg site, indicated that the biomass production and plant composition may not be the first parameter to mirror effects of climate changes on the ecosystem. Thus, it is possible that some of the observed changes in other ecosystem processes with time will also affect the aboveground plant biomass, and this way the manipulations may still affect plant biomass but in a more indirect way. This confirms that studies on climate change and ecosystems should cover the entire ecosystem. Further, it signifies the importance of longer term studies, and emphasises that conclusions based on only one or two seasons should be drawn with outmost caution.

In general I did not see many statistically significant interactions between the climate parameters. This indicates that interactions did not influence the effects of single manipulations on the aboveground vegetation. However, we do have to keep in mind that the experiment has so far only lasted five years, and that effects of interactions between single factors may appear only after a longer period of climate manipulation. Further, we deliberately did not push the system above a certain drought threshold in order to avoid a full tissue dieback, and we therefore may not see the full effects of the future drought patterns. I expect the ecosystem would show long-lasting responses if the drought period exceeds such a threshold.

Together with results from the regeneration of *C. vulgaris* at the Mols site, this leads us to conclude that both the state of the *C. vulgaris* population and the magnitude and frequency of disturbances determine the effects of future climate change on the plant community.

FURTHER RESEARCH

This thesis deals with the effects of climate change on the aboveground vegetation in a Danish heathland. However, the studies presented only lasted for 1- 5 years, which is a very short time in a climate change perspective. All results in this thesis are therefore short term responses and preliminary conclusions. Longer term studies are needed to overcome some of the differences between seasons and to survey a possible tipping point. Further, the

age of the vegetation was shown to be important and more time is needed to study responses and recovery from a disturbance in all stages of the *C. vulgaris* life cycle.

Changes in biodiversity, both new species and species that disappear, also need a longer time scale than 5 years before they become apparent. Furthermore, potentially invasive species are not dealt with in this thesis, but could, on a longer time scale, be extremely relevant to both the species composition and ecosystem functioning.

Studies on flowering pattern, seed production and seed quality for both species all needs to be studied closer. For instance, a germination experiment within the climate manipulation should be implemented. Also methods (for instance NDVI) for observing phenological patterns has to be developed. Further, this thesis does not cover any genetic variation, acclimation or adaptation.

Finally, possible interactions between climate change and other parameters, including nitrogen deposition and management, are presently only assumptions, and since this type of ecosystem is only maintained by management actions, studies with a management perspective are needed.

REFERENCES

Aerts R (1993) Competition between dominant plant species in heathlands. In *Heathlands: Patterns and Processes in a Changing Environment*. Springer.

Aerts R (1995) The Advantages of Being Evergreen. *Trends in Ecology & Evolution* 10:402-407

Aerts R, Berendse F, Decaluwe H, Schmitz M (1990) Competition in Heathland Along An Experimental Gradient of Nutrient Availability. *Oikos* 57:310-318

Aerts R, Cornelissen JHC, Dorrepaal E (2006) Plant performance in a warmer world: General responses of plants from cold, northern biomes and the importance of winter and spring events. *Plant Ecology* 182:65-77

Ainsworth EA, Long SP (2005) What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist* 165:351-371

Albert K, Ro-Poulsen H, Mikkelsen T, Michelsen A, van der Linden L, Beier C (2011a) Effects of elevated CO₂, warming and drought episodes on plant carbon uptake in a temperate heath ecosystem are controlled by soil water status. *Plant Cell and Environment* 34:1207-1222

Albert KR, Ro-Poulsen H, Mikkelsen TN, Michelsen A, van der Linden L, Beier C (2011b) Interactive effects of elevated CO₂, warming, and drought on photosynthesis of *Deschampsia flexuosa* in a temperate heath ecosystem. *Journal of Experimental Botany* 62:4253-4266

Albert KR, Mikkelsen TN, Michelsen A, Ro-Poulsen H, van der Linden L (2011c) Interactive effects of drought, elevated CO₂ and warming on photosynthetic capacity and photosystem performance in temperate heath plants. *Journal of plant physiology* 168:1550-1561

Albert KR, Kongstad J, Schmidt IK, Ro-Poulsen H, Mikkelsen TN, Michealsen A, Beier C (2012) Temperate heath plant response to dry conditions depends on growth strategy and less on physiology. *Acta Oecologica* (submitted).

Andalo C, Godelle B, Lefranc M, Mousseau M, TillBottraud I (1996) Elevated CO₂ decreases seed germination in *Arabidopsis thaliana*. *Glob Change Biol* 2:129-135

Andersen JM, Boutrup S, Bili L, van der Svendsen LM, Bøgestrand J, Grant R, Lauridsen TL, Ellermann T, Ærtebjerg G, Nielsen KE, Søgaard B, Jørgensen LF, Dalgren K (2005) Vandmiljø og Natur 2004. Tilstand og udvikling - faglig sammenfatning. pp 1-138. Danmarks Miljøundersøgelser, Miljøministeriet, Denmark.

Andresen LC, Michelsen A, Jonasson S, Schmidt IK, Mikkelsen TN, Ambus P, Beier C (2010a) Plant nutrient mobilization in temperate heathland responds to elevated CO₂, temperature and drought. *Plant and Soil* 328:381-396

Andresen LC, Konestabo HS, Maraldo K, Holmstrup M, Ambus P, Beier C, Michelsen A (2011) Organic matter flow in the food web at a temperate heath under multifactorial climate change. *Rapid Communications in Mass Spectrometry* 25:1485-1496

Andresen LC, Michelsen A, Ambus P, Beier C (2010b) Belowground heathland responses after 2 years of combined warming, elevated CO₂ and summer drought. *Biogeochemistry* 101:27-42

Arft AM, Walker MD, Gurevitch J, Alatalo JM, Bret-Harte MS, Dale M, Diemer M, Gugerli F, Henry GHR, Jones MH, Hollister RD, Jonsdottir IS, Laine K, Levesque E, Marion GM, Molau U, Molgaard P, Nordenhall U, Raszhivin V, Robinson CH, Starr G, Stenstrom A, Stenstrom M, Totland O, Turner PL, Walker LJ, Webber PJ, Welker JM, Wookey PA (1999) Responses of tundra plants to experimental warming: Meta-analysis of the international tundra experiment. *Ecological Monographs* 69:491-511

Assmann T, Janssen J (1999) The effects of habitat changes on the endangered ground beetle *Carabus nitens* (Coleoptera: Carabidae). *Journal of Insect Conservation* 3:107-116

Barker CG, Power SA, Bell JNB, Orme CDL (2004) Effects of habitat management on heathland response to atmospheric nitrogen deposition. *Biological Conservation* 120:41-52

Beier C (2004) Climate change and ecosystem function - full-scale manipulations of CO₂ and temperature. *New Phytologist* 162:243-245

Beier C, Emmett B, Gundersen P, Tietema A, Penuelas J, Estiarte M, Gordon C, Gorissen A, Llorens L, Roda F, Williams D (2004) Novel approaches to study climate change effects on terrestrial ecosystems in the field: Drought and passive nighttime warming. *Ecosystems* 7:583-597

Beier C, Emmett BA, Tietema A, Schmidt IK, Penuelas J, Lang EK, Duce P, de Angelis P, Gorissen A, Estiarte M, de Dato GD, Sowerby A, Kroel-Dulay G, Lellei-Kovacs E, Kull O, Mand P, Petersen H, Gjelstrup P, Spano D (2009) Carbon and nitrogen balances for six shrublands across Europe. *Global Biogeochemical Cycles* 23: Article Number: GB4008 DOI: 10.1029/2008GB003381

Beierkuhnlein C, Thiel D, Jentsch A, Willner E, Kreyling J (2011) Ecotypes of European grass species respond differently to warming and extreme drought. *Journal of Ecology* 99:703-713

Berdowski J J M (1994) The effect of external stress and disturbance factors on *Calluna*-dominated heathland vegetation. In *Heathlands; Patterns and Processes in a Changing Environment*. Eds. R Aerts and Heil G.W. pp. 85-124. Kluwer Academic Publisher.

Bobbink R, Hornung M, Roelofs JGM (1998) The effects of air-borne nitrogen pollutants on species diversity in natural and semi-natural European vegetation. *Journal of Ecology* 86:717-738

Chapin FS, Shaver GR (1996) Physiological and growth responses of arctic plants to a field experiment simulating climatic change. *Ecology* 77:822-840

Christensen JH, Hewitson B, Busuioc A, Chen A, Gao X, Held I, Jones R, Kolli RK, Kwon W-T, Laprise I, Magaña Rueda V, Mearns L, Menéndez CG, Räisänen J, Rinke A, Sarr A, Whetton P (2007) Regional Climate Projections. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (ed. S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller). Cambridge, United Kingdom and New York, NY, USA.

Cleland EE, Chiariello NR, Loarie SR, Mooney HA, Field CB (2006) Diverse responses of phenology to global changes in a grassland ecosystem. *Proceedings of the National Academy of Sciences of the United States of America* 103:13740-13744

Cornelissen JHC, Aerts R, Cerabolini B, Werger MJA, van der Heijden MGA (2001) Carbon cycling traits of plant species are linked with mycorrhizal strategy. *Oecologia* 129:611-619

Crawford R (2008) Woody plants at the margin. In *Plants at the Margin: Ecological Limits and Climate Change*. pp 309-337. Cambridge Univ Press, 32 Avenue of the Americas, New York, NY 10013-2473 USA.

Damgaard C (1999) A test of asymmetric competition in plant monocultures using the maximum likelihood function of a simple growth model. *Ecological Modelling* 116:285-292

de Graaff MA, van Groenigen KJ, Six J, Hungate B, van Kessel C (2006) Interactions between plant growth and soil nutrient cycling under elevated CO₂: a meta-analysis. *Glob Change Biol* 12:2077-2091

Ellenberg H (1991) *Indicator Values of Plants in Central Europe 1. Indicator Values of Vascular Plants Not Including Rubus*. *Scripta Geobotanica* 18:9-166

Emmett BA, Beier C, Estiarte M, Tietema A, Kristensen HL, Williams D, Penuelas J, Schmidt I, Sowerby A (2004) The response of soil processes to climate change: Results from manipulation studies of shrublands across an environmental gradient. *Ecosystems* 7:625-637

Falk K, Friedrich U, von Oheimb G, Mischke K, Merkle K, Meyer H, Hardtle W (2010) *Molinia caerulea* responses to N and P fertilisation in a dry heathland ecosystem (NW-Germany). *Plant Ecology* 209:47-56

Fosaa AM, Sykes MT, Lawesson JE, Gaard M (2004) Potential effects of climate change on plant species in the Faroe Islands. *Global Ecology and Biogeography* 13:427-437

Friedrich U, von Oheimb G, Dzedek C, Kriebitzsch WU, Selbmann K, Haerdtle W (2011) Mechanisms of purple moor-grass (*Molinia caerulea*) encroachment in dry heathland ecosystems with chronic nitrogen inputs. *Environmental Pollution* 159:3553-3559

Gimingham C H (1972) *Ecology of Heathlands*. Chapman and Hall, London.

Gordon C, Woodin SJ, Alexander IJ, Mullins CE (1999) Effects of increased temperature, drought and nitrogen supply on two upland perennials of

contrasting functional type: *Calluna vulgaris* and *Pteridium aquilinum*. *New Phytologist* 142:243-258

Graglia E, Jonasson S, Michelsen A, Schmidt IK, Havstrom M, Gustavsson L (2001) Effects of environmental perturbations on abundance of subarctic plants after three, seven and ten years of treatments. *Ecography* 24:5-12

Grime J (2002) *Plant Strategies, Vegetation Processes, and Ecosystem Properties*. John Wiley & Sons, pp. 456.

Grime J, Fridley JD, Askew AP, Thompson K, Hodgson JG, Bennett CR (2008) Long-term resistance to simulated climate change in an infertile grassland. *Proceedings of the National Academy of Sciences of the United States of America* 105:10028-10032

Hara Y, Toriyama K (1998) Seed nitrogen accelerates the rates of germination, emergence, and establishment of rice plants. *Soil Science and Plant Nutrition* 44:359-366

Heil GW, Bruggink M (1987) Competition for Nutrients Between *Calluna-Vulgaris* (L) Hull and *Molinia-Caerulea* (L) Moench. *Oecologia* 73:105-108

Heil GW, Diemont WH (1983) Raised Nutrient Levels Change Heathland Into Grassland. *Vegetatio* 53:113-120

Hikosaka K, Kinugasa T, Oikawa S, Onoda Y, Hirose T (2011) Effects of elevated CO₂ concentration on seed production in C₃ annual plants. *Journal of Experimental Botany* 62:1523-1530

Hovenden MJ, Newton P, Carran R, Theobald P, Wills K, Schoor J, Williams A, Osanai Y (2008) Warming prevents the elevated CO₂-induced reduction in available soil nitrogen in a temperate, perennial grassland. *Glob Change Biol* 14:1018-1024

Hudson JM, Henry GH (2010) High Arctic plant community resists 15 years of experimental warming. *Journal of Ecology* 98:1035-1041

Jablonski LM, Wang XZ, Curtis PS (2002) Plant reproduction under elevated CO₂ conditions: a meta-analysis of reports on 79 crop and wild species. *New Phytologist* 156:9-26

Jonasson S (1988) Evaluation of the Point Intercept Method for the Estimation of Plant Biomass. *Oikos* 52:101-106

Jonasson S, Michelsen A, Schmidt IK, Nielsen EV, Callaghan TV (1996) Microbial biomass C, N and P in two arctic soils and responses to addition of NPK fertilizer and sugar: Implications for plant nutrient uptake. *Oecologia* 106:507-515

Kongstad J, Schmidt I, Riis-Nielsen T, Arndal M, Mikkelsen T, Beier C (2011) High Resilience in Heathland Plants to Changes in Temperature, Drought, and CO₂ in Combination: Results from the CLIMAITE Experiment. *Ecosystems* 1-15

Larsen KS, Andresen LC, Beier C, Jonasson S, Albert KR, Ambus P, Arndal MF, Carter MS, Christensen S, Holmstrup M, Ibrom A, Kongstad J, van der Linden L, Maraldo K, Michelsen A, Mikkelsen TN, Pilegaard K, Prieme A, Ro-Poulsen H, Schmidt IK, Selsted MB, Stevnbak K (2011) Reduced N cycling in response to elevated CO₂, warming, and drought in a Danish heathland: Synthesizing results of the CLIMAITE project after two years of treatments. *Glob Change Biol* 17:1884-1899

Leakey ADB, Ainsworth EA, Bernacchi CJ, Rogers A, Long SP, Ort DR (2009) Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *Journal of Experimental Botany* 60:2859-2876

Luo YQ, Gerten D, Le Maire G, Parton WJ, Weng ES, Zhou XH, Keough C, Beier C, Ciais P, Cramer W, Dukes JS, Emmett B, Hanson PJ, Knapp A, Linder S, Nepstad D, Rustad L (2008) Modeled interactive effects of precipitation, temperature, and [CO₂] on ecosystem carbon and water dynamics in different climatic zones. *Glob Change Biol* 14:1986-1999

Maraldo K, Holmstrup M (2009a) Recovery of enchytraeid populations after severe drought events. *Applied Soil Ecology* 42:227-235

Maraldo K, Krogh PH, van der Linden L, Christensen B, Mikkelsen TN, Beier C, Holmstrup M (2010a) The counteracting effects of elevated atmospheric CO₂ concentrations and drought episodes: Studies of enchytraeid communities in a dry heathland. *Soil Biology & Biochemistry* 42:1958-1966

Maraldo K, Krogh PH, van der Linden L, Christensen B, Mikkelsen TN, Beier C, Holmstrup M (2010b) The counteracting effects of elevated atmospheric CO₂ concentrations and drought episodes: Studies of enchytraeid communities in a dry heathland. *Soil Biology & Biochemistry* 42:1958-1966

Maraldo K, Holmstrup M (2009b) Recovery of enchytraeid populations after severe drought events. *Applied Soil Ecology* 42:227-235

Menzel A, Sparks TH, Estrella N, Koch E, Aasa A, Ahas R, Alm-Kubler K, Bissolli P, Braslavska O, Briede A, Chmielewski FM, Crepinsek Z, Curnel Y, Dahl A, Defila C, Donnelly A, Filella Y, Jatcza K, Mage F, Mestre A, Nordli O, Penuelas J, Pirinen P, Remisova V, Scheifinger H, Striz M, Susnik A, Van Vliet AJH, Wielgolaski FE, Zach S, Zust A (2006) European phenological response to climate change matches the warming pattern. *Glob Change Biol* 12:1969-1976

Michelsen A, Graglia E, Schmidt IK, Jonasson S, Sleep D, Quarmby C (1999) Differential responses of grass and a dwarf shrub to long-term changes in soil microbial biomass C, N and P following factorial addition of NPK fertilizer, fungicide and labile carbon to a heath. *New Phytologist* 143:523-538

Miglietta F, Hoosbeek MR, Foot J, Gigon F, Hassinen A, Heijmans M, Peressotti A, Saarinen T, van Breemen N, Wallen B (2001) Spatial and temporal performance of the MiniFACE (Free Air CO₂ Enrichment) system on bog ecosystems in northern and central Europe. *Environmental Monitoring and Assessment* 66:107-127

Mikkelsen TN, Beier C, Jonasson S, Holmstrup M, Schmidt IK, Ambus P, Pilegaard K, Michelsen A, Albert K, Andresen LC, Arndal MF, Bruun N, Christensen S, Danbaek S, Gundersen P, Jorgensen P, Linden LG, Kongstad J, Maraldo K, Prieme A, Riis-Nielsen T, Ro-Poulsen H, Stevnbak K, Selsted MB, Sorensen P, Larsen KS, Carter MS, Ibrom A, Martinussen T, Miglietta F, Sverdrup H (2008) Experimental design of multifactor climate change experiments with elevated CO₂, warming and drought: the CLIMAITE project. *Functional Ecology* 22:185-195

Miller GR, Cummins RP (1987) Role of Buried Viable Seeds in the Recolonization of Disturbed Ground by Heather (*Calluna-Vulgaris* [L] Hull) in the Cairngorm Mountains, Scotland, Uk. *Arctic and Alpine Research* 19:396-401

Miyagi KM, Kinugasa T, Hikosaka K, Hirose T (2007) Elevated CO₂ concentration, nitrogen use, and seed production in annual plants. *Glob Change Biol* 13:2161-2170

Nielsen KE, Degn HJ, Damgaard C, Bruus M, Nygaard B (2011) A Native Species with Invasive Behaviour in Coastal Dunes: Evidence for Progressing Decay and Homogenization of Habitat Types. *Ambio* 40:819-823

Nielsen PL, Andresen LC, Michelsen A, Schmidt IK, Kongstad J (2009) Seasonal variations and effects of nutrient applications on N and P and microbial biomass under two temperate heathland plants. *Applied Soil Ecology* 42:279-287

Norby RJ, Luo YQ (2004) Evaluating ecosystem responses to rising atmospheric CO₂ and global warming in a multi-factor world. *New Phytologist* 162:281-293

Penuelas J, Boada M (2003) A global change-induced biome shift in the Montseny mountains (NE Spain). *Glob Change Biol* 9:131-140

Penuelas J, Gordon C, Llorens L, Nielsen T, Tietema A, Beier C, Bruna P, Emmett B, Estiarte M, Gorissen A (2004) Nonintrusive field experiments show different plant responses to warming and drought among sites, seasons, and species in a north-south European gradient. *Ecosystems* 7:598-612

Penuelas J, Prieto P, Beier C, Cesaraccio C, de Angelis P, de Dato G, Emmett BA, Estiarte M, Garadnai J, Gorissen A, Lang EK, Kroel-Dulay G, Llorens L, Pellizzaro G, Riis-Nielsen T, Schmidt IK, Sirca C, Sowerby A, Spano D, Tietema A (2007) Response of plant species richness and primary productivity in shrublands along a north-south gradient in Europe to seven years of experimental warming and drought: reductions in primary productivity in the heat and drought year of 2003. *Glob Change Biol* 13:2563-2581

Penuelas J, Sardans J, Llusia J, Owen SM, Carnicer J, Giambelluca TW, Rezende EL, Waite M, Niinemets U (2010) Faster returns on 'leaf economics' and different biogeochemical niche in invasive compared with native plant species. *Global Change Biology* 16:2171-2185

Power SA, Ashmore MR, Cousins DA, Sheppard LJ (1998) Effects of nitrogen addition on the stress sensitivity of *Calluna vulgaris*. *New Phytologist* 138:663-673

Rustad LE, Campbell JL, Marion GM, Norby RJ, Mitchell MJ, Hartley AE, Cornelissen JHC, Gurevitch J, Gcte NEWS (2001) A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming. *Oecologia* 126:543-562

Sala OE, Chapin FS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber-Sanwald E, Huenneke LF, Jackson RB, Kinzig A, Leemans R, Lodge DM, Mooney HA, Oesterheld M, Poff NL, Sykes MT, Walker BH, Walker M, Wall DH (2000) Biodiversity - Global biodiversity scenarios for the year 2100. *Science* 287:1770-1774

Schmidt IK, Jonasson S, Michelsen A (1999) Mineralization and microbial immobilization of N and P in arctic soils in relation to season, temperature and nutrient amendment. *Applied Soil Ecology* 11:147-160

Schmidt IK, Tietema A, Williams D, Gundersen P, Beier C, Emmett BA, Estiarte M (2004) Soil solution chemistry and element fluxes in three European heathlands and their responses to warming and drought. *Ecosystems* 7:638-649

Selsted MB, van der Linden L, Ibrom A, Michelsen A, Larsen KS, Pedersen JK, Mikkelsen TN, Pilegaard K, Beier C, Ambus P (2012) Soil respiration is stimulated by elevated CO₂ and reduced by summer drought: three years of measurements in a multifactor ecosystem manipulation experiment in a temperate heathland (CLIMAITE). *Glob Change Biol.* doi: 10.1111/j.1365-2486.2011.02634.x

Shaw MR, Zavaleta ES, Chiariello NR, Cleland EE, Mooney HA, Field CB (2002) Grassland responses to global environmental changes suppressed by elevated CO₂. *Science* 298:1987-1990

Sheppard LJ, Leith ID, Crossley A, Van Dijk N, Fowler D, Sutton M, Woods C (2008) Stress responses of *Calluna vulgaris* to reduced and oxidised N applied under 'real world conditions'. *Environmental Pollution* 154:404-413

Smith SE, Read DJ (2008) *Mycorrhizal Symbiosis*. Academic Press, London.

Sowerby A, Emmett BA, Tietema A, Beier C (2008) Contrasting effects of repeated summer drought on soil carbon efflux in hydric and mesic heathland soils. *Glob Change Biol* 14:2388-2404

Terry AC, Ashmore MR, Power SA, Allchin EA, Heil GW (2004) Modelling the impacts of atmospheric nitrogen deposition on *Calluna*-dominated ecosystems in the UK. *Journal of Applied Ecology* 41:897-909

Thurig B, Korner C, Stocklin J (2003) Seed production and seed quality in a calcareous grassland in elevated CO₂. *Glob Change Biol* 9:873-884

van Heerwaarden LM, Toet S, van Logtestijn RSP, Aerts R (2005) Internal nitrogen dynamics in the graminoid *Molinia caerulea* under higher N supply and elevated CO₂ concentrations. *Plant and Soil* 277:255-264

van Meeteren M, Tietema A, van Loon E, Verstraten J (2008) Microbial dynamics and litter decomposition under a changed climate in a Dutch heathland. *Applied Soil Ecology* 38:119-127

Vergeer P, Rengelink R, Ouborg NJ, Roelofs JGM (2003) Effects of population size and genetic variation on the response of *Succisa pratensis* to eutrophication and acidification. *Journal of Ecology* 91:600-609

von Oheimb G, Hardtle W, Falk K, Gerke AK, Meyer H, Drees C, Matern A (2009) Is *Calluna vulgaris* a suitable bio-monitor of management-mediated nutrient pools in heathland ecosystems? *Ecological Indicators* 9:1049-1055

Walther GR, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin JM, Hoegh-Guldberg O, Bairlein F (2002) Ecological responses to recent climate change. *Nature* 416:389-395

Wamelink G, van Dobben H, Mol-Dijkstra J, Schouwenberg E, Kros J, De Vries W, Berendse F (2009) Effect of nitrogen deposition reduction on biodiversity and carbon sequestration. *Forest Ecology and Management* 258:1774-1779

Woodward FI, Williams BG (1987) Climate and Plant-Distribution at Global and Local Scales. *Vegetatio* 69:189-197

Zak DR, Pregitzer KS, Curtis PS, Teeri JA, Fogel R, Randlett DL (1993) Elevated Atmospheric CO₂ and Feedback Between Carbon and Nitrogen Cycles. *Plant and Soil* 151:105-117

LIST OF PAPERS

- I. Jane Kongstad, Inger K. Schmidt, Torben Riis-Nielsen, Marie F. Arndal, Teis N. Mikkelsen and Claus Beier. **High Resilience in Heathland Plants to Changes in Temperature, Drought and CO₂ in Combination:** Results from the CLIMAITE Experiment. *Ecosystems* (2011) DOI: 10.1007/s10021-011-9508-9.
- II. Jane Kongstad, Sven Jonasson, Inger K. Schmidt, Pia L. Nielsen and Anders Michelsen. **Changes from heathland to grassland; a study on the impact of fertilizer on phenology, chemistry and growth of the heathland grass *Deschampsia flexuosa*.** Submitted to *Oecologia*, Jan 2012.
- III. Jane Kongstad, Inger K. Schmidt and Johannes Ransijn J. **Altered flower phenology and allocation of resources to reproductive structures in response to drought, warming and CO₂ enrichment.** Manuscript to be submitted to *Plant Ecology*
- IV. Inger K. Schmidt, Torben Riis-Nielsen T, Jane Kongstad, Johannes Ransijn, Christian Damgaard, Claus Beier. **Vulnerability of *Calluna vulgaris* in response to long-term climate manipulation and ecosystem disturbances.** Early draft of manuscript to be submitted to *Journal of Ecology*

The four papers are not included in this version of the thesis due to copyright.

Former issues of Forest & Landscape Research

(No. 1 - No. 26 issued in "The Research Series" (Forskningsserien))

- No. 1-1993 Stofkredsløb i sitkagran, rødgran og bøgebevoksninger i Danmark.
Lars Bo Pedersen
ISBN 87-89822-14-5. *Out of print*
- No. 2-1993 Provenienser af stilkeg (*Quercus robur* L.) og vintereg (*Quercus petraea* (Matthuschka) Liebl.) i Danmark.
Jan Svejgaard Jensen
ISBN 87-89822-16-1. DKK 150.00
- No. 3-1993 Growth and Yield Estimation from Successive Forest Inventories. Proceedings from the IUFRO Conference, held in Copenhagen, 14-17 June 1993.
J.K. Vanclay, J.P. Skovsgaard & G.Z. Gertner (ed.)
ISBN 87-89822-19-6. DKK 150.00
- No. 4-1993 Vanris på dansk stilkeg (*Quercus robur* L.).
Jan Svejgaard Jensen
ISBN 87-89822-22-6. DKK 100.00
- No. 5-1994 The Use of Sludge in Forestry and Agriculture. A Comparison of the Legislation in Different Countries.
Merete Morsing
ISBN 87-89822-24-2. DKK 100.00
- No. 6-1994 Marginaljorder og landskabet - marginaliseringsdebatten 10 år efter. Rapport fra et tværfagligt seminar afholdt af Dansk Landskabsøkologisk Forening i samarbejde med Institut for Økonomi, Skov og Landskab, KVL, Fredag den 25. september 1992.
Jesper Brandt & Jørgen Primdahl (ed.)
ISBN 87-89822-28-5. *Out of print*
- No. 7-1994 Landskabsøkologiske problemer i kystzonen. Rapport fra et tværfagligt seminar afholdt af Dansk Landskabsøkologisk Forening i samarbejde med Institut for Økonomi, Skov og Landskab, KVL, Fredag den 8. oktober 1993.
Erling Andersen (ed.)
ISBN 87-89822-29-3. DKK 75.00

- No. 8-1994 Throughfall and Canopy Interactions in Spruce Forest.
Karin Hansen
ISBN 87-89822-30-7. DKK 150.00
- No. 9-1994 Genetisk variation i sitkagran (*Picea sitchensis* (Bong.) Carr.) i højdevækst, stammeform og frosthærdighed - vurderet ud fra danske proveniens-, afkoms- og klonforsøg.
Ulrik Bräuner Nielsen
ISBN 87-89822-35-8. DKK 200.00
- No. 10-1994 Density Variations and Demarcation of the Juvenile Wood in Norway Spruce.
Frede Danborg
ISBN 87-89822-36-6. DKK 100.00
- No. 11-1995 Genetics and Silviculture of Beech. Proceedings from the 5th Beech Symposium of the IUFRO Project Group P1.10-00, 19-24 September 1994, Mogenstrup, Denmark.
Søren F. Madsen (ed.)
ISBN 87-89822-38-2. DKK 200.00
- No. 12-1995 Naturbeskyttelse og det almindelige agerland. Rapport fra det 4. landskabsøkologiske seminar afholdt af Dansk Landskabsøkologisk Forening i samarbejde med Institut for Økonomi, Skov og Landskab, KVL, 11. november 1994.
Jesper Brandt & Jørgen Primdahl (ed.)
ISBN 87-89822-43-9. *Out of print*
- No. 13-1996 Bøgeforyngelser i Østjylland.
J.P. Skovsgaard & M. Morsing (ed.)
ISBN 87-89822-45-5. DKK 250.00
- No. 14-1996 Bynære landbrugsområder i hovedstadsregionen 1994.
Susanne Ogstrup & Jørgen Primdahl
ISBN 87-89822-47-1. *Out of print*
- No. 15-1996 Plantevækst i forbindelse med byggeri. Planlægningens og projekterings indflydelse på vedplanters vækstvilkår i utilsigtet komprimerede jorder.
Thomas Barfoed Randrup
ISBN 87-89822-49-8. DKK 300.00

- No. 16-1996 Virkning af slamgødskning på det omgivende miljø og på biomassekvantitet og -kvalitet i energiskove af pil.
Keld Hauge Nielsen
ISBN 87-89822-58-7. DKK 150.00
- No. 17-1996 Træers forhold til salinitet. En behandling af træers reaktion på salt- og ionstress med vægt på arter fra den tempererede klimazone.
Jens Dragsted
ISBN 87-89822-64-1. DKK 300.00
- No. 18-1996 Juvenile Wood in Norway and Sitka Spruce. Anatomy, density, drying properties, visual grading and strength properties.
Frede Danborg
ISBN 87-89822-69-2. *Out of print*
- No. 19-1997 Tyndingsfri drift af sitkagran. En analyse af bevoksningsstruktur og vedmasseproduktion i utyndede bevoksninger af sitkagran (*Picea sitchensis* (Bong.) Carr.) i Danmark.
Jens Peter Skovsgaard
ISBN 87-89822-78-1. DKK 300.00
- No. 20-1997 Friluftsliv i skovene 1976/77 - 1993/94.
Frank Søndergaard Jensen & Niels Elers Koch
ISBN 87-89822-89-7. DKK 250.00
- No. 21-1997 Decline of mires in four Danish state forests during the 19th and 20th century.
Flemming Rune
ISBN 87-89822-94-3. DKK 100.00
- No. 22-1997 Fire artikler om nåletræer.
Ellen Juel Christensen (ed.)
ISBN 87-7903-003-3. DKK 125.00
- No. 23-1998 Vitalization of mature Norway spruce stands by fertilization and liming.
Morten Ingerslev
ISBN 87-7903-008-4. DKK 250.00
- No. 24-1998 Natural forest management among indigenous peoples in Latin America.
Søren Gram
ISBN 87-7903-020-3. DKK 125.00

- No. 25-1998 Friluftsliv i det åbne land 1994/95.
Frank Søndergaard Jensen
ISBN 87-7903-021-1. DKK 175.00
- No. 26-1999 Forest recreation in Denmark from the 1970s to the 1990s.
Frank Søndergaard Jensen
ISBN 87-7903-058-0. DKK 175.00
- No. 27-2000 Offentlige områdeudpegninger i jordbrugslandskabet.
Helle Tegner Anker, Ole Hjort Caspersen, Berit Hasler & Jørgen Primdahl
ISBN 87-7903-068-8. DKK 100.00
- No. 28-2000 Anvendelse og betydning af byens parker og grønne områder
(Use and importance of urban parks).
Stine Holm
ISBN 87-7903-075-0. DKK 300.00
- No. 29-2001 Recirkulering af organisk affald ved kompostering i lokale og større kredsløb.
Ulrik Reeh
ISBN 87-7903-115-3. DKK 150.00
- No. 30-2001 Vedvarende græsarealer – landbruget og reguleringen.
Erling Andersen
ISBN 87-7903-135-8. DKK 250.00
- No. 31-2002 Landskab, forandring og forvaltning - fem landskabsstudier fra Bjerringbro og Hvorslev
Ole H. Caspersen & Bo Fritzboeger (ed.)
ISBN 87-7903-142-0. DKK 250.00
- No. 32-2002 Implementation of Landscape Ecological Methods to GIS-based Procedures for Land Use Analysis and Decision Support in Rural Areas
Ole H. Caspersen
ISBN 87-7903-141-2. DKK 250.00
- No. 33-2003 Næringsstofkredsløb i skove - Ionbalanceprojektet
Karin Hansen (ed.)
ISBN 87-7903-156-0. DKK 300.00

- No. 34-2003 A study of factors affecting germination, establishment, and competition of the turfgrass species red fescue (*Festuca rubra* L. spp. *litoralis* Vasey), perennial ryegrass (*Lolium perenne* L.), and Kentucky bluegrass (*Poa pratensis* L.)
Søren Ugilt Larsen
ISBN 87-7903-202-8. DKK 250.00
- No. 35-2005 Nørholm Hede. En langtidsundersøgelse af hedens vegetationsudvikling og tilgroning
Torben Riis-Nielsen, Inger Kappel Schmidt, Bjarke Frandsen & Trine Binding
ISBN 87-7903-249-4. DKK 300.00
- No. 36-2006 Understanding and Communication forest stand structures. Lifting barriers for nature-based forest management
Anders Busse Nielsen
ISBN-10: 87-7903-285-0.
ISBN-13: 978-87-7903-285-0. DKK 250.00
- No. 37-2006 Bynære landbrugsområder I hovedstadsregionen 2004. Udvikling i landbrug, landskab og bebyggelse 1984-2004
Jørgen Primdahl, Anne Gravsholt Busck & Casper Lindemann
ISBN-10: 87-7903-299-0.
ISBN-13: 978-87-7903-299-6. DKK 200.00
- No. 38-2007 Skovopfattelse blandt danskere og i skovlovgivningen.
Liv Oustrup
ISBN 978-87-7903-308-5. DKK 300.00
- No. 39-2008 Transpiration and water budgets of European beech (*Fagus sylvatica* L.) dominated stands in relation to canopy structure
Lise Dalsgaard
ISBN 978-87-7903-350-4. DKK 300.00
- No. 40-2008 Partnerskaber i parkforvaltningen: Grøn drift på kontrakt
Andrej Christian Lindholst
ISBN 978-87-7903-378-8. DKK 250.00
- No. 41-2008 Status and prospects for urban green structure planning in China – Weihai city as a case study
Li Liu
ISBN 978-87-7903-380-1. DKK 300.00


- No. 42-2008 Enzymatic hydrolysis of lignocellulose. Substrate interactions and high solids loadings
Jan Bach Kristensen
ISBN 978-87-7903-393-1. DKK 200.00
- No. 43-2009 Marketizing municipal park management organisations in Denmark - A study of policymaking and organisational change in the period 1985-2005
Christine Nuppenau
ISBN 978-87-7903-395-5. DKK 250.00
- No. 44-2009 Unges friluftsliv i Danmark i det 21. århundrede – Unges friluftsliv som personligt, socialt og kulturelt identitetsprojekt på friluftslivets felt
Hans Jørgen Fisker
ISBN 978-87-7903-460-0. DKK 300.00
- No. 45-2010 Use of urban green space
Jasper Schipperijn
ISBN 978-87-7903-462-4. DKK 250.00
- No. 46-2011 Methods for physical characterisation of solid biofuels - a basis for standardisation
Peter Daugbjerg Jensen
ISBN 978-87-7903-518-8. DKK 200.00
- No. 47-2012 Root dynamics and below ground carbon input in a changing climate
Marie Frost Arndal
ISBN 978-87-7903-567-6
- No. 48-2012 Plant community responses to climate change
Jane Kongstad
ISBN 978-87-7903-575-1

Single issues are available from:

Forest & Landscape Denmark
University of Copenhagen
Rolighedsvej 23
DK-1958 Frederiksberg C
Tel. +45 3533 1500
SL-publikationer@life.ku.dk



Plant community responses to climate change



Forest & Landscape Denmark
University of Copenhagen
Rolighedsvej 23
DK-1958 Frederiksberg C
Tel. +45 3533 1500
sl@life.ku.dk
<http://sl.life.ku.dk/english>

National centre for
research, education and
advisory services within
the fields of forest and
forest products, landscape
architecture and landscape
management, urban planning
and urban design
