



Genome Sequence of the Acidophilic Bacterium *Acidocella* sp. Strain MX-AZ02

Servín-Garcidueñas, Luis E.; Garrett, Roger A.; Amils, Ricardo; Martínez-Romero, Esperanza

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.00041-12](https://doi.org/10.1128/genomeA.00041-12)

Publication date:
2013

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Servín-Garcidueñas, L. E., Garrett, R. A., Amils, R., & Martínez-Romero, E. (2013). Genome Sequence of the Acidophilic Bacterium *Acidocella* sp. Strain MX-AZ02. *Genome Announcements*, 1(1), 1-2.
<https://doi.org/10.1128/genomeA.00041-12>

Genome Sequence of the Acidophilic Bacterium *Acidocella* sp. Strain MX-AZ02

Luis E. Servín-Garcidueñas,^a Roger A. Garrett,^b Ricardo Amils,^c Esperanza Martínez-Romero^a

Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico^a; Department of Biology, Archaea Centre, University of Copenhagen, Copenhagen, Denmark^b; Centro de Biología Molecular Severo Ochoa, Universidad Autónoma de Madrid, Madrid, Spain^c

Here, we report the draft genome sequence of *Acidocella* sp. strain MX-AZ02, an acidophilic and heterotrophic alphaproteobacterium isolated from a geothermal lake in western Mexico.

Received 12 October 2012 Accepted 26 October 2012 Published 15 January 2013

Citation Servín-Garcidueñas LE, Garrett RA, Amils R, Martínez-Romero E. 2013. Genome sequence of the acidophilic bacterium *Acidocella* sp. strain MX-AZ02. *Genome Announc.* 1(1):e00041-12. doi:10.1128/genomeA.00041-12.

Copyright © 2013 Servín-Garcidueñas et al. This is an open-access article distributed under the terms of the [Attribution 3.0 Unported Creative Commons License](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Luis E. Servín-Garcidueñas, luis.e.servin@gmail.com.

Acidocella sp. strain MX-AZ02 was isolated from a naturally acidic (pH 2.3) and heavy metal-containing shallow lake in the Los Azufres National Park in western Mexico. The *Acidocella* genus comprises aerobic, acidophilic, Gram-negative bacteria belonging to the class *Alphaproteobacteria* (1). *Acidocella* relatives have been identified both in natural and acid mine drainage environments exhibiting high heavy-metal levels (2–6). *Acidocella* has also been detected among *Sphagnum* moss microbiota growing under varying acidic conditions (7, 8). Currently, the genus contains three reference strains isolated from acidic environments (9–11).

DNA was isolated from *Acidocella* sp. MX-AZ02, which yields smooth, round, and translucent colonies on DSMZ medium 35a. The organism is maintained at the Center of Genomic Sciences in the culture collection of the Ecological Genomics Department, National University of Mexico (UNAM). The sequencing was performed with the Roche 454 GS-FLX titanium technology generating 58.04 Mbp (~16-fold coverage) from a mate-paired library with 3-kb inserts. The reads were assembled *de novo* using Newbler assembler 2.3 (454 Life Sciences). The assembly produced 303 contigs of >500 bp each with an N₅₀ size of 22.12 kb. Nine scaffolds were generated containing 250 contigs. Genome annotation was performed using the NCBI Prokaryotic Genomes Automatic Annotation Pipeline (PGAAP) (<http://www.ncbi.nlm.nih.gov/genomes/static/Pipeline.html>).

The genome of *Acidocella* sp. MX-AZ02 was estimated to be 3.6 Mbp with a G+C content of 64.1% and it carried 3,553 open reading frames (ORFs). The 16S rRNA gene phylogeny indicated that the strain is closely related to type strains *Acidocella facilis* PW2, *Acidocella aluminiidurans* AL46, and *Acidocella aminolytica* 101, sharing 99.86%, 99.50%, and 97.93% sequence identities, respectively, over 1,407 bp.

Metal resistance determinants have been identified for *Acidocella* strains (12, 13, 14). The *Acidocella* sp. MX-AZ02 genome codes for arsenic, chromium, copper, and cobalt-zinc-cadmium transporters, as well as heavy-metal sensor signal transduction histidine kinases and chaperones. Carbonic anhydrases were also

encoded, which may provide a means to cope with the low CO₂ levels in acidic waters.

One *Acidocella* strain was shown to metabolize fructose from medium containing cell-free algal exudates, but it was unable to metabolize mannitol or glucose (15). *Acidocella* sp. MX-AZ02 may use glucose in the isolation medium as a carbon source. An acidophilic and abundant unicellular green alga was recently characterized from the same lake from which *Acidocella* sp. MX-AZ02 was isolated (16). Possibly, *Acidocella* sp. MX-AZ02 utilizes organic compounds from the alga, as was proposed previously for acidophilic microalgae and acidophilic heterotrophic bacteria (15).

The draft genome of *Acidocella* sp. MX-AZ02 will facilitate the identification of metal resistance determinants and may help us understand bacterial–algal interactions. This is the first isolated bacterial genome for an *Acidocella* strain and is the first sequenced bacterial genome from the Los Azufres National Park.

Nucleotide sequence accession number. The draft of the genome sequence is deposited at DDBJ/EMBL/GenBank under the accession no. [AMPS000000000](https://www.ncbi.nlm.nih.gov/nuclink/AMPS000000000).

ACKNOWLEDGMENTS

We thank the Next Generation Sequencing unit from Macrogen Inc. for pyrosequencing. The Comisión Federal de Electricidad (CFE) personnel provided a sampling permit.

This research was supported by PAPIIT IN205412 from DGAPA, UNAM. L.E.S.G. received a PhD scholarship from CONACYT (Mexico) under the Programa de Doctorado en Ciencias Biomédicas from UNAM. L.E.S.G. had a “Beca Mixta” from CONACYT and a fellowship from PAEP during a research internship at the Danish Archaea Centre.

REFERENCES

1. Kishimoto N, Kosako Y, Wakao N, Tano T, Hiraishi A. 1995. Transfer of *Acidiphilium facilis* and *Acidiphilium aminolytica* to the genus *Acidocella* gen. nov. and emendation of the genus *Acidiphilium*. *Syst. Appl. Microbiol.* 18:85–91.
2. Hallberg KB, Coupland K, Kimura S, Johnson DB. 2006. Macroscopic streamer growths in acidic, metal-rich mine waters in north Wales consist of novel and remarkably simple bacterial communities. *Appl. Environ. Microbiol.* 72:2022–2030.

3. Hamamura N, Olson SH, Ward DM, Inskeep WP. 2005. Diversity and functional analysis of bacterial communities associated with natural hydrocarbon seeps in acidic soils at rainbow springs, Yellowstone National Park. *Appl. Environ. Microbiol.* 71:5943–5950.
4. Johnson DB, Rolfe S, Hallberg KB, Iversen E. 2001. Isolation and phylogenetic characterization of acidophilic microorganisms indigenous to acidic drainage waters at an abandoned Norwegian copper mine. *Environ. Microbiol.* 3:630–637.
5. Lear G, Niyogi D, Harding J, Dong Y, Lewis G. 2009. Biofilm bacterial community structure in streams affected by acid mine drainage. *Appl. Environ. Microbiol.* 75:3455–3460.
6. Lu S, Gischkat S, Reiche M, Akob DM, Hallberg KB, Küsel K. 2010. Ecophysiology of Fe-cycling bacteria in acidic sediments. *Appl. Environ. Microbiol.* 76:8174–8183.
7. Bragina A, Maier S, Berg C, Müller H, Chobot V, Hadacek F, Berg G. 2011. Similar diversity of alphaproteobacteria and nitrogenase gene amplicons on two related *Sphagnum* mosses. *Front. Microbiol.* 2:275.
8. Opelt K, Berg G. 2004. Diversity and antagonistic potential of bacteria associated with bryophytes from nutrient-poor habitats of the Baltic sea coast. *Appl. Environ. Microbiol.* 70:6569–6579.
9. Kimoto K, Aizawa T, Urai M, Ve NB, Suzuki K, Nakajima M, Sunairi M. 2010. *Acidocella aluminiidurans* sp. nov., an aluminium-tolerant bacterium isolated from *Panicum repens* grown in a highly acidic swamp in actual acid sulfate soil area of Vietnam. *Int. J. Syst. Evol. Microbiol.* 60:764–768.
10. Kishimoto N, Kosako Y, Tano T. 1993. *Acidiphilium aminolytica* sp. nov.: an acidophilic chemoorganotrophic bacterium isolated from acidic mineral environment. *Curr. Microbiol.* 27:131–136.
11. Wichlacz PL, Unz RF, Langworthy TA. 1986. *Acidiphilium angustum* sp. nov., *Acidiphilium facilis* sp. nov., and *Acidiphilium rubrum* sp. nov.: acidophilic heterotrophic bacteria isolated from acidic coal mine drainage. *Int. J. Syst. Bacteriol.* 36:197–201.
12. Ghosh S, Mahapatra NR, Banerjee PC. 1997. Metal resistance in *Acidocella* strains and plasmid-mediated transfer of this characteristic to *Acidiphilium multivorum* and *Escherichia coli*. *Appl. Environ. Microbiol.* 63:4523–4527.
13. Ghosh S, Mahapatra NR, Ramamurthy T, Banerjee PC. 2000. Plasmid curing from an acidophilic bacterium of the genus *Acidocella*. *FEMS Microbiol. Lett.* 183:271–274.
14. Ghosh S, Mahapatra NR, Nandi S, Banerjee PC. 2005. Integration of metal-resistant determinants from the plasmid of an *Acidocella* strain into the chromosome of *Escherichia coli* DH5alpha. *Curr. Microbiol.* 50:28–32.
15. Nancucheo I, Barrie Johnson D. 2012. Acidophilic algae isolated from mine-impacted environments and their roles in sustaining heterotrophic acidophiles. *Front. Microbiol.* 3:325.
16. Servín-Garcidueñas LE, Martínez-Romero E. 2012. Complete mitochondrial and plastid genomes of the green microalga Trebouxiophyceae sp. strain MX-AZ01 isolated from a highly acidic geothermal lake. *Eukaryot. Cell* 11:1417–1418.