



Taenia solium cysticercosis - an emerging foodborne zoonosis in sub-Saharan Africa

Johansen, Maria Vang; Lekule , Faustin ; Pondja , Alberto ; Ngowi, Helena ; Assane , Yunus ; Kimbi, Eliakunda; Magnussen, Pascal; Mejer, Helena; Braae, Uffe Christian; Trevisan, Chiara; Saarnak, Christopher; Thamsborg, Stig Milan

Publication date:
2013

Document version
Peer reviewed version

Citation for published version (APA):
Johansen, M. V., Lekule , F., Pondja , A., Ngowi, H., Assane , Y., Kimbi, E., ... Thamsborg, S. M. (2013). *Taenia solium cysticercosis - an emerging foodborne zoonosis in sub-Saharan Africa*. Poster session presented at MED VET NET Association International Scientific Conference , Lyngby, Denmark.

Taenia solium cysticercosis

an emerging foodborne zoonosis in sub-Saharan Africa

Johansen, M.V., Lekule, F., Pondja, A., Ngowi, H., Assane, Y., Kimbi, E., Magnussen, P. Mejer, H., Braae, U. C., Trevisan, C., Saarnak, C. L., Thamsborg, S. M.

¹Section for Parasitology and Aquatic Diseases, Department of Veterinary Disease Biology, Faculty of Health and Medical Sciences, University of Copenhagen. ²Sokoine University of Agriculture, Faculty of Veterinary Medicine, Department of Veterinary Medicine and Public Health, Morogoro, Tanzania. ³Eduardo Mondlane University, Maputo, Mozambique


Taenia solium – a vicious tapeworm

Declared eradicable in 1993, among the Neglected Zoonotic Diseases, to be controlled by 2020 (WHO/FAO/OIE)

In theory: Eradicable

Existing control tools:

- Vaccine for porcine cysticercosis
- Drugs for porcine cysticercosis
- Drugs for human taeniosis



In reality: Emerging

- Number of pigs and people increase significantly in sub-Saharan Africa
- Demand for pork is significantly increasing
- Knowledge regarding *T. solium* cysticercosis is almost non-existing
- Meat inspection is either non-existing or inappropriate
- Open defecation is highly prevalent

However, no tools are available in sub-Saharan Africa

T. solium cysts in pork

Cyst in pig brain



"White nodules"




"Measly pork"



Photo: U. Braae

T. solium cysts in human brain



Symptoms like severe headache and epilepsy are very common




Cross-disciplinary Risk Assessment of Taenia solium Cysticercosis in Eastern & Southern Africa (CESA) 2006 - 2010

Conclusion

High prevalence of epilepsy, taeniosis, human cysticercosis and porcine cysticercosis were found in the studied communities of Mozambique and Tanzania

Lack of knowledge regarding the disease and its transmission is a major obstacle for control and prevention



Securing Rural Livelihoods through Improved Smallholder Pig Production in Mozambique and Tanzania (SLIPP) 2010 -2015

SLIPP USES A MULTI FACTORIAL APPROACH to address the complexity of changing not only the production system but also perceptions and prejudices. This involves awareness generation to improve overall knowledge and community based research to document effect and feasibility at village level to persuade stakeholders to adapt solutions long-term

SLIPP tools:

- Health education
- Pig management education
- Total pig confinement
- Anthelmintic treatment of pigs at strategic time points
- Pig management including model houses, water and feed