

UNIVERSITY OF COPENHAGEN



Bacterial community structure in High-Arctic snow and freshwater as revealed by pyrosequencing of 16S rRNA genes and cultivation

Møller, Annette K.; Søborg, Ditte A.; Abu Al-Soud, Waleed; Sørensen, Søren Johannes; Kroer, Niels

Published in:
Polar Research

DOI:
[10.3402/polar.v32i0.17390](https://doi.org/10.3402/polar.v32i0.17390)

Publication date:
2013

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Møller, A. K., Søborg, D. A., Abu Al-Soud, W., Sørensen, S. J., & Kroer, N. (2013). Bacterial community structure in High-Arctic snow and freshwater as revealed by pyrosequencing of 16S rRNA genes and cultivation. *Polar Research*, 32, [17390]. <https://doi.org/10.3402/polar.v32i0.17390>

Yu Y., Lee C., Kim J. & Hwang S. 2005. Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. *Biotechnology and Bioengineering* 89, 670–679.

Zhao J.S., Manno D. & Hawari J. 2009. *Psychrilyobacter atlanticus* gen. nov., sp. nov., a marine member of the phylum Fusobacteria that produces H₂ and degrades

nitramine explosives under low temperature conditions. *International Journal of Systematic and Evolutionary Microbiology* 59, 491–497.

Zwart G., Crump B.C., Agterveld M.P.K.V., Hagen F. & Han S.K. 2002. Typical freshwater bacteria: an analysis of available 16S rRNA gene sequences from plankton of lakes and rivers. *Aquatic Microbial Ecology* 28, 141–155.