



Genome sequences of three highly copper-resistant *Salmonella enterica* subsp. I serovar typhimurium strains isolated from pigs in Denmark

Qin, Yanan; Hasman, Henrik; Aarestrup, Frank Møller; Alwathnani, Hend A. ; Rensing, Christopher Günther T

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.01334-14](https://doi.org/10.1128/genomeA.01334-14)

Publication date:
2014

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Qin, Y., Hasman, H., Aarestrup, F. M., Alwathnani, H. A., & Rensing, C. G. T. (2014). Genome sequences of three highly copper-resistant *Salmonella enterica* subsp. I serovar typhimurium strains isolated from pigs in Denmark. *Genome Announcements*, 2(6), [e01334-14]. <https://doi.org/10.1128/genomeA.01334-14>

Genome Sequences of Three Highly Copper-Resistant *Salmonella enterica* subsp. I Serovar Typhimurium Strains Isolated from Pigs in Denmark

Yanan Qin,^a Henrik Hasman,^b Frank M. Aarestrup,^b Hend A. Alwathnani,^c Christopher Rensing^{a,d}

Department of Plant and Environmental Science, University of Copenhagen, Frederiksberg, Denmark^a; National Food Institute, Technical University of Denmark, Lyngby, Denmark^b; Department of Botany and Microbiology, King Saud University, Riyadh, Saudi Arabia^c; Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, China^d

***Salmonella typhimurium* is the causative agent of typhoid fever, which causes nearly 21.7 million illnesses and 217,000 deaths around the world each year. Here, we describe the draft genome sequences of the *Salmonella typhimurium* strains S7, S15, and S23, isolated from copper-fed pigs in Denmark and containing additional putative determinants conferring resistances to copper and other metals and metalloids.**

Received 12 November 2014 Accepted 19 November 2014 Published 24 December 2014

Citation Qin Y, Hasman H, Aarestrup FM, Alwathnani HA, Rensing C. 2014. Genome sequences of three highly copper-resistant *Salmonella enterica* subsp. I serovar Typhimurium strains isolated from pigs in Denmark. *Genome Announc.* 2(6):e01334-14. doi: 10.1128/genomeA.01334-14.

Copyright © 2014 Qin et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Christopher Rensing, chres@life.ku.dk.

Salmonella enterica subsp. I serovar Typhimurium (*S. typhimurium*) is one of the most important food-borne bacterial pathogens, with a broad host range, including food animals and humans, infecting 21.7 million people and causing 217,000 deaths annually (1–3). In addition, the magnitude of multidrug resistance in *Salmonella* and other pathogens at the human-animal and ecosystem interface has been a major concern globally.

Three *S. typhimurium* (S7, S15, S23) strains were isolated from copper-fed pigs as part of the Danish Integrated Antimicrobial Resistance Monitoring (DANMAP) surveillance program (4). Genomic DNA (gDNA) was purified from the isolates using the Easy-DNA extraction kit (Invitrogen), and DNA concentrations were determined using the Qubit dsDNA BR assay kit (Invitrogen). Whole-genome shotgun sequencing of *S. typhimurium* strains S7, S15, and S23 was performed using the Illumina MiSeq platform to generate 12,037,844, 12,039,576, and 12,061,925 paired-end reads, resulting in ~165-fold, ~170-fold, and ~168-fold coverage of the genomes, respectively. Genome assemblies were constructed *de novo* using CLCBio Genomic Workbench version 5.1 (5) (CLCBio, Denmark), which generated 146, 148, and 148 contigs, respectively.

The resulting contigs were uploaded into the RAST server (6,7) to predict the open reading frames (ORFs). In brief, the predicted ORFs were annotated by searching against clusters of SEED (8) databases. The draft genome size of *S. typhimurium* S7 is approximately 4,927,534 bp in length, with an average GC content of 52.0% and comprises 4,907 predicted coding sequences with an average length of 810 bp. *S. typhimurium* S15 is 4,914,811 bp in length, with an average GC content of 52.1%, and comprises 4,898 predicted coding sequences with an average length of 847 bp. *S. typhimurium* S23 is 4,914,318 bp in length, with an average GC content of 52.1%, and comprises 4,908 predicted coding sequences with an average length of 803 bp.

The three *S. typhimurium* genomes contain several copper-

resistance genes, including the Cu(I)-translocating P-type ATPase CopA, multicopper oxidase CueO, and a copper-responsive MerR-family transcriptional regulator CueR (9–11), which were also reported in *S. typhi* strains Ty2 and CT18 (12–14). All three strains contained a mobile 20-gene copper resistance determinant that contained the previously described *pco* and *sil* determinants (15, 16). There are two additional genes between these two determinants; *pcoG* encodes a putative M23B metalloproteinase, and *pcoF* encodes a putative copper binding protein.

The sequences also revealed genes encoding the *Salmonella*-specific P-type ATPase, *GolT*, *golBS*, and *gesABC* (17, 18, 19) genes. Most *Salmonella* strains lack an RND-type CusCFBA system; instead, CueP sequesters copper to neutralize toxicity (20–23). However, *S. typhimurium* (S7, S15, S23) contained both an RND type system and CueP. In addition, the genome also revealed a six-gene cluster *merEDAPTR* encoding proteins conferring mercury resistance and the five-gene *ars* operon *arsRDABC* (24).

Nucleotide sequence accession numbers. These whole-genome shotgun projects have been deposited in DDBJ/EMBL/GenBank under the accession numbers [JRGS00000000](https://ncbi.nlm.nih.gov/nucl/JRGS00000000), [JRGR00000000](https://ncbi.nlm.nih.gov/nucl/JRGR00000000), [JRGT00000000](https://ncbi.nlm.nih.gov/nucl/JRGT00000000). The version described in this paper is the first version. The BioProject designations for those projects are PRJNA260777, PRJNA260771, and PRJNA260778.

ACKNOWLEDGMENTS

This work was supported by the Center for Environmental and Agricultural Microbiology (CREAM) funded by the Villum Kann Rasmussen Foundation.

REFERENCES

- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. 2011. Foodborne illness acquired in the United States—major pathogens. *Emerg. Infect. Dis.* 17:7–15. <http://dx.doi.org/10.3201/eid1701.P11101>.
- Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, Jones

- TF, Fazil A, Hoekstra RM. 2010. The global burden of nontyphoidal *Salmonella* gastroenteritis. *Clin. Infect. Dis.* 50:882–889. <http://dx.doi.org/10.1086/650733>.
3. Crump JA, Mintz ED. 2010. Global trends in typhoid and paratyphoid fever. *Clin. Infect. Dis.* 50:241–246. <http://dx.doi.org/10.1086/649541>.
 4. Danish Integrated Antimicrobial Resistance Monitoring. 2005. DANMAP 2004—use of antimicrobial agents and the occurrence of antimicrobial resistance in bacteria from food animals, foods and humans in Denmark. DANMAP, Copenhagen.
 5. Yap KP, Gan HM, Teh CSJ, Baddam R, Chai LC, Kumar N, Tiruvayipati SA, Ahmed N, Thong KL. 2012. Genome sequence and comparative pathogenomics analysis of a *Salmonella enterica* serovar Typhi strain associated with a typhoid carrier in Malaysia. *J. Bacteriol.* 194:5970–5971. <http://dx.doi.org/10.1128/JB.01416-12>.
 6. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
 7. Meyer F, Paarmann D, D'ouza M, Olson R, Glass EM, Kubal M, Paczian T, Rodriguez A, Stevens R, Wilke A, Wilkening J, Edwards RA. 2008. The metagenomics RAST server—a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* 9:386. <http://dx.doi.org/10.1186/1471-2105-9-386>.
 8. Hemmerich C, Buechlein A, Podicheti R, Revanna KV, Dong Q. 2010. An Ergatis-based prokaryotic genome annotation web server. *Bioinformatics* 26:1122–1124. <http://dx.doi.org/10.1093/bioinformatics/btq090>.
 9. Zhu L, Elguindi J, Rensing C, Ravishankar S. 2012. Antimicrobial activity of different copper alloy surfaces against copper resistant and sensitive *Salmonella enterica*. *Food Microbiol.* 30:303–310. <http://dx.doi.org/10.1016/j.fm.2011.12.001>.
 10. Espariz M, Checa SK, Audero MEP, Pontel LB, Soncini FC. 2007. Dissecting the *Salmonella* response to copper. *Microbiology* 153: 2989–2997. <http://dx.doi.org/10.1099/mic.0.2007/006536-0>.
 11. Hodgkinson V, Petris MJ. 2012. Copper homeostasis at the host-pathogen interface. *J. Biol. Chem.* 287:13549–13555. <http://dx.doi.org/10.1074/jbc.R111.316406>.
 12. Deng W, Liou SR, Plunkett G, Mayhew GF, Rose DJ, Burland V, Kodoyianni V, Schwartz DC, Blattner FR. 2003. Comparative genomics of *Salmonella enterica* serovar Typhi strains Ty2 and CT18. *J. Bacteriol.* 185:2330–2337. <http://dx.doi.org/10.1128/JB.185.7.2330-2337.2003>.
 13. Holt KE, Parkhill J, Mazzoni CJ, Roumagnac P, Weill FX, Goodhead I, Rance R, Baker S, Maskell DJ, Wain J, Dolecek C, Achtman M, Dougan G. 2008. High-throughput sequencing provides insights into genome variation and evolution in *Salmonella* Typhi. *Nat. Genet.* 40:987–993. <http://dx.doi.org/10.1038/ng.195>.
 14. Parkhill J, Dougan G, James KD, Thomson NR, Pickard D, Wain J, Churcher C, Mungall KL, Bentley SD, Holden MTG, Sebahia M, Baker S, Basham D, Brooks K, Chillingworth T, Connor P, Cronin A, Davis P, Davies RM, Dowd L, White N, Farrar J, Feltwell T, Hamlin N, Haque A, Hien TT, Holroyd S, Jagels K, Krogh A, Larsen TS, Leather S, Moule S, O'Gaora P, Parry C, Quail M, Rutherford K, Simmonds M, Skelton J, Stevens K, Whitehead S, Barrell BG. 2001. Complete genome sequence of a multiple drug resistant *Salmonella enterica* serovar Typhi CT18. *Nature* 413:848–852. <http://dx.doi.org/10.1038/35101607>.
 15. Zimmermann M, Udagedara SR, Sze CM, Ryan TM, Howlett GJ, Xiao Z, Wedd AG. 2012. PcoE—a metal sponge expressed to the periplasm of copper resistance *Escherichia coli*. implication of its function role in copper resistance. *J. Inorg. Biochem.* 115:186–197. <http://dx.doi.org/10.1016/j.jinorgbio.2012.04.009>.
 16. Gupta A, Matsui K, Lo JF, Silver S. 1999. Molecular basis for resistance to silver cations in *Salmonella*. *Nat. Med.* 5:5183–5188.
 17. Kim JS, Kim MH, Joe MH, Song SS, Lee IS, Choi SY. 2002. The sctR of *Salmonella enterica* serovar Typhimurium encoding a homologue of MerR protein is involved in the copper-responsive regulation of *cuiD*. *FEMS Microbiol. Lett.* 210:99–103. <http://dx.doi.org/10.1111/j.1574-6968.2002.tb11166.x>.
 18. Checa SK, Espariz M, Audero ME, Botta PE, Spinelli SV, Soncini FC. 2007. Bacterial sensing of and resistance to gold salts. *Mol. Microbiol.* 63:1307–1318. <http://dx.doi.org/10.1111/j.1365-2958.2007.05590.x>.
 19. Pontel LB, Audero ME, Espariz M, Checa SK, Soncini FC. 2007. GolS controls the response to gold by the hierarchical induction of *Salmonella*-specific genes that include a CBA efflux-coding operon. *Mol. Microbiol.* 66:814–825. <http://dx.doi.org/10.1111/j.1365-2958.2007.05963.x>.
 20. Porcheron G, Garenaux A, Proulx J, Sabri M, Dozois CM. 2013. Iron, copper, zinc, and manganese transport and regulation in pathogenic Enterobacteria: correlations between strains site of infection and the relative importance of the different metal transport systems for virulence. *Front. Cell. Infect. Microbiol.* 3:90. <http://dx.doi.org/10.3389/fcimb.2013.00090>.
 21. Dupont CL, Grass G, Rensing C. 2011. Copper toxicity and the origin of bacterial resistance—new insights and applications. *Metallomics* 3:1109–1118. <http://dx.doi.org/10.1039/c1mt00107h>.
 22. Osman D, Waldron KJ, Denton H, Taylor CM, Grant AJ, Mastroeni P, Robinson NJ, Cavet JS. 2010. Copper homeostasis in *Salmonella* is atypical and copper-CueP is a major periplasmic metal complex. *J. Biol. Chem.* 285:25259–25268. <http://dx.doi.org/10.1074/jbc.M110.145953>.
 23. Pontel LB, Soncini FC. 2009. Alternative periplasmic copper-resistance mechanisms in Gram negative bacteria. *Mol. Microbiol.* 73:212–225. <http://dx.doi.org/10.1111/j.1365-2958.2009.06763.x>.
 24. Carlin A, Shi W, Dey S, Rosen BP. 1995. The ars operon of *Escherichia coli* confers arsenical and antimonial resistance. *J. Bacteriol.* 177:981–986.