



4,8-bisallyl-2,6-dimethylnaphthalene-1,5-diyl diacetate

Christensen, Jørn Bolstad; Sørensen, Jeanett N.; Schaumburg, Kjeld; Bechgaard, Klaus

Published in:
Molbank

DOI:
[10.3390/M812](https://doi.org/10.3390/M812)

Publication date:
2014

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Christensen, J. B., Sørensen, J. N., Schaumburg, K., & Bechgaard, K. (2014). 4,8-bisallyl-2,6-dimethylnaphthalene-1,5-diyl diacetate. *Molbank*, 2014(1), [M812]. <https://doi.org/10.3390/M812>

Short Note

4,8-Bisallyl-2,6-dimethylnaphthalene-1,5-diyl Diacetate

Jørn B. Christensen *, Jeanett N. Sørensen, Kjeld Schaumburg and Klaus Bechgaard

Department of Chemistry, University of Copenhagen, Thorvaldsensvej 40, DK-1871 Frederiksberg C, Denmark

* Author to whom correspondence should be addressed; E-Mail: jbc@chem.ku.dk.

Received: 4 December 2013 / Accepted: 8 January 2014 / Published: 13 January 2014

Abstract: 4,8-Diallyl-2,6-dimethylnaphthalene-1,5-diyl diacetate (**1**) which is a highly substituted naphthalene derivative has been synthesized in two steps starting from 2,6-dimethyl-1,5-naphthalenediol (**3**) using a modified Claisen-rearrangement.

Keywords: 1,2,4,5,6,8-hexasubstituted naphthalene; Claisen-rearrangement


Peri-substituted aromatic compounds [1] such as 1,4,5,8-tetrasubstituted naphthalenes are notoriously difficult to synthesize due to the short distance between the peri-positions and in compounds such as 1,4,5,8-tetramethyl naphthalene the methyl groups are interlocking giving rise to restricted rotation [2]. Peri-strain also affects the reactivity of the compounds as for example seen in 4,8-dibromo-1,5-dimethoxynaphthalene, which isomerizes to 2,7-dibromo-1,5-dimethoxy- and finally 2,6-dibromo-1,5-dimethoxynaphthalene upon even short heating in acetic acid [3].

1,4,5,8-tetrasubstituted naphthalenes can serve as starting materials for the synthesis of pyrenes and heterocyclic pyrenes such as dioxapyrenes, which are interesting compounds for conducting radical cation salts. We [4–7] and other groups [8–11] have previously reported different methodologies for the synthesis of 1,6-dioxa- and 1,6-dithiapyrenes from naphthalene derivatives and while the 1,6-dithiapyrenes can be synthesized from 1,5-disubstituted naphthalenes through an acid catalyzed ring closure, the synthesis of 1,6-dioxapyrenes requires either 1,2,5,6-tetrasubstituted naphthalenes or 1,4,5,8-tetrasubstituted naphthalenes as intermediates.

The Claisen-rearrangement is an efficient method for introducing substituents ortho to a phenol via the allyl aryl ether, however if the ortho-positions are blocked, the product is the corresponding 4-allyl substituted compound. [12] This led us to consider the double Claisen-rearrangement of 2,6-disubstituted bis allyl ethers of 1,5-naphthalenediols as a synthetic route to *bis*peri-substituted naphthols for subsequent uses for the synthesis of 1,6-dioxapyrenes.

2,6-Dimethyl-1,5-naphthalenediol (**3**) [4] was alkylated with allyl bromide in DMF with K_2CO_3 as the base to give the crude bisallyl ether (**2**), which was subjected to a modified Claisen-rearrangement [13] by reflux in a mixture of *N,N*-diethylaniline and acetic anhydride (Scheme 1). Under these conditions, the highly air sensitive tetrasubstituted naphthalenediol is immediately protected by acetylation to give a stable product, that can be purified without any special precautions.

Scheme 1. The synthesis of 4,8-Bisallyl-2,6-dimethylnaphthalene-1,5-diyl Diacetate.



4,8-Bisallyl-2,6-dimethylnaphthalene-1,5-diyl diacetate (1)

2,6-Dimethyl-1,5-dihydroxynaphthalene (**3**) [4] (10.2 g; 0.05 mol) and allyl bromide (10 mL; 14 g; 0.12 mol) was added to a degassed suspension of dry K_2CO_3 (0.15 mol) in DMF (50 mL). The mixture was stirred at $20^\circ C$ under N_2 overnight. The reaction mixture was poured into water (500 mL) and extracted with diethyl ether (100 mL). The organic phase was dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude bis(allyl ether) was dissolved in a mixture of *N,N*-diethyl aniline (50 mL) and acetic anhydride (20 mL) and refluxed under N_2 for 3 days ($200-210^\circ C$). The product separated upon cooling to room temperature was isolated by filtration, washed with petroleum ether (Bp. $35^\circ C$) until free of *N,N*-diethyl aniline and acetic acid and air dried to give **1** as an off-white powder.

Yield: 9.6 g (55%).

An analytical sample was crystallized from EtOH. Mp. $166-168^\circ C$.

Calcd. for C₂₂H₂₄O₄, C, 74.98; H, 6.86. Found: C, 74.56; H, 6.70.

¹H-NMR (500 MHz; CDCl₃): δ 7.08 (s, 2 H); 6.07 (m; 2H); 5.03 (m; 2H); 4.92 (m; 2H); 3.78 (m, 4H); 2.30 (s, 6H); 2.13 (s, 6H).

¹³C-NMR (125 MHz; CDCl₃): δ 168.9; 143.9; 137.7; 132.5; 132.1; 127.4; 126.9; 116.2; 40.3; 21.5; 16.7.

Acknowledgments

We would to thank University of Copenhagen for technical support.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Balasubramanian, V. Peri interaction in naphthalene derivatives. *Chem. Rev.* **1966**, *66*, 567–641.
2. Imashiro, F.; Takegoshi, K.; Saika, A.; Taira, Z.; Asahi, Y. Molecular structure of 1,4,5,8-tetramethylnaphthalene and in-plane molecular rotation of some methyl-substituted naphthalenes in solids. *J. Am. Chem. Soc.* **1985**, *107*, 2341–2346.
3. Sylvester-Hvid, K.; Sørensen, J.; Schaumburg, K.; Bechgaard, K.; Christensen, J.B. Preparation of some 4,8-Dimethoxy-diformylnaphthalenes. *Synth. Commun.* **1993**, *23*, 1905–1914.
4. Christensen, J.B.; Johannsen, I.; Bechgaard, K. Synthesis and properties of substituted 1,6-dioxapyrene donors. *J. Org. Chem.* **1991**, *56*, 7055–7058.
5. Mortensen, M.B.; Schlüter, A.; Sørensen, J.; Christensen, J.B. A convenient preparation of 1,6-dioxapyrene-2,7-dicarboxylic acid diethylester. *Acta Chem. Scand.* **1997**, *51*, 807–809.
6. Christensen, J.B.; Bechgaard, K. The chemistry of 1,6-dioxapyrenes part 3 [1,2]: Scope and limitations of an acid catalyzed ring-closing reaction. *J. Het. Chem.* **2003**, *40*, 757–761.
7. Tyson, D.S.; Fabrizio, E.F.; Panzner, M.J.; Kinder, J.D.; Buisson, J.-P.; Christensen, J.B.; Meador, M.A. Synthesis, characterization, and optical properties of a cyano-functionalized 4,5,9,10-tetraaryl-1,6-dioxapyrene. *J. Photochem. Photobiol. A* **2005**, *172*, 97–107.
8. Tilak, B.D. Synthesis of thiophenes and thiapyrans. IV. Thiophenes and thiapyrans from naphthalenethiols. *Proc. Indian Acad. Sci.* **1951**, *33A*, 71–77.
9. Nakasuji, K.; Kubota, H.; Kotani, T.; Murata, I.; Saito, G.; Enoki, T.; Imaeda, K.; Inokuchi, H.; Honda, M.; Katayama, C.; *et al.* Novel peri-condensed Weitz-type donors: Synthesis, physical properties, and crystal structures of 3,10-dithiapyrene (DTPR), 1,6-dithiapyrene (DTPY), and some of their CT complexes. *J. Am. Chem. Soc.* **1986**, *108*, 3460–3466.
10. Nakasuji, K.; Sasaki, M.; Kotani, T.; Murata, I.; Enoki, T.; Imaeda, K.; Inokuchi, H.; Kawamoto, A.; Tanaka, J. Methylthio- and ethanediyldithio-substituted 1,6-dithiapyrenes and their charge-transfer complexes: New organic molecular metals. *J. Am. Chem. Soc.* **1987**, *109*, 6970–6975.
11. Miyazaki, E.; Morita, Y.; Nakasuji, K. 2-Iodo-1,6-dithiapyrene: Syntheses, crystal structures and physical properties of CT complexes and salt. *Polyhedron* **2005**, *24*, 2632–2638.

12. Tarbell, D.S. The Claisen Rearrangement. In *Organic Reactions*; Wiley: New York, NY, USA, 1944; Volume 2, pp. 1–48.
13. Fieser, L.F.; Campbell, W.P.; Fry, E.M. Synthesis of quinones related to Vitamins K₁ and K₂. *J. Am. Chem. Soc.* **1939**, *61*, 2206–2218.

© 2014 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).