



Growth performance, meat quality and carcass composition of broilers fed rapeseed-enriched diets

Karlsson, Anders H; Bredie, Wender; Petersen, Mikael Agerlin; Nielsen, Belinda; Danielsen, Bente; Petersen, J. S.; Katholm, A.; Dickow, Mia Schødt

Publication date:
2015

Document version
Peer reviewed version

Citation for published version (APA):
Karlsson, A. H., Bredie, W., Petersen, M. A., Nielsen, B., Danielsen, B., Petersen, J. S., ... Dickow, M. S. (2015). *Growth performance, meat quality and carcass composition of broilers fed rapeseed-enriched diets*. Poster session presented at ICoMST 2015, Clermont-Ferrand, France.

GROWTH PERFORMANCE, MEAT QUALITY AND CARCASS COMPOSITION OF BROILERS FED RAPESEED-ENRICHED DIETS

Anders H. Karlsson¹, W. Bredie, M.A. Petersen¹, B. Nielsen¹, B. Danielsen¹, J.S. Petersen², A. Katholm³, M. Schødt Dickow³

¹Dept. of Food Science, University of Copenhagen, Rolighedsvej 26, 1958, Frederiksberg C, Denmark, ²SEGES, Agro Food Park 15, 8200 Aarhus N, Denmark, ³DLG, Vesterbrogade 4A, 1620 København V, Denmark

Abstract

This paper describes the effect of a diet containing 15% rapeseed meal on growth performance, carcass composition, fatty acid composition, volatiles and sensory meat quality attributes on broilers pectoralis meat compared to 0%. The overall results show that the broilers performed well on the rapeseed meal diet used.

Key Words – Rapeseed-enriched broiler feed, Growth performance, Meat and carcass quality.

Introduktion

The majority of European broiler production's need for protein is today covered by imported soybean meal. Using imported soy in animal production is questioned, and limiting its use for European meat production is therefore in focus. Replacing imported soy protein by domestic protein, could be one way to strengthen the European broiler brand. Rapeseed is a protein crop that is grown in many EU states, including Denmark, and there are therefore great opportunities to increase its use in broiler production. Furthermore, replacing soybean meal with rapeseed could also result in cheaper broiler feed. The aim of the present study was to compare the growth performance, foot pad quality, carcass composition and meat flavour in broilers fed diets including 0, or 15% rapeseed meal from days 11 to day 32. The broilers were slaughtered at day 34.

Material and Methods

Birds and production: In total 720 broilers of the breed Ross 308 were divided into 12 pens, each containing 60 broilers. There were 30 males and 30 females in each pen. The birds were raised on a conventional broiler farm.

Broiler diets: The two diets tested were both based on soybean meal, wheat and corn and both contained the coccidiostat salinomycin. The control diet contained 0% rapeseed meal, while the test diet contained 15% rapeseed meal. Diets were optimized to be equal regarding energy and protein. The rapeseed in the test diet replaced a part of the wheat, soy, and maize gluten meal. From day 6 to the end of the experiment, the feed was diluted with whole wheat starting with an inclusion rate of 6 % and then increasing to a whole wheat inclusion rate of 24%. The last 2 days before slaughter, the birds were fed a finishing diet without any rapeseed meal and without any coccidiostat.

Dietary fatty acid composition: The lipids were extracted using chloroform/methanol solution (2:1, v/v), followed by homogenization. The chloroform phase was used for fatty acid composition (FAME) analysis. The methylated fatty acids were analysed on the gas chromatograph using an Omega wax column and FID detection. The data was analysed using Chemstation software (Agilent Technologies) and the fatty acid methyl esters were identified by comparing retention times with known standards. The results were expressed as % fatty acid of the total content of detected fatty acids.

Performance and carcass quality: For carcass studies, 7 males and 7 females were randomly marked in each of the boxes at day 0. All the broilers in each pen were weighed during their time in the box on days 0 (start), 7, 14, 28 and 32. Feed consumption, mortality-rate and foot pad health were also registered. Feed conversion corrected for mortality (FCR-corr.) was calculated based on a standard curve for daily weight gain. On day 33, the broilers were placed in slaughter boxes, and on the next day (day 34) they arrived at the commercial slaughter plant "Sødam Øko Fjerkræslægteri" where they were electrically stunned and slaughtered. On Day 36, the carcasses were cut according to the standard method described by Darré and Claudi-Magnussen et al. (DMRI, unpublished).

Analysis of volatiles: When meat samples were cooked for sensory testing, extra samples were prepared simultaneously for analysis of volatiles. For each of the two test diets, 12 samples stored for 6 days were analysed together with four samples stored for either 7, 8 or 9 days (48 samples in total). Dynamic headspace sampling followed by thermal desorption of traps to a GC-MS system was applied (procedure slightly modified from [1]).

Sensory evaluation: A descriptive sensory analysis was carried out with a trained sensory panel (n=10). In order to simulate the fresh meat quality typically found on shop display, the samples from both feeding regimes were stored at +2°C for 6, 7, 8 and 10 consecutive days prior to cooking. The pectoralis major muscles were partitioned in a left and right part. For each broiler one part was frozen (control day 6) and the other part stored refrigerated at +2°C

until the respective storage day. On the day of assessment the frozen control samples were thawed out and together with the refrigerated samples cooked sous-vide in a water bath to a core temperature of 63°C. Samples were subsequently held at 10 min (60°C) before unpacking, slicing and serving to the panel. All samples were evaluated in four replicates over four sessions. The day 6 control (with and without rapeseed-enriched diet) were present in all sessions. Within each session the sample presentation order was randomised.

Data analysis: The data was analysed using ANOVA (Proc. Mixed, SAS, 9.4) where the fixed effects of diet and sex were included in the model. For the analyses of performance during growth the model included the fixed effect of Day of measurement. Least-squares means and standard error of the means were calculated. A p-value less than 0.05 was considered as a significant difference.

Results and Discussion

Performance and carcass quality: Using rapeseed in the diet had no effect on the performance indicators registered (Tab. 1-2) or on carcass quality attributes (Tab. 3). Expected differences between sex were also found in this study (Tab. 4).

All broilers performed well and the health status was good. The foot pad score was very low in both treatments indicating that the test diet did not have any negative impact on the digestion. FCR was highest on the test diet and overall production performance was best on the control diet.

Replacing part of soybean meal with 15% rapeseed meal did not have an influence on total fat content of the meat, as well as the concentrations of the saturated fatty acids found (C16:0 and C18:1) in the diets. The concentration of MUFA (C18:1) was reduced when using rapeseed meal. Regarding concentrations of PUFA, C18:2 was increased and C18:3 was reduced in the diet when using rapeseed meal (Tab. 5). It is well-known that the fatty acid composition found in meat from birds, is to a very high degree reflecting the fatty acid composition in the diet. Therefore it was expected to find the similar fat profile in the meat when using rapeseed meal in the bird's diet.

Sensory study: The effect of broiler diet showed a significant effect at day 10 for **crumbliness** (p=.025) with the rapeseed fed broilers scoring significantly higher on this attribute. The effect of storage days at each diet regime was significant for the attribute **sweet** (p=.024) for the control diet only. In this case the sweetness was scored significantly higher after ageing the meat refrigerated for 10 days. When modelling within the sensory evaluation sessions more sensory differences could be observed, but these could not be clearly separated from the variability in the assessment of the control samples (Day 6). Therefore the overall measurable sensory differences were judged to be relatively small with only effects occurring at the prolonged (Day 10) refrigerated storage.

Volatiles: In total, 69 volatile compounds could be detected in the cooked meat samples. No effect was seen from storage prior to cooking, but 27 compounds were significantly influenced by the broilers' diet. Almost all of these compounds are related to lipid oxidation (aldehydes, alcohols, ketones and alkanes; Fig. 1), and they had all higher levels in meat from broilers given the 15%-rapeseed meal diet. Many of the compounds are suspected to cause off-flavour in meat, for example hexanal [2], but when comparing with the results of the sensory test, it can be concluded that the levels were not high enough to be detected sensorily.

Conclusion

Performance: The broilers performed well on a 15% rapeseed meal diet. **Volatiles:** Increased oxidation after feeding rapeseed diets, but not enough to be detected sensorily. **Fatty acid composition:** Feeding rapeseed did not influence total fat content, as well as the contents of the saturated fatty acids C16:0 and C18. MUFA (C18:1) was reduced in the diet containing rapeseed. PUFA C18:2 was increased, and C18:3 was reduced in the rapeseed meal based diet. These results are expected as the fatty acid profile found in chicken meat reflect the fatty acids found in the diet.

Sensory: The overall measurable sensory differences were relatively small with only effects occurring at the prolonged refrigerated storage.

Based on these results, there seem to be no **economical advantage** to use 15% rapeseed meal in the broiler diet as a replacement for soybean meal. However, the price for rapeseed and soy varies over time, and in this light, the results can be of importance. If the price difference between rapeseed meal and soybean meal will be large, the overall production economy could benefit from the use of up to 15 % rapeseed meal in the diet.

Table 1. The effect of diet overall and at 32 days age on broiler performance (LS-means and SE).

Attribute	Diet		SE	p-value
	No-Rapeseed	15%-Rapeseed		
Mortality, %	3.4	3.9	0.62	0.97
Mortality-Day32, %	3.9	4.7	1.42	0.71
FCR, kg/kg	1.14 ^a	1.15 ^b	0.003	0.017*
FCR-Day32, kg/kg	1.42	1.44	0.007	0.055
FCR-corr, kg/kg	1.13 ^a	1.14 ^b	0.003	0.050*
FCR-corr-Day32, kg/kg	1.41	1.43	0.008	0.138
Food pad score _{Day28}	0.4	1.6	1.2	0.48
Weight-Day32, g	1957	1906	29	0.24

Table 2. The overall effect of age on performance (LS-means and SE).

Attribute	Day				SE	p-value
	7	14	28	32		
Mortality, %	2.8	3.5	4.2	4.3	12.93	0.59
Weight, g	171 ^a	450 ^b	1574 ^c	1931 ^d	20.8	<0.0001***
FCR, kg/kg	0.75 ^a	1.06 ^b	1.34 ^c	1.43 ^d	0.004	<0.0001***
FCR-corr, kg/kg	0.74 ^a	1.05 ^b	1.33 ^c	1.42 ^d	0.005	<0.0001***

Table 3. The overall effect of diet on carcass quality (LS-means and SE).


Attribute	Diet		SE	p-value
	No-Rapeseed	15%-Rapeseed		
Live wt, g	1982	1916	35.5	0.22
Carcass wt, g	1438	1377	28.0	0.15
Dressing-%	72.2	72.0	0.3	0.55
Fillet, g	486.5	498.6	3.5	0.99
Fillet, %	34.3	34.4	0.2	0.71
Thigh, g	250.6	251.4	1.9	0.78
Drumstick, g	179.7	178.9	1.2	0.70
Wings, g	123.4	124.7	0.7	0.21
Neck+wingpoints, g	17.3 ^a	18.2 ^b	0.3	0.045*
Skin+Skinfat, g	45.5 ^a	42.9 ^b	0.8	0.047*

Table 4. The overall effect of sex on slaughter quality (LS-means and SE).

Attribute	Sex		SE	p-value
	Male	Female		
Live wt, g	2085 ^a	1814 ^b	31.2	<0.0001***
Carcass wt, g	1509 ^a	1306 ^b	25.0	<0.0001***
Dressing-%	71.6 ^a	72.6 ^b	0.3	0.0033**
Fillet, g	474.4 ^a	498.6 ^b	3.5	<0.0001***
Fillet, %	33.3 ^a	35.3 ^b	0.2	<0.0001***
Thigh, g	252.9	249.1	1.8	0.12
Drumstick, g	184.0	174.6	1.2	<0.0001***
Wings, g	125.6 ^a	122.5 ^b	0.7	0.0058*
Neck+wingpoints, g	18.5 ^a	17.1 ^b	0.2	<0.0001***
Skin+Skinfat, g	41.8 ^a	46.6 ^b	0.9	0.0003***

Table 5. Effect of diet on fatty acid composition of the meat (LS-means and SE)

Diet	C16:0, % Palmitic	C18:0, % Stearic	C18:1, % Oleic	C18:2, % Linoleic	C18:3, % α-linoleic	Total fat, %
0% rapeseed	13.43±0.03	1.72±0.12	28.36±0.15	51.39±0.06	5.10±0.01	7.20±0.70
15% rapeseed	13.38±0.03	1.40±0.12	25.41±0.15	54.91±0.06	4.90±0.01	6.30±0.70
p-value	0.3906	0.2026	0.0051**	0.0006***	0.0097**	0.4573



Acknowledgements

This project was funded by a grant in 2013 from the Danish Poultry Levy Fund (*Fjerkræafgiftsfonden*) and the Programme for Green Development and Demonstration (*GUDP*).