



C-di-GMP regulates *Pseudomonas aeruginosa* stress response to tellurite during both planktonic and biofilm modes of growth

Chua, Song Lin; Sivakumar, Krishnakumar; Rybtke, Morten Levin; Yuan, Mingjun; Andersen, Jens Bo; Nielsen, Thomas E.; Givskov, Michael; Tolker-Nielsen, Tim; Cao, Bin; Kjelleberg, Staffan; Yang, Liang

Published in:
Scientific Reports

DOI:
[10.1038/srep10052](https://doi.org/10.1038/srep10052)

Publication date:
2015

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Chua, S. L., Sivakumar, K., Rybtke, M. L., Yuan, M., Andersen, J. B., Nielsen, T. E., ... Yang, L. (2015). C-di-GMP regulates *Pseudomonas aeruginosa* stress response to tellurite during both planktonic and biofilm modes of growth. *Scientific Reports*, 5, [10052]. <https://doi.org/10.1038/srep10052>

SCIENTIFIC REPORTS



OPEN

C-di-GMP regulates *Pseudomonas aeruginosa* stress response to tellurite during both planktonic and biofilm modes of growth

Received: 23 January 2014

Accepted: 20 March 2015

Published: 20 May 2015

Song Lin Chua^{1,2}, Krishnakumar Sivakumar^{1,3}, Morten Rybtke⁴, Mingjun Yuan¹, Jens Bo Andersen⁴, Thomas E. Nielsen¹, Michael Givskov^{1,4}, Tim Tolker-Nielsen⁴, Bin Cao^{1,5}, Staffan Kjelleberg^{1,6} & Liang Yang^{1,7}

Stress response plays an important role on microbial adaptation under hostile environmental conditions. It is generally unclear how the signaling transduction pathway mediates a stress response in planktonic and biofilm modes of microbial communities simultaneously. Here, we showed that metalloid tellurite (TeO_3^{2-}) exposure induced the intracellular content of the secondary messenger cyclic di-GMP (c-di-GMP) of *Pseudomonas aeruginosa*. Two diguanylate cyclases (DGCs), SadC and SiaD, were responsible for the increased intracellular content of c-di-GMP. Enhanced c-di-GMP levels by TeO_3^{2-} further increased *P. aeruginosa* biofilm formation and resistance to TeO_3^{2-} . *P. aeruginosa* $\Delta\text{sadC}\Delta\text{siaD}$ and $\text{PAO1}/p_{\text{lac}}\text{-yhjH}$ mutants with low intracellular c-di-GMP content were more sensitive to TeO_3^{2-} exposure and had low relative fitness compared to the wild-type PAO1 planktonic and biofilm cultures exposed to TeO_3^{2-} . Our study provided evidence that c-di-GMP level can play an important role in mediating stress response in microbial communities during both planktonic and biofilm modes of growth.

Microorganisms display a striking ability to adapt to unfavorable conditions such as exposure to UV radiation, heavy metals and antibiotic treatments, by inducing stress responses and forming surface-attached biofilms^{1,2}. Biofilms consist of microbial cells embedded in their self-produced extracellular polymeric substances (EPS). The EPS can account for up to 90% of the biofilm biomass and serve as physical barriers to protect biofilm cells³. Hence, biofilms dramatically increase the tolerance of bacterial cells towards environmental stress and immune attack during the course of infections^{4,5}. Extensive intercellular communication and interactions have been observed within biofilms, and cells with distinct physiology may play different roles under stress conditions^{6–8}.

Bis-(3'-5')-cyclic dimeric guanosine monophosphate (C-di-GMP) plays an important role in biofilm formation of a wide range of bacteria⁹. Bacterial intracellular c-di-GMP content is determined by diguanylate cyclases (DGCs) that catalyze the formation of c-di-GMP and phosphodiesterases (PDEs), which degrade c-di-GMP⁹. When intracellular c-di-GMP content is high, bacterial cells reduce motility and

¹Singapore Centre on Environmental Life Sciences Engineering (SCElse), Nanyang Technological University, Singapore 637551. ²NUS Graduate School of Integrative Sciences and Engineering, National University of Singapore, Singapore 117543. ³Interdisciplinary Graduate School, Nanyang Technological University, Singapore 637551. ⁴Costerton Biofilm Center, Department of International Health, Immunology and Microbiology, University of Copenhagen, 2200 København N, Denmark. ⁵School of Civil and Environmental Engineering, Nanyang Technological University, Singapore 639798. ⁶Center for Marine Bio-Innovation and School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney 2052, Australia. ⁷School of Biological Sciences, Nanyang Technological University, Singapore 637551. Correspondence and requests for materials should be addressed to B.C. (email: bincao@ntu.edu.sg) or L.Y. (email: yangliang@ntu.edu.sg)

increase synthesis of EPS matrix, resulting in biofilm formation^{10,11}. In contrast, biofilm cells increase their motility and disperse from biofilms when the intracellular c-di-GMP content is low^{12,13}. C-di-GMP signaling can be induced by stress conditions such as antimicrobial exposure^{14,15}. The impact of c-di-GMP on mediating stress response by microbial communities during both planktonic and biofilm modes of growth remains unclear.

Anthropogenic activities have resulted in serious metal(loid) pollution, especially in industrialized countries and regions. The natural ecosystems are often direct or indirect recipients of toxic metal(loid)s such as TeO_3^{2-} . Many environmental bacteria including *Pseudomonas aeruginosa* are capable of surviving in the presence of TeO_3^{2-} at low concentrations by reducing TeO_3^{2-} to Te(0) nanomaterials, as a result of either detoxification, redox maintenance or respiration^{16–19}. Although the toxic effects of metal(oid)s on environmental microorganisms at individual cell levels have been extensively studied²⁰, little is known about the impacts of metal(loid)s on bacterial social behaviours²¹.

In the present study, we investigated the role of c-di-GMP in mediating stress responses by the opportunistic pathogen *Pseudomonas aeruginosa* to a toxic metalloid, tellurite (TeO_3^{2-}). TeO_3^{2-} is highly toxic to most microbes and had been previously described by Alexander Fleming as an antimicrobial agent²². Bacterial cells take up TeO_3^{2-} and subsequently reduce it to tellurium nanoparticles, which can be easily tracked by the black precipitates on the bacterial cell surface. Quantification of intracellular c-di-GMP and proteomic analysis indicated that c-di-GMP levels were induced by TeO_3^{2-} exposure, which enhanced *P. aeruginosa* TeO_3^{2-} resistance and biofilm formation. SadC and SiaD were found to be important in the induction of c-di-GMP by TeO_3^{2-} exposure. We showed that mutants with low intracellular c-di-GMP content could be outcompeted by the wild-type strain from biofilm and planktonic cultures under metalloid stress condition.

Results

Stress responses of *P. aeruginosa* to TeO_3^{2-} induced c-di-GMP signaling. Cultivation of different bacterial species in the presence of sub-lethal concentrations of antimicrobial agents is a widely employed method to investigate their stress responses^{23–25}. The MIC of *P. aeruginosa* to TeO_3^{2-} is 100 $\mu\text{g}/\text{ml}$ in ABTGC medium. Large aggregates (approximately 1–3 mm) were formed when *P. aeruginosa* was grown in ABTGC media containing 10 $\mu\text{g}/\text{ml}$ TeO_3^{2-} at 37 °C (Fig. 1a). Further analysis of the TeO_3^{2-} -induced aggregates by field emission scanning electron microscopy (FE-SEM) and energy-dispersive X-ray spectroscopy (EDX) revealed the presence of tellurium-containing precipitates around the bacterial cells (Fig. 1b,c). No tellurium-containing precipitates were observed for *P. aeruginosa* cells growing in medium without TeO_3^{2-} . Thus, the tellurium-containing precipitates might generate conditions of membrane-associated stress for *P. aeruginosa* cells.

TeO_3^{2-} and oxyanions such as selenate/selenite are well known to exert their toxic effects on microorganisms via generation of reactive oxygen species (ROS)^{26,27}. We measured the generation of ROS by *P. aeruginosa* cells exposed to sub-lethal concentrations of TeO_3^{2-} as well as SeO_3^{2-} and SeO_4^{2-} by using the OxiSelect™ *in vitro* ROS/RNS assay kit. As anticipated, exposure of *P. aeruginosa* cells to the TeO_3^{2-} , SeO_3^{2-} and SeO_4^{2-} significantly increased their cytoplasmic ROS content regardless of the nutrient conditions (Fig. 1d).

Proteomic analysis of TeO_3^{2-} stressed *P. aeruginosa* cells. Oxidative stress response by *P. aeruginosa* leading to aggregate formation, recently reported to resemble the biofilm physiology²⁸ has not been documented. We thus investigated the overall impact of TeO_3^{2-} on *P. aeruginosa* cells using a comparative proteomic approach for cells cultivated with and without 10 $\mu\text{g}/\text{ml}$ TeO_3^{2-} .

Using a p-value cut-off of 0.05 and a fold change cut-off of 5 (as described in the Materials and Methods), 129 proteins were significantly affected by TeO_3^{2-} exposure with 64 proteins upregulated (Supplementary table 1) and 65 proteins being down-regulated (Supplementary table 2).

The expression of several of outer membrane associated proteins was induced by TeO_3^{2-} treatment, including OprQ (PA2760, 28.8-fold), OprI precursor (PA2853, 15-fold), probable outer membrane protein precursor (PA2391, 10.9-fold), OprM (PA0427, 10.5-fold), OprL precursor (PA0973, 9.8-fold), OprD precursor (PA0958, 9.8-fold), OprB (PA3186, 9.7-fold) and OprC (PA3790, 8.1-fold) (Supplementary table 1). The membrane transporter CdrB of the large extracellular protein CdrA²⁹ was induced 25.8-fold by exposure to TeO_3^{2-} (Supplementary table 1). CdrAB expression has been used as a c-di-GMP indicator³⁰ and reported to promote biofilm formation and auto-aggregation in a Psl polysaccharide dependent manner²⁹, and co-immunoprecipitation experiments have clearly shown that CdrA binds to Psl²⁹. HPLC analysis showed that *P. aeruginosa* PAO1 cultivated in ABTGC medium with 10 $\mu\text{g}/\text{ml}$ TeO_3^{2-} treatment had a higher relative intracellular c-di-GMP concentration compared to untreated control samples (approximately 2.5-fold) (Fig. 1e).

SadC and SiaD contribute to c-di-GMP induction by TeO_3^{2-} . CdrAB belongs to a family of bacterial proteins secreted by the two-partner secretion system³¹. Recently, two other members of this family, XacFhaB from *Xanthomonas axonopodis* pv. Citri and FHA from *Bordetella pertussis* have also been implicated in biofilm formation^{32,33}. These large inter-bacterial adhesins may play a key role in establishing structured biofilm communities under stress conditions. The *cdrA* promoter is positively regulated by the c-di-GMP concentration, and the expression of $P_{cdrA}\text{-gfp}$ has been recently used as a biosensor




Figure 1. Aggregates formed by *P. aeruginosa* wild-type PAO1 in ABTGC medium with and without 10 µg/ml TeO₃²⁻ under shaking condition after 1 d (a). Aggregates formed in TeO₃²⁻ containing medium were analyzed by FE-SEM (b) and energy-dispersive X-ray spectroscopy (c). Arrows in the FE-SEM image indicate the bacterial cell and nanoparticles on the cell surface. ROS generation by *P. aeruginosa* PAO1 cells after exposure to sub-lethal concentration of TeO₃²⁻, SeO₄²⁻ and SeO₃²⁻ (d). Relative intracellular c-di-GMP content of PAO1 cultures in ABTGC medium with and without 10 µg/ml TeO₃²⁻ was quantified by HPLC (e). Means and standard deviations of three replicates are shown. Student's *t*-test was performed for testing differences between groups. * *P* < 0.05.

of the intracellular content of c-di-GMP in *P. aeruginosa*³⁰. We tested the expression of the P_{cdrA}-gfp reporter in *P. aeruginosa* cultures with and without the presence of TeO₃²⁻ and found that TeO₃²⁻ exposure significantly increased the expression of fluorescence in a dose dependent manner (Fig. 2a). This result is in accordance with our HPLC quantification and indicates that TeO₃²⁻ exposure increases the intracellular content of c-di-GMP and that TeO₃²⁻ induced aggregates might carry physiological traits similar to those of biofilms.

Recently, both SadC and SiaD, were shown to be able to transduce an extracellular signal generated by the toxic detergent SDS and catalyze synthesis of c-di-GMP for facilitating biofilm formation by *P. aeruginosa*^{34,35}. The defect environmental signaling Δ*sadC* and Δ*siaD* mutants were severely impaired in expression of the P_{cdrA}-gfp reporter in the presence of TeO₃²⁻ (Fig. 2a). SiaD appears to be more important than SadC for P_{cdrA}-gfp induction by TeO₃²⁻ since the Δ*sadC* mutant still displayed a slight induction of P_{cdrA}-gfp by TeO₃²⁻ (Fig. 2a).

Exopolysaccharides are the major EPS components of *P. aeruginosa* biofilms and are well known to be induced by high intracellular c-di-GMP content in *P. aeruginosa*. We examined the expression of a lacZ-based biosensor of the Pel synthesis operon (mini-CTX-*pel-lacZ*³⁶) in *P. aeruginosa* strains under TeO₃²⁻ stress. As with P_{cdrA}-gfp fusion, the expression of the *pel-lacZ* fusion was induced by TeO₃²⁻ treatment, with SiaD essential for this induction (Fig. 2b). However, there was a slight induction of the *pel-lacZ* fusion by tellurite even in the Δ*sadC*Δ*siaD* double mutant (Fig. 2b).

Consistent with our observation of TeO₃²⁻-induced aggregation, *P. aeruginosa* grown in the presence of TeO₃²⁻ formed more biofilms than cells grown without TeO₃²⁻ (Fig. 3). The induction of biofilm formation was dependent on the presence of Pel and Psl polysaccharides (Fig. 3).

Induction of c-di-GMP confers a growth advantage under tellurite exposure during planktonic cultures. Since c-di-GMP signaling was induced by TeO₃²⁻ exposure, we examined whether induction of c-di-GMP signaling would confer a growth advantage of *P. aeruginosa* under TeO₃²⁻ exposure. There was no growth defect of Δ*sadC*, Δ*siaD* and Δ*sadC*Δ*siaD* mutants under normal growth condition as compared to PAO1 control (Fig. 4a). However, the *P. aeruginosa* Δ*sadC*, Δ*siaD* single or double mutants were more sensitive to TeO₃²⁻ (Fig. 4b). Similarly, the PAO1/*p_{lac}-yhjH* mutant, which




Figure 2. Expression of biosensor $P_{cdrA-gfp}$ (a) and $P_{pel-lacZ}$ (b) by *P. aeruginosa* strains in ABTGC medium with and without the presence of 10 µg/ml TeO_3^{2-} . The $P_{cdrA-gfp}$ expression was shown as relative fluorescence units (RFU) per OD_{600} . The $P_{pel-lacZ}$ expression was shown as Miller Unit. Means and standard deviations of three replicates are shown. Student's *t*-test was performed for testing differences between groups. * $P < 0.05$.

contains a PBBRMCS-2 plasmid with a constitutively expressed phosphodiesterase gene *yhjH* fused to and expressed by the *lac* promoter and thus has a low intracellular of *c*-di-GMP content¹², was also more sensitive to TeO_3^{2-} (Fig. 4). These results showed that intracellular *c*-di-GMP content determines the tolerance of *P. aeruginosa* to TeO_3^{2-} exposure during planktonic cultures.

Low intracellular *c*-di-GMP mutants lose fitness under stress during both planktonic and bio-film modes of growth. When *cfp*-tagged PAO1 and *yfp*-tagged $\Delta sadC\Delta siaD$ mutant strains were combined 1:1 (vol/vol) for planktonic co-cultivation experiments, the wild-type showed higher survival rates and gained a higher level of relative fitness than the $\Delta sadC\Delta siaD$ mutant in the presence of TeO_3^{2-} than without TeO_3^{2-} (Fig. 5a). Since diverse phenotypic and genotypic variants coexist in bacterial biofilms^{37,38}, we tested whether TeO_3^{2-} exposure-induced biofilm formation by high *c*-di-GMP containing cells would lead to protection of mutants with low intracellular *c*-di-GMP content in co-cultures. Here, PAO1 displayed a higher relative fitness than the $\Delta sadC\Delta siaD$ mutant in biofilm co-cultures with and without the presence of TeO_3^{2-} (Fig. 5b). However, the relative fitness of $\Delta sadC\Delta siaD$ compared to PAO1 in biofilm co-cultures was slightly higher with the presence of TeO_3^{2-} than in its absence (Fig. 5b). This suggests TeO_3^{2-} could potentially induce expression of other DGC harboring proteins in the $\Delta sadC\Delta siaD$ mutant and partly restore the intracellular *c*-di-GMP levels and biofilm formation.




Figure 3. Biofilm formation by *P. aeruginosa* PAO1, $\Delta pelA$, $\Delta pslBCD$ and $\Delta pelA\Delta pslBCD$ in medium containing 0, 10, 25 and 50 $\mu g/ml$ TeO_3^{2-} under static conditions after 1 d incubation. Biofilms were firstly stained with 0.01% (w/v) crystal violet (a) and then quantified by dissolving in 96% ethanol and measuring absorbance at 590 nm (b). Means and standard deviations of three replicates are shown. Student's *t*-test was performed for testing differences between groups. * $P < 0.05$.

When we mixed *cfp*-tagged PAO1 and *yfp*-tagged PAO1/*p_{lac}-yhjH* strains 1:1 (vol:vol) for planktonic co-cultivation experiments, the wild-type PAO1 strain gained a higher level of relative fitness than the *c*-di-GMP depleted PAO1/*p_{lac}-yhjH* strain with and without exposure to TeO_3^{2-} (Fig. 6a). Moreover, PAO1/*p_{lac}-yhjH* was fully outcompeted by PAO1 in biofilm co-cultures supplemented with TeO_3^{2-} (Fig. 6b). These results suggest that variants with low intracellular *c*-di-GMP content are unlikely to be protected and maintained by both *P. aeruginosa* planktonic and biofilm communities when *c*-di-GMP is required for stress response.

Discussion

Bacterial cells face various types of stress during the colonization of natural environments and hosts. A series of stress response mechanisms has evolved in bacteria to cope with these harmful conditions. One well characterized stringent stress response mechanism is SpoT-mediated ppGpp accumulation, which can be provoked by nutritional stress caused by harmful conditions such as antibiotic treatment and UV irradiation³⁹. ppGpp is able to bind directly to the bacterial RNA polymerase and further regulate transcriptional activity of many genes.

In addition to the stringent stress response, bacteria employ a wide range of social behaviors for surviving under unfavorable environmental conditions and these responses also contribute to bacterial pathogenesis⁴⁰. For example, the *Staphylococcus aureus* agr quorum-sensing system is involved in the oxidative stress response⁴¹. Biofilm formation is evoked as a stress response mechanism by a wide range of bacteria⁴². It involves encasing bacterial cells inside the densely packed EPS matrix components and attaching firmly to biotic and abiotic surfaces. Biofilms are up to 1,000 times more resistant to antimicrobial agents compared to their planktonic counterparts⁴³.

Recently, bacteria were found to form floating biofilm-resembling aggregates that are resistant to antimicrobials and phagocytosis²⁸. Our work here showed that TeO_3^{2-} exposure can elevate the *c*-di-GMP level in *P. aeruginosa* and lead to the formation of floating aggregates. TeO_3^{2-} -induced floating aggregate formation requires Pel and Psl polysaccharides as well as extracellular DNA (eDNA) (Fig. S1), in accordance with the Psl polysaccharide-eDNA interaction enabling the formation of skeleton of *P. aeruginosa*




Figure 4. Growth curve (a) and TeO_3^{2-} tolerance assay (b). *P. aeruginosa* PAO1, $\Delta sadC$, $\Delta siaD$, $\Delta sadC\Delta siaD$, and PAO1/ $p_{lac-yhjH}$ strains were cultivated in ABTGC medium at 37 °C with shaking for growth measurement. For TeO_3^{2-} tolerance assay, *P. aeruginosa* PAO1, $\Delta sadC$, $\Delta siaD$, $\Delta sadC\Delta siaD$, and PAO1/ $p_{lac-yhjH}$ strains were cultivated in ABTGC medium with the presence of 20 $\mu g/ml$ TeO_3^{2-} overnight followed by CFU determination. Means and standard deviations of three replicates are shown.

biofilms⁴⁴. In addition to serving as matrix scaffolds, the polysaccharides could also induce synthesis of iron siderophore pyoverdine via the Gac/Rsm pathway in the floating aggregates, as we had previously demonstrated⁴⁵. The formation of stress-induced biofilm-resembling aggregates might contribute to the dissemination of infection in the host.

The results presented here demonstrate that *P. aeruginosa* mutants with low c-di-GMP content were more sensitive to TeO_3^{2-} exposure in planktonic cultures and thus their growth was negatively affected by TeO_3^{2-} exposure, as compared to c-di-GMP containing wild-type strain (Fig. 4). Consistent with this finding, a recent study on biodegradation of 3-chloroaniline by *Comamonas testosteroni* reported that, compared with the wild type, the strain with an elevated c-di-GMP level exhibited a better growth on the toxic substrate at high concentrations⁴⁶. In addition to TeO_3^{2-} , the detergent Na-dodecylsulfate (SDS)³⁵ also raised the c-di-GMP levels and caused aggregation of *P. aeruginosa*. In accordance with the TeO_3^{2-} findings, the $\Delta siaD$ mutant with low intracellular c-di-GMP content was more sensitive to SDS during planktonic growth³⁵. Together, these studies highlight that c-di-GMP signaling is involved in multiple stress response mechanisms, which might due to multiple DGCs and PDEs being encoded by many bacterial species.

Finally, we found that wild-type PAO1 strain biofilms prevented the attachment of mutants with low intracellular c-di-GMP content in both normal and TeO_3^{2-} stress co-cultures. Our previous study revealed that the polysaccharides in *P. aeruginosa* biofilms could not be shared, for structural or functional benefits, by mutants that are defective in their synthesis³⁸. These latter findings corroborate with the results presented here, and c-di-GMP mediated synthesis of polysaccharides may form another strategy to repress the proliferation and maintenance of c-di-GMP defective variants in biofilms. Considering




Figure 5. Relative fitness of $\Delta sadC\Delta siaD$ mutant to PAO1 in planktonic co-cultures and biofilm co-cultures in ABTGC medium with and without the presence of $10\ \mu\text{g/ml}$ TeO_3^{2-} (a). Means and standard deviations of three replicates are shown. Student's *t*-test was performed for testing differences between groups. * $P < 0.05$. CLSM images of biofilm co-cultures formed by *cfp*-tagged *P. aeruginosa* PAO1 and *yfp*-tagged $\Delta sadC\Delta siaD$ mutant in ABTGC medium with and without the presence of $10\ \mu\text{g/ml}$ TeO_3^{2-} (b). Representative image from triplicate experiments was shown for each condition. Bars, $50\ \mu\text{m}$.

that polysaccharides with similar structure to the *P. aeruginosa* polysaccharides are widely distributed in natural bacterial species, our results might reflect a conserved strategy employed by a range of bacterial species to repress the spreading of variants which cannot respond to environmental conditions by moderating their own c-di-GMP levels.

Methods

Bacterial strains and growth medium. The bacterial strains, plasmids, and primers used in this study are listed in Table 1. *Escherichia coli* DH5a strain was used for standard DNA manipulations. LB medium⁴⁷ was used to cultivate *E. coli* strains. Batch cultivation of *P. aeruginosa* was carried out at 37°C in ABT minimal medium⁷ supplemented with $5\ \text{g glucose l}^{-1}$ (ABTG) or $2\ \text{g glucose l}^{-1}$ and $2\ \text{g casamino acids l}^{-1}$ (ABTGC). For plasmid maintenance in *E. coli*, the medium was supplemented with $100\ \mu\text{g ampicillin (Ap) ml}^{-1}$, $15\ \mu\text{g gentamicin (Gm) ml}^{-1}$, $15\ \mu\text{g tetracycline (Tc) ml}^{-1}$, or $8\ \mu\text{g chloramphenicol (Cm) ml}^{-1}$. For marker selection in *P. aeruginosa*, $30\ \mu\text{g Gm ml}^{-1}$, $50\ \mu\text{g Tc ml}^{-1}$, and $200\ \mu\text{g carbenicillin (Cb) ml}^{-1}$ were used, when appropriate. Antibiotics were not added to *P. aeruginosa* cultures for c-di-GMP,




Figure 6. Relative fitness of PAO1/p_{lac}-yhjH mutant to PAO1 in planktonic co-cultures and biofilm co-cultures in ABTGC medium with and without the presence of 10 µg/ml TeO₃²⁻ (a). Means and standard deviations of three replicates are shown. Student's *t*-test was performed for testing differences between groups. * *P* < 0.05. CLSM images of biofilm co-cultures formed by cfp-tagged *P. aeruginosa* PAO1 and yfp-tagged PAO1/p_{lac}-yhjH mutant in ABTGC medium with and without the presence of 10 µg/ml TeO₃²⁻ (b). Representative image from triplicate experiments was shown for each condition. Bars, 50 µm.

stress response and biofilm assays as the plasmids we used were highly stable for these short-term experiments.

Construction of *P. aeruginosa* mutants. The $\Delta pelA$, $\Delta pslBCD$ and $\Delta pelA\Delta pslBCD$ mutants defective for Pel and/or Psl polysaccharide biogenesis were constructed by allelic displacement as previously described⁴⁸. The $\Delta sadC$, $\Delta siaD$ and $\Delta sadC\Delta siaD$ mutants defective for SadC and/or SiaD diguanylate cyclase were constructed by allelic displacement as previously described³⁴.

Quantification of static biofilms. The microtitre tray biofilm formation assay was performed as described by O'Toole & Kolter⁴⁹. Briefly, overnight cultures grown in ABTG medium were diluted to OD₆₀₀ = ~0.001 with fresh ABTG medium and transferred to the wells of polystyrene 96-well microtitre trays (200 µl per well) and incubated for 24 h at 37 °C. Liquid culture was removed from each well and the wells were washed twice with 0.9% NaCl followed by staining with 0.1% crystal violet and washing twice with 0.9% NaCl. The crystal violet-stained biofilms were then resuspended in 96% ethanol, and the absorbance of biofilm-associated dye was measured at 600 nm. Experiments were performed in triplicate, and the results are shown as the mean ± sd.

Strain(s) or plasmid	Relevant characteristic(s)	Source or reference
<i>P. aeruginosa</i> strains		
PAO1	Prototypic wild-type strain	55
$\Delta pelA$	Gm ^r ; <i>pelA</i> derivative of PAO1 constructed by allelic exchange	38
$\Delta pslBCD$	Gm ^r ; <i>pslBCD</i> derivative of PAO1 constructed by allelic exchange	38
$\Delta pelA\Delta pslBCD$	Gm ^r ; <i>pelA/pslBCD</i> derivative of PAO1 constructed by allelic exchange	38
$\Delta sadC$	Gm ^r ; <i>sadC</i> derivative of PAO1 constructed by allelic exchange	This study
$\Delta siaD$	Gm ^r ; <i>siaD</i> derivative of PAO1 constructed by allelic exchange	This study
$\Delta sadC\Delta siaD$	Gm ^r ; <i>sadC/siaD</i> derivative of PAO1 constructed by allelic exchange	This study
PAO1/ <i>p_{cdrA}-gfp</i>	Gm ^r ; PAO1 carrying the <i>p_{cdrA}-gfp</i> report	30
$\Delta sadC$ / <i>p_{cdrA}-gfp</i>	Gm ^r ; $\Delta sadC$ carrying the <i>p_{cdrA}-gfp</i> report	This study
$\Delta siaD$ / <i>p_{cdrA}-gfp</i>	Gm ^r ; $\Delta siaD$ carrying the <i>p_{cdrA}-gfp</i> report	This study
$\Delta sadC\Delta siaD$ / <i>p_{cdrA}-gfp</i>	Gm ^r ; $\Delta sadC\Delta siaD$ carrying the <i>p_{cdrA}-gfp</i> report	This study
PAO1/ <i>p_{lac}-yhjH</i>	Tc ^r ; PAO1 containing the <i>p_{lac}-yhjH</i> vector	12
PAO1/ <i>p_{pel}-lacZ</i>	Tc ^r ; PAO1 carrying the mini-CTX- <i>pelA-lacZ</i> report	This study
$\Delta sadC$ / <i>p_{pel}-lacZ</i>	Tc ^r ; $\Delta sadC$ carrying the mini-CTX- <i>pelA-lacZ</i> report	This study
$\Delta siaD$ / <i>p_{pel}-lacZ</i>	Tc ^r ; $\Delta siaD$ carrying the mini-CTX- <i>pelA-lacZ</i> report	This study
$\Delta sadC\Delta siaD$ / <i>p_{pel}-lacZ</i>	Tc ^r ; $\Delta sadC\Delta siaD$ carrying mini-CTX- <i>pelA-lacZ</i> report	This study
<i>E. coli</i> strain		
DH5 α	F ⁻ , $\phi 80dlacZ\Delta M15$, $\Delta(lacZYA-argF)U169$, <i>deoR</i> , <i>recA1</i> , <i>endA1</i> , <i>hsdR17</i> (rK ⁻ , mK ⁺), <i>phoA</i> , <i>supE44</i> , λ^- , <i>thi-1</i> , <i>gyrA96</i> , <i>relA1</i>	Laboratory collection
Plasmids		
pUCP22	Ap ^r ; Gm ^r ; Broad-host-range cloning vector	56
pMPELA	Ap ^r ; Gm ^r ; <i>pelA</i> allelic replacement vector	57
pMPSL-KO1	Ap ^r ; Gm ^r ; <i>pslBCD</i> allelic replacement vector	58
pEX18Gm:: $\Delta sadC$	Gm ^r ; <i>sadC</i> allelic replacement vector	34
pEX18Gm:: $\Delta siaD$	Gm ^r ; <i>siaD</i> allelic replacement vector	34
pFLP2	Ap ^r ; Source of FLP recombinase	59
<i>p_{cdrA}-gfp</i>	Ap ^r ; Gm ^r ; pUCP22 carrying the <i>p_{cdrA}-gfp</i> fusion	30
pRK600	Cm ^r ; <i>ori</i> ColE1 RK2-Mob ⁺ RK2-Tra ⁺ ; helper vector for conjugation	60
<i>p_{lac}-yhjH</i>	Tc ^r ; pBBR1MCS3 carrying the <i>yhjH</i> gene	12
Mini-CTX- <i>pel-lacZ</i>	Tc ^r ; mini-CTX vector carrying the <i>pel-lacZ</i> fusion	36

Table 1. Strains and plasmids used in this study.

Field emission scanning electron microscopy (FE-SEM) and energy-dispersive X-ray spectroscopy (EDX). The aggregates were dried and coated with platinum (Pt) using a vacuum electric sputter coater (JEOL JFC-1300, JEOL Asia Pte Ltd, Singapore). SEM images were taken using a field emission scanning electron microscope (FE-SEM, JSM-7600, JEOL Asia Pte Ltd, Singapore) at a voltage of 2.0–5.0 kV and EDX spectra were obtained using an energy-dispersive X-ray spectroscope (AZtecEnergy, Oxford Instruments, Oxfordshire, UK) as previously described⁵⁰. Experiments were performed in triplicate, and representative images were shown.

Reactive oxygen species (ROS) assay. PAO1 cultures were grown in ABTGC or LB medium controls and media with $10\ \mu\text{g ml}^{-1}$ TeO_3^{2-} , SeO_3^{2-} and SeO_4^{2-} , respectively. The ROS content of 1 ml stationary phase bacterial cells were then measured by using the OxiSelect™ *in vitro* ROS/RNS assay kit (Green Fluorescence), accordingly to manufacturer's instructions. 2', 7'-dichlorodihydrofluorescein (DCF) was used as a standard and the concentrations of ROS from PAO1 cultures were estimated according to the DCF standard curve. The fluorescence of the samples was read by the Tecan Infinite 2000 Microplate Reader at 480 nm excitation/530 nm emission. Experiments were performed in triplicate, and the results are shown as the mean \pm sd. Student's *t*-test was performed for testing differences between groups.

iTRAQ-based proteomics analyses. *P. aeruginosa* PAO1 was grown in ABTG medium with and without $10\ \mu\text{g/ml}$ TeO_3^{2-} at 37°C with shaking until stationary phase was reached. Cells were harvested and iTRAQ-based proteomics analyses were carried out as previously described¹².

Determination of minimal inhibitory concentration (MIC). The MIC assays employed a microtiter broth dilution method as previously described in the NCSLA guidelines⁵¹. Briefly, fresh ~16 h cultures of *P. aeruginosa* were diluted in ABTG medium. For determination of MIC, potassium tellurite was dissolved in water at a concentration 10 times higher than the required range by serial dilutions from a stock solution. $10\ \mu\text{l}$ of each concentration were added to each corresponding well of a 96-well microtiter plate (polypropylene, Costar Corp.) and $90\ \mu\text{l}$ of bacterial culture ($\sim 1 \times 10^5$ cells) in ABTG medium were added. The plate was incubated at 37°C for 16–18 h. MIC was taken as the lowest concentration where no visual growth (based on OD_{600}) of bacteria was detected. Experiments were performed in triplicate and representative results were shown.

TeO_3^{2-} tolerance assay. Overnight cultures of different *P. aeruginosa* strains were inoculated into ABTGC medium containing $20\ \mu\text{g/ml}$ TeO_3^{2-} and cultivated overnight (24 h). Overnight cultures were serially diluted and plated onto LB agar media. LB plates were incubated at 37°C overnight before CFU calculation. Experiments were performed in triplicate, and the results are shown as the mean \pm sd.

Beta-galactosidase activity assay. A classical β -galactosidase assay⁵² was used to measure expression of the P_{pel} -*lacZ* fusion in *P. aeruginosa* strains transformed with the mini-CTX-*pel*-*lacZ* fusion³⁶, which carries the *pel* promoter fused to the *E. coli lacZ* gene. Experiments were performed in triplicate, and the results are shown as the mean \pm sd. Student's *t*-test was performed for testing differences between groups.

Gfp reporter fusion assay. The expression of the c-di-GMP P_{cdrA} -*gfp* biosensor³⁰ in *P. aeruginosa* strains in the presence and absence of TeO_3^{2-} was monitored by using a Tecan Infinite 2000 Microplate Reader. Monitoring strains were cultivated in 24-well microtiter plate with ABTGC medium with different concentrations of TeO_3^{2-} at 37°C with shaking. OD_{600} and GFP fluorescence (in relative fluorescence units, rfu) were measured every hour until the culture reach stationary growth phase. Experiments were performed in triplicate, and the results are shown as the mean \pm sd. Student's *t*-test was performed for testing differences between groups.

Quantification of c-di-GMP concentration. Extraction of c-di-GMP was conducted as previously described⁴⁵. $10\ \text{ml}$ of *P. aeruginosa* cells in the early stationary phase from the ABTGC medium with and without $10\ \mu\text{g/ml}$ TeO_3^{2-} were washed twice with $1\ \text{mM}$ ammonium acetate. Cells were lysed with $0.6\ \text{M}$ HClO_4 on ice for 30 min. Cell debris was removed by centrifugation and supernatant was neutralized to pH 6.0 with the addition of $2.5\ \text{M}$ KHCO_3 . The precipitated KClO_4 was removed by centrifugation and the supernatant was used for relative quantification of c-di-GMP. The concentration was measured by High Performance Liquid Chromatography (HPLC), the injection volume is $20\ \mu\text{l}$ with 254 nm as detection wavelength. Reverse-phase C18 Targa column ($2.1 \times 40\ \text{mm}$, $5\ \mu\text{m}$) (catalog number: TR-0421-C185) was used with solvent A ($10\ \text{mM}$ ammonium acetate in water) and solvent B ($10\ \text{mM}$ ammonium acetate in methanol) at a flow rate of $0.2\ \text{ml min}^{-1}$. Eluent gradient is as follows: 0 to 8 min, 1% B; 8 to 14 min, 15% B; 14 to 16 min, 19% B; 16 to 24 min, 100% B; 24 to 32 min, 100% B; 32 to 40 min, 1% B; 40 to 42 min, 1% B. The retention time of c-di-GMP is around 14.0 min. The c-di-GMP concentration was normalized by total protein concentration. The relative c-di-GMP concentrations of cells treated with $10\ \mu\text{g ml}^{-1}$ tellurite against cells in ABTGC only were shown. Experiments were performed in triplicate, and the results are shown as the mean \pm sd. Student's *t*-test was performed for testing differences between groups.

Competition assay. Competition assays were performed in both planktonic and biofilm co-cultures. In planktonic co-cultures, *cfp*-tagged wild-type PAO1 was mixed 1:1 (vol/vol) with *yfp*-tagged PAO1/ P_{lac} -*yhjH* (or *yfp*-tagged $\Delta\text{sadC}\Delta\text{siaD}$) and the mixtures inoculated into fresh ABTGC medium with and without the presence of $10\ \mu\text{g/ml}$ TeO_3^{2-} . For relative fitness calculation, co-cultures were plated in LB agar plates after 24 h cultivation at 37°C with shaking. Colony-forming units (CFUs) N_t were determined from three individual experiments and the number of PAO1 and PAO1/ P_{lac} -*yhjH* (or $\Delta\text{sadC}\Delta\text{siaD}$) colonies were determined based on their specific fluorescence at times $t=0$ and at $t=T$. Relative fitness was

determined as $r_{ij} = [N_i(T) - N_i(0)] / [N_j(T) - N_j(0)]$ as previously described with modification⁵³, resulting in a fitness of '1' when competing organisms are equally fit. Experiments were performed in triplicate, and the results are shown as the mean \pm sd. Student's *t*-test was performed for testing differences between groups.

In biofilm co-cultures, *cfp*-tagged wild-type PAO1 cells were mixed with *yfp*-tagged PAO1/*p_{lac}-yjhH* (or *yfp*-tagged Δ *sadC* Δ *siaD*) cells at 1:1 (vol/vol) and the mixtures were inoculated into fresh ABTGC medium with and without the presence of 10 μ g/ml TeO_3^{2-} . Static biofilms were cultivated on cover slides at 37 °C for 24 h as previously described⁵⁴. Biofilms were imaged with a Zeiss LSM780 confocal laser scanning microscope (CLSM) equipped with detectors and filter sets for monitoring of Cfp and Yfp fluorescence. Images were obtained using a 40 \times /1.4 objective. Simulated three-dimensional images and sections as well as biovolumes were generated using the Imaris software package (Bitplane AG)⁸. The biovolume V_i of each strain in the biofilm mode was determined from three individual experiments based on their fluorescence at times $t = 0$ and at $t = T$. Relative fitness was determined as $r_{ij} = [V_i(T) - V_i(0)] / [V_j(T) - V_j(0)]$ as previously described with modification⁵³, resulting in a fitness of '1' when competing organisms are equally fit. Experiments were performed in triplicate, and the results are shown as the mean \pm sd. Student's *t*-test was performed for testing differences between groups.

References

- Schwartz, T., Hoffmann, S. & Obst, U. Formation of natural biofilms during chlorine dioxide and u.v. disinfection in a public drinking water distribution system. *J Appl Microbiol.* **95**, 591–601 (2003).
- Baker-Austin, C., Wright, M. S., Stepanauskas, R. & McArthur, J. V. Co-selection of antibiotic and metal resistance. *Trends Microbiol.* **14**, 176–182 (2006).
- Flemming, H. C., Neu, T. R. & Wozniak, D. J. The EPS matrix: the “house of biofilm cells”. *J Bacteriol* **189**, 7945–7947 (2007).
- Stewart, P. S. & Costerton, J. W. Antibiotic resistance of bacteria in biofilms. *Lancet* **358**, 135–138 (2001).
- Yang, L. *et al.* Combating biofilms. *FEMS Immunol Med Microbiol* **65**, 146–157 (2012).
- Herrmann, G. *et al.* Colistin-tobramycin combinations are superior to monotherapy concerning the killing of biofilm *Pseudomonas aeruginosa*. *J Infect Dis.* **202**, 1585–1592, doi:10.1086/656788 (2010).
- Yang, L. *et al.* Effects of iron on DNA release and biofilm development by *Pseudomonas aeruginosa*. *Microbiology* **153**, 1318–1328 (2007).
- Chua, S. L. *et al.* Dispersed cells represent a distinct stage in the transition from bacterial biofilm to planktonic lifestyles. *Nature communications* **5**, 4462 (2014).
- Hengge, R. Principles of c-di-GMP signalling in bacteria. *Nat Rev Microbiol.* **7**, 263–273 (2009).
- Lory, S., Merighi, M. & Hyodo, M. Multiple activities of c-di-GMP in *Pseudomonas aeruginosa*. *Nucleic Acids Symp Ser (Oxf)* **53**, 51–52 (2009).
- Yang, L. *et al.* Pattern differentiation in co-culture biofilms formed by *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *FEMS Immunol Med Microbiol.* **62**, 339–347 (2011).
- Chua, S. L. *et al.* Bis-(3'-5')-cyclic dimeric GMP regulates antimicrobial peptide resistance in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* **57**, 2066–2075 (2013).
- Gjermansen, M., Nilsson, M., Yang, L. & Tolker-Nielsen, T. Characterization of starvation-induced dispersion in *Pseudomonas putida* biofilms: genetic elements and molecular mechanisms. *Mol Microbiol.* **75**, 815–826 (2010).
- Kaplan, J. B. Antibiotic-induced biofilm formation. *Int J Artif Organs* **34**, 737–751 (2011).
- Klebansberger, J., Lautenschlager, K., Bressler, D., Wingender, J. & Philipp, B. Detergent-induced cell aggregation in subpopulations of *Pseudomonas aeruginosa* as a preadaptive survival strategy. *Environ Microbiol.* **9**, 2247–2259 (2007).
- Klonowska, A., Heulin, T. & Vermeglio, A. Selenite and tellurite reduction by *Shewanella oneidensis*. *Appl Environ Microbiol.* **71**, 5607–5609 (2005).
- Yurkov, V., Jappé, J. & Verméglio, A. Tellurite resistance and reduction by obligately aerobic photosynthetic bacteria. *Appl Environ Microbiol.* **62**, 4195–4198 (1996).
- Moscoso, H., Saavedra, C., Loyola, C., Pichuantes, S. & Vasquez, C. Biochemical characterization of tellurite-reducing activities of *Bacillus stearothermophilus* V. *Res Microbiol.* **149**, 389–397 (1998).
- Trutko, S. M. *et al.* Involvement of the respiratory chain of gram-negative bacteria in the reduction of tellurite. *Arch Microbiol.* **173**, 178–186 (2000).
- Zannoni, D. Bacterial processing of metalloids as a tool in Biotechnology. *J Biotechnol.* **150**, S52–S53 (2010).
- Mohanty, A., Liu, Y., Yang, L. & Cao, B. Extracellular biogenic nanomaterials inhibit pyoverdine production in *Pseudomonas aeruginosa*: A novel insight into impacts of metal(loid)s on environmental bacteria. *Applied Microbiology and Biotechnology.* **99**, 1957–1966 (2015).
- Fleming, A. & Young, M. Y. The inhibitory action of potassium tellurite on coliform bacteria. *J Pathol Bacteriol* **51**, 29–35 (1940).
- Cummins, J., Reen, F. J., Baysse, C., Mooij, M. J. & O'Gara, F. Subinhibitory concentrations of the cationic antimicrobial peptide colistin induce the *pseudomonas* quinolone signal in *Pseudomonas aeruginosa*. *Microbiology* **155**, 2826–2837 (2009).
- Brazas, M. D. & Hancock, R. E. Ciprofloxacin induction of a susceptibility determinant in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* **49**, 3222–3227 (2005).
- Thompson, D. K. *et al.* Proteomics reveals a core molecular response of *Pseudomonas putida* F1 to acute chromate challenge. *BMC Genomics* **11**, 311 (2010).
- Perez, J. M. *et al.* Bacterial toxicity of potassium tellurite: unveiling an ancient enigma. *PLoS One* **2**, e211 (2007).
- Bebien, M., Chauvin, J. P., Adriano, J. M., Grosse, S. & Vermeglio, A. Effect of selenite on growth and protein synthesis in the phototrophic bacterium *Rhodospira rubra*. *Appl Environ Microbiol.* **67**, 4440–4447 (2001).
- Alhede, M. *et al.* Phenotypes of non-attached *Pseudomonas aeruginosa* aggregates resemble surface attached biofilm. *PLoS One* **6**, e27943 (2011).
- Borlee, B. R. *et al.* *Pseudomonas aeruginosa* uses a cyclic-di-GMP-regulated adhesin to reinforce the biofilm extracellular matrix. *Mol Microbiol.* **75**, 827–842 (2010).
- Rybtke, M. T. *et al.* Fluorescence-Based Reporter for Gauging Cyclic Di-GMP Levels in *Pseudomonas aeruginosa*. *Appl Environ Microbiol.* **78**, 5060–5069 (2012).
- Jacob-Dubuisson, F., Loch, C. & Antoine, R. Two-partner secretion in Gram-negative bacteria: a thrifty, specific pathway for large virulence proteins. *Mol Microbiol.* **40**, 306–313 (2001).
- Gottig, N., Garavaglia, B. S., Garofalo, C. G., Orellano, E. G. & Ottado, J. A filamentous hemagglutinin-like protein of *Xanthomonas axonopodis* pv. citri, the phytopathogen responsible for citrus canker, is involved in bacterial virulence. *PLoS One* **4**, e4358 (2009).

33. Serra, D. O. *et al.* FHA-mediated cell-substrate and cell-cell adhesions are critical for *Bordetella pertussis* biofilm formation on abiotic surfaces and in the mouse nose and the trachea. *PLoS One* **6**, e28811 (2011).
34. Irie, Y. *et al.* Self-produced exopolysaccharide is a signal that stimulates biofilm formation in *Pseudomonas aeruginosa*. *P Natl Acad Sci USA* **109**, 20632–20636 (2012).
35. Klebensberger, J., Birkenmaier, A., Geffers, R., Kjelleberg, S. & Philipp, B. SiaA and SiaD are essential for inducing autoaggregation as a specific response to detergent stress in *Pseudomonas aeruginosa*. *Environ Microbiol.* **11**, 3073–3086 (2009).
36. Sakuragi, Y. & Kolter, R. Quorum-sensing regulation of the biofilm matrix genes (pel) of *Pseudomonas aeruginosa*. *Journal of bacteriology* **189**, 5383–5386 (2007).
37. Chew, S. C. *et al.* Dynamic remodeling of microbial biofilms by functionally distinct exopolysaccharides. *mBio.* **5**, e01536–01514 (2014).
38. Yang, L. *et al.* Distinct roles of extracellular polymeric substances in *Pseudomonas aeruginosa* biofilm development. *Environ Microbiol* **13**, 1705–1717 (2011).
39. Potrykus, K. & Cashel, M. (p)ppGpp: still magical? *Annu Rev Microbiol.* **62**, 35–51 (2008).
40. West, S. A., Diggle, S. P., Buckling, A., Gardner, A. & Griffins, A. S. The social lives of microbes. *Annu Rev Ecol Evol S* **38**, 53–77 (2007).
41. Sun, F. *et al.* Quorum-sensing agr mediates bacterial oxidation response via an intramolecular disulfide redox switch in the response regulator AgrA. *Proc Natl Acad Sci U S A* **109**, 9095–9100 (2012).
42. O'Toole, G. A. & Stewart, P. S. Biofilms strike back. *Nature biotechnology* **23**, 1378–1379 (2005).
43. Hoiby, N., Bjarnsholt, T., Givskov, M., Molin, S. & Ciofu, O. Antibiotic resistance of bacterial biofilms. *International journal of antimicrobial agents* **35**, 322–332 (2010).
44. Wang, S. *et al.* The exopolysaccharide Psl-eDNA interaction enables the formation of a biofilm skeleton in *Pseudomonas aeruginosa*. *Environ Microbiol Rep*, **7**, 330–340 (2015).
45. Chen, Y. *et al.* Multiple diguanylate cyclase-coordinated regulation of pyoverdine synthesis in *Pseudomonas aeruginosa*. *Environ Microbiol Rep*, In Press (2015).
46. Wu, Y., Ding, Y., Cohen, Y. & Cao, B. Elevated level of the second messenger c-di-GMP in *Comamonas testosteroni* enhances biofilm formation and biofilm-based biodegradation of 3-chloroaniline. *Appl Microbiol Biotechnol.* **99**, 1967–1976 (2015).
47. Bertani, G. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. *J Bacteriol* **62**, 293–300 (1951).
48. Starkey, M. *et al.* *Pseudomonas aeruginosa* rugose small-colony variants have adaptations that likely promote persistence in the cystic fibrosis lung. *J Bacteriol* **191**, 3492–3503 (2009).
49. O'Toole, G. A. & Kolter, R. Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development. *Mol Microbiol.* **30**, 295–304 (1998).
50. Ng, C. K. *et al.* Influence of outer membrane c-type cytochromes on particle size and activity of extracellular nanoparticles produced by *Shewanella oneidensis*. *Biotechnol Bioeng.* **110**, 1831–1837 (2013).
51. Wiegand, I., Hilpert, K. & Hancock, R. E. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat Protoc.* **3**, 163–175 (2008).
52. Smale, S. T. Beta-galactosidase assay. *Cold Spring Harbor protocols* **2010**, pdb prot5423 (2010).
53. Flohr, R. C., Blom, C. J., Rainey, P. B. & Beaumont, H. J. Founder niche constrains evolutionary adaptive radiation. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 20663–20668 (2013).
54. Liu, Y., Yang, L. & Molin, S. Synergistic activities of an efflux pump inhibitor and iron chelators against *Pseudomonas aeruginosa* growth and biofilm formation. *Antimicrob Agents Chemother* **54**, 3960–3963 (2010).
55. Holloway, B. W. & Morgan, A. F. Genome organization in *Pseudomonas*. *Annu Rev Microbiol.* **40**, 79–105 (1986).
56. West, S. E., Schweizer, H. P., Dall, C., Sample, A. K. & Runyen-Janecky, L. J. Construction of improved *Escherichia-Pseudomonas* shuttle vectors derived from pUC18/19 and sequence of the region required for their replication in *Pseudomonas aeruginosa*. *Gene* **148**, 81–86 (1994).
57. Starkey, M. *et al.* *Pseudomonas aeruginosa* Rugose Small-Colony Variants Have Adaptations That Likely Promote Persistence in the Cystic Fibrosis Lung. *J Bacteriol* **191**, 3492–3503 (2009).
58. Kirisits, M. J., Prost, L., Starkey, M. & Parsek, M. R. Characterization of colony morphology variants isolated from *Pseudomonas aeruginosa* biofilms. *Appl Environ Microbiol.* **71**, 4809–4821 (2005).
59. Hoang, T. T., Karkhoff-Schweizer, R. R., Kutchma, A. J. & Schweizer, H. P. A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* **212**, 77–86 (1998).
60. Kessler, B., de Lorenzo, V. & Timmis, K. N. A general system to integrate lacZ fusions into the chromosomes of gram-negative eubacteria: regulation of the Pm promoter of the TOL plasmid studied with all controlling elements in monocopy. *Mol Gen Genet* **233**, 293–301 (1992).

Acknowledgments

We thank Dr. Matthew R. Parsek (University of Washington, Seattle, USA) for providing the pEX18Gm:: $\Delta sadC$ and pEX18Gm:: $\Delta siaD$ plasmids for constructing the $\Delta sadC$ and $\Delta siaD$ mutants. This research was supported by the National Research Foundation and Ministry of Education Singapore under its Research Centre of Excellence Programme and the Start-up Grants (M4080847.030) and (M4330002.C70) from Nanyang Technological University, Singapore. A portion of this work was supported by grants from the Danish Council for Strategic Research (M.G.) and AcRF Tier 2 (MOE2014-T2-2-172) from Ministry of Education, Singapore.

Author Contributions

T.T.N., B.C., S.K. and L.Y. designed the project. S.L.C., M.T.R., J.B.A., M.J.Y. and K.S. performed the experiments. T.E.N., M.G., B.C. and L.Y. interpreted data. B.C. and L.Y. wrote the main manuscript text. All authors reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Chua, S. L. *et al.* C-di-GMP regulates *Pseudomonas aeruginosa* stress response to tellurite during both planktonic and biofilm modes of growth. *Sci. Rep.* **5**, 10052; doi: 10.1038/srep10052 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>