



## Genome sequence of *Kocuria varians* G6 isolated from a slaughterhouse in Denmark

Raghupathi, Prem Krishnan; Herschend, Jakob; Røder, Henriette Lyng; Sørensen, Søren Johannes; Burmølle, Mette

*Published in:*  
Genome Announcements

*DOI:*  
[10.1128/genomeA.00076-16](https://doi.org/10.1128/genomeA.00076-16)

*Publication date:*  
2016

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Raghupathi, P. K., Herschend, J., Røder, H. L., Sørensen, S. J., & Burmølle, M. (2016). Genome sequence of *Kocuria varians* G6 isolated from a slaughterhouse in Denmark. *Genome Announcements*, 4(2), [e00076-16]. <https://doi.org/10.1128/genomeA.00076-16>

# Genome Sequence of *Kocuria varians* G6 Isolated from a Slaughterhouse in Denmark

Prem K. Raghupathi, Jakob Herschend, Henriette L. Røder, Søren J. Sørensen, Mette Burmølle

Section for Microbiology, Department of Biology, University of Copenhagen, Copenhagen, Denmark

We report here the first draft genome sequence of *Kocuria varians* G6, which was isolated from a meat chopper at a small slaughterhouse in Denmark. The 2.90-Mb genome sequence consists of 95 contigs and contains 2,518 predicted protein-coding genes.

Received 16 January 2016 Accepted 12 February 2016 Published 31 March 2016

**Citation** Raghupathi PK, Herschend J, Røder HL, Sørensen SJ, Burmølle M. 2016. Genome sequence of *Kocuria varians* G6 isolated from a slaughterhouse in Denmark. *Genome Announc* 4(2):e00076-16. doi:10.1128/genomeA.00076-16.

**Copyright** © 2016 Raghupathi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mette Burmølle, burmolle@bio.ku.dk.

*Kocuria varians* is a Gram-positive bacterium belonging to the family *Micrococcineae*. *Kocuria* species are ubiquitous and are highly adapted to their ecological niches (1). *K. varians* is predominantly isolated from different food-processing plants, e.g., milk-processing (2), fermented meat (3), and beef-processing plants (4). In these industrial settings, *K. varians* is often found in biofilms, and it is reported to favor both attachment and detachment of pathogens, namely, *Listeria monocytogenes* (2, 5). *K. varians* was also reported to cause brain abscess (6). Further, the beneficial effects of *K. varians* include improvement of the flavor profile in fermented sausage (7) and the degradation of putrescine (3). From a review of above-mentioned literature, *K. varians* has a central role in various complex interactions. Currently, there is no genome sequence information available for *K. varians*.

In this report, we announce the first draft assembly genome of *K. varians* isolated from a slaughterhouse in Denmark (8). The draft genome consists of 95 contigs, with an average G+C content of 70.5%. The whole-genome sequencing libraries were prepared using the Nextera XT kit (Illumina, USA), according to the manufacturer's recommendations, and then sequenced as a part of the flow cell, as 2 × 250-base paired-end reads with Illumina MiSeq technology. The reads were cleaned and trimmed using CLC Genomics Workbench 7 (CLC bio, Denmark). The processed reads were assembled using SPAdes version 3.5.0 (9). The assembled genomes were uploaded to the RAST (10) server to perform functional annotations and to check and screen for noncoding rRNAs and tRNAs. rRNA genes were predicted by RNAmmer 1.2 (11).

The annotated results predicted 2,518 coding sequences (1,189 coding sequences [CDSs] have functional predictions), 50 RNA genes, 15 tRNA-coding genes, and 3 rRNA-coding genes. The genome has single predicted copies of 5S-16S-23S rRNA genes. The number of genes transcribed from the positive strand was 1,394, while 1,124 genes were transcribed from the negative strand. There are 364 predicted subsystems in the genome of G6, and we used this to reconstruct the metabolic network. The annotated genome has 22 genes involved in virulence, disease, and defense, including 10 genes coding for antibiotic resistance and 7 clustered

regularly interspaced short palindromic repeat (CRISPR) elements, indicating the influence of phage exposure on the adaptation of this strain. Sixty-eight genes are involved in the stress responses of the bacterium. Functional comparison on the RAST server revealed the closest neighbors of *K. varians* to be *Kocuria rhizophila* DC2201, followed by *K. rhizophila* P7-4, *Rothia dentocariosa* ATCC 17931, and *Arthrobacter aurescens* TC1. Further work with this genome and comparisons to other *Kocuria* species will give more insights into the adaptation and evolution of *K. varians* to different environments.

**Nucleotide sequence accession numbers.** The whole-genome shotgun project for *K. varians* G6 has been deposited at the European Nucleotide Archive (ENA) under the contig accession numbers CZJX01000001 to CZJX01000095. The version described in this paper is the first version.

## FUNDING INFORMATION

This work was funded partly by the Danish Council for Independent Research and the Villum Foundation.

## REFERENCES

- Kim WJ, Kim YO, Kim DS, Choi SH, Kim DW, Lee JS, Kong HJ, Nam BH, Kim BS, Lee SJ, Park HS, Chae SH. 2011. Draft genome sequence of *Kocuria rhizophila* 7-4. *J Bacteriol* 193:4286–4287.
- Carpentier B, Chassaing D. 2004. Interactions in biofilms between *Listeria monocytogenes* and resident microorganisms from food industry premises. *Int J Food Microbiol* 97:111–122. <http://dx.doi.org/10.1016/j.jfoodmicro.2004.03.031>.
- Callejón S, Sendra R, Ferrer S, Pardo I. 2015. Ability of *Kocuria varians* LTH 1540 to degrade putrescine: identification and characterization of a novel amine oxidase. *J Agric Food Chem* 63:41704178. <http://dx.doi.org/10.1021/jf5026967>.
- Leroy S, Giammarinaro P, Chacornac JP, Lebert I, Talon R. 2010. Biodiversity of indigenous staphylococci of naturally fermented dry sausages and manufacturing environments of small-scale processing units. *Food Microbiol* 27:294–301. <http://dx.doi.org/10.1016/j.fm.2009.11.005>.
- Midelet G, Kobilinsky A, Carpentier B. 2006. Construction and analysis of fractional multifactorial designs to study attachment strength and transfer of *Listeria monocytogenes* from pure or mixed biofilms after contact with a solid model food. *Appl Environ Microbiol* 72:2313–2321. <http://dx.doi.org/10.1128/AEM.72.4.2313-2321.2006>.

6. Tsai C-Y, Su SH, Cheng Y-H, Chou YL, Tsai T-H, Lieu A-S. 2010. *Kocuria varians* infection associated with brain abscess: a case report. *BMC Infect Dis* 10:102. <http://dx.doi.org/10.1186/1471-2334-10-102>.
7. Tremonte P, Succi M, Reale A, Di Renzo T, Sorrentino E, Coppola R. 2007. Interactions between strains of *Staphylococcus xylosus* and *Kocuria varians* isolated from fermented meats. *J Appl Microbiol* 103:743–751. <http://dx.doi.org/10.1111/j.1365-2672.2007.03315.x>.
8. Røder HL, Raghupathi PK, Herschend J, Brejnrod A, Knøchel S, Sørensen SJ, Burmølle M. 2015. Interspecies interactions result in enhanced biofilm formation by co-cultures of bacteria isolated from a food processing environment. *Food Microbiol* 51:18–24. <http://dx.doi.org/10.1016/j.fm.2015.04.008>.
9. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477.
10. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
11. Lagesen K, Hallin P, Rødland EA, Staerfeldt H-H, Rognes T, Ussery DW. 2007. RNAMmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.