

UNIVERSITY OF COPENHAGEN



## Managing Brain Extracellular K(+) during Neuronal Activity The Physiological Role of the Na(+)/K(+)-ATPase Subunit Isoforms

Larsen, Brian Roland; Stoica, Anca; MacAulay, Nanna

*Published in:*  
Frontiers in Physiology

*DOI:*  
[10.3389/fphys.2016.00141](https://doi.org/10.3389/fphys.2016.00141)

*Publication date:*  
2016

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[Unspecified](#)

*Citation for published version (APA):*  
Larsen, B. R., Stoica, A., & MacAulay, N. (2016). Managing Brain Extracellular K(+) during Neuronal Activity: The Physiological Role of the Na(+)/K(+)-ATPase Subunit Isoforms. *Frontiers in Physiology*, 7, [141].  
<https://doi.org/10.3389/fphys.2016.00141>



# Managing Brain Extracellular $K^+$ during Neuronal Activity: The Physiological Role of the $Na^+/K^+$ -ATPase Subunit Isoforms

Brian Roland Larsen, Anca Stoica and Nanna MacAulay\*

Department of Neuroscience and Pharmacology, University of Copenhagen, Copenhagen, Denmark

## OPEN ACCESS

### Edited by:

Sigrid A. Langhans,  
Al duPont Hospital for Children, USA

### Reviewed by:

Francisco F. De-Miguel,  
Universidad Nacional Autonoma de  
Mexico, Mexico  
Gustavo Blanco,  
Kansas University Medical Center,  
USA

### \*Correspondence:

Nanna MacAulay  
macaulay@sund.ku.dk

### Specialty section:

This article was submitted to  
Membrane Physiology and Membrane  
Biophysics,  
a section of the journal  
Frontiers in Physiology

**Received:** 14 January 2016

**Accepted:** 04 April 2016

**Published:** 22 April 2016

### Citation:

Larsen BR, Stoica A and MacAulay N  
(2016) Managing Brain Extracellular  
 $K^+$  during Neuronal Activity: The  
Physiological Role of the  
 $Na^+/K^+$ -ATPase Subunit Isoforms.  
*Front. Physiol.* 7:141.  
doi: 10.3389/fphys.2016.00141

During neuronal activity in the brain, extracellular  $K^+$  rises and is subsequently removed to prevent a widespread depolarization. One of the key players in regulating extracellular  $K^+$  is the  $Na^+/K^+$ -ATPase, although the relative involvement and physiological impact of the different subunit isoform compositions of the  $Na^+/K^+$ -ATPase remain unresolved. The various cell types in the brain serve a certain temporal contribution in the face of network activity; astrocytes respond directly to the immediate release of  $K^+$  from neurons, whereas the neurons themselves become the primary  $K^+$  absorbers as activity ends. The kinetic characteristics of the catalytic  $\alpha$  subunit isoforms of the  $Na^+/K^+$ -ATPase are, partly, determined by the accessory  $\beta$  subunit with which they combine. The isoform combinations expressed by astrocytes and neurons, respectively, appear to be in line with the kinetic characteristics required to fulfill their distinct physiological roles in clearance of  $K^+$  from the extracellular space in the face of neuronal activity. Understanding the nature, impact and effects of the various  $Na^+/K^+$ -ATPase isoform combinations in  $K^+$  management in the central nervous system might reveal insights into pathological conditions such as epilepsy, migraine, and spreading depolarization following cerebral ischemia. In addition, particular neurological diseases occur as a result of mutations in the  $\alpha 2$ - (familial hemiplegic migraine type 2) and  $\alpha 3$  isoforms (rapid-onset dystonia parkinsonism/alternating hemiplegia of childhood). This review addresses aspects of the  $Na^+/K^+$ -ATPase in the regulation of extracellular  $K^+$  in the central nervous system as well as the related pathophysiology. Understanding the physiological setting in non-pathological tissue would provide a better understanding of the pathological events occurring during disease.

**Keywords:**  $K^+$  clearance, astrocytes, glutamate, brain ion homeostasis, extracellular space


## $K^+$ MANAGEMENT IN THE EXTRACELLULAR SPACE OF THE CENTRAL NERVOUS SYSTEM

Neuronal activity, in the form of propagating action potentials, results in a transient release of  $K^+$  into the extracellular space (Frankenhaeuser and Hodgkin, 1956). Prolonged accumulation of  $K^+$  in the extracellular space of the central nervous system (CNS) causes wide-spread depolarization of neurons and glia which results in compromised synaptic transmission, neuronal firing, and

neurotransmitter re-uptake. Consequently, it follows that extracellular K<sup>+</sup> must be closely managed and therefore cleared from the extracellular space following neuronal excitation. Brain extracellular K<sup>+</sup>, [K<sup>+</sup>]<sub>o</sub>, has indeed proven to be tightly regulated under physiological conditions; from a basal level of ~3 mM, [K<sup>+</sup>]<sub>o</sub> rarely rises above ~12 mM. This concentration is denoted as the ceiling level and more intense or longer electrical stimulation of neuronal pathways generally cannot push [K<sup>+</sup>]<sub>o</sub> above this point (Krnjevic and Morris, 1972; Prince et al., 1973; Futamachi et al., 1974; Heinemann and Lux, 1977). However, typical physiologically occurring neuronal activity in the CNS is estimated to cause [K<sup>+</sup>]<sub>o</sub> transients of only 0.2–0.4 mM above the baseline (Syková et al., 1974; Singer and Lux, 1975).


During neuronal activity, in which K<sup>+</sup> is released to the extracellular space and the neuronal structures thus lose part of their intracellular K<sup>+</sup>, an elevation in [K<sup>+</sup>]<sub>o</sub> is observed along with a parallel rise in intracellular [K<sup>+</sup>] in the neighboring glia cells (Ballanyi et al., 1987; Grafe and Ballanyi, 1987). The activity-dependent glial K<sup>+</sup> accumulation points to this cellular compartment acting as a temporal K<sup>+</sup> sink during neuronal release of K<sup>+</sup> into the extracellular space (Ballanyi et al., 1987; Grafe and Ballanyi, 1987). Shortly following termination of neuronal activity, the glial [K<sup>+</sup>]<sub>i</sub> declines, indicative of steady glial K<sup>+</sup> efflux, while the neurons re-gain their [K<sup>+</sup>]<sub>i</sub> via re-uptake into the neuronal structures (Ballanyi et al., 1987; Grafe and Ballanyi, 1987, see **Figure 1**). Surely, part of the neuronally-released K<sup>+</sup> simply diffuses away in the extracellular space (Gardner-Medwin, 1983) but the molecular mechanism(s) that exercise the astrocytic clearance of extracellular K<sup>+</sup> have been investigated and debated intensely ever since the original findings in the mid-1900s:

- (I) Glial cells surrounding neurons respond directly to an increase in extracellular K<sup>+</sup> by a membrane depolarization (Kuffler and Nicholis, 1964; Kuffler and Potter, 1964; Orkand et al., 1966). The fact that the glial cells in essence behaved as K<sup>+</sup> electrodes, due to their high permeability for K<sup>+</sup>, indicated that they might act out a protective role toward neurons in connection to K<sup>+</sup> by redistribution of K<sup>+</sup> away from the site of activity (Orkand et al., 1966). This concept of K<sup>+</sup> channel-mediated spatial buffering of activity-evoked K<sup>+</sup> release relies on local differences in K<sup>+</sup> equilibrium- and membrane potentials leading to Kir4.1-mediated glial K<sup>+</sup> influx, electrotonic transfer of K<sup>+</sup> through the gap junction-coupled glia syncytium and Kir4.1-mediated release at a distant site (Karwoski et al., 1989; Walz, 2000; Kofuji and Newman, 2004). Although the spatial buffer currents are well-documented, the quantitative contribution to extracellular K<sup>+</sup> management of this channel-mediated mechanism may be limited and is under continuous debate, see (MacAulay and Zeuthen, 2012; Larsen and MacAulay, 2014; Larsen et al., 2014). This mode of K<sup>+</sup> clearance will not be discussed further here.
- (II) The Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup> cotransporter of the subtype 1 (NKCC1) is functionally expressed in cultured astrocytes in which the NKCC1 is responsible for around half of the overall cellular K<sup>+</sup> uptake, increasing its fractional



- uptake as the extracellular K<sup>+</sup> increases (Walz and Hertz, 1984; Larsen et al., 2014). Taken together with its low apparent K<sup>+</sup> affinity (~25 mM Larsen et al., 2014), which would allow it to increase its transport activity when faced with stimulus-evoked K<sup>+</sup> transients, NKCC1 has been suggested as a factor involved in clearance of K<sup>+</sup> from the extracellular space (Su et al., 2000; Kofuji and Newman, 2004; Hertz et al., 2015). In a recent study in rat hippocampal slices, we were nevertheless unable to demonstrate a participating role of NKCC1 in clearance of K<sup>+</sup> from the extracellular space of this particular brain region (Larsen et al., 2014). As negligible transcript and protein of NKCC1 is present in astrocytes *in vivo* (Plotkin et al., 1997; Clayton et al., 1998; Zhang et al., 2014) and NKCC1 is recognized to be upregulated in cultured cells of different origin (Raaf et al., 1996), NKCC1 appears to be active in cultured astrocytes but not involved in stimulus-evoked K<sup>+</sup> clearance in the rodent hippocampus under approached physiological conditions.
- (III) The Na<sup>+</sup>/K<sup>+</sup>-ATPase is comprised of an α and a β subunit (1:1) and drives uptake of 2 K<sup>+</sup> from the extracellular space in exchange for 3 Na<sup>+</sup> from the intracellular compartment. As this transport process translocates both ions against


their electrochemical gradients, the Na<sup>+</sup>/K<sup>+</sup>-ATPase relies on ATP hydrolysis as an integral step in its enzymatic cycle. Constant activity of the Na<sup>+</sup>/K<sup>+</sup>-ATPase is required to uphold the transmembrane ionic gradients, which create and maintain the membrane potential. The Na<sup>+</sup>/K<sup>+</sup>-ATPase is, in addition, involved in control of neuronal excitability (Haglund and Schwartzkroin, 1990; Wang et al., 2012; Gullledge et al., 2013) and has been indicated as a key contributor to management of extracellular [K<sup>+</sup>] following activity-induced K<sup>+</sup> release: Inhibition of the Na<sup>+</sup>/K<sup>+</sup>-ATPase yielded prolonged [K<sup>+</sup>]<sub>o</sub> recovery times upon electrical stimulation in the rat optic nerve (Ransom et al., 2000) and hippocampus (D'Ambrosio et al., 2002; Larsen et al., 2014, see **Figure 2**). In support of a dominating role of astrocytes in K<sup>+</sup> clearance during neuronal activity, astrocytes display a maximal Na<sup>+</sup>/K<sup>+</sup>-ATPase transport activity (V<sub>max</sub>) considerably larger than that of their neuronal counterpart (Kimmelberg et al., 1978; Walz and Hertz, 1982; White et al., 1992; Hajek et al., 1996). The astrocytic Na<sup>+</sup>/K<sup>+</sup>-ATPase isoform combination is, in addition, of a composition that allows astrocytes to increase their Na<sup>+</sup>/K<sup>+</sup>-ATPase activity as a function of elevated extracellular [K<sup>+</sup>] (Grisar et al., 1980; Walz and Hertz, 1982; Hajek et al., 1996; Larsen et al., 2014) and thus allows astrocytes to respond specifically to the K<sup>+</sup> transients observed in connection with neuronal activity. As different isoforms and/or subunit compositions of the Na<sup>+</sup>/K<sup>+</sup>-ATPase, each with distinct kinetic characteristics, are expressed in different cellular structures of the CNS, one may speculate whether these combinations are assigned specific roles in regard to management of activity-induced extracellular K<sup>+</sup> transients.



This review aims at presenting the CNS localization, role, and physiological impact of the different subunit isoform combinations of the Na<sup>+</sup>/K<sup>+</sup>-ATPase in the management of extracellular K<sup>+</sup> in the brain.

## LOCALIZATION OF THE Na<sup>+</sup>/K<sup>+</sup>-ATPase ISOFORMS IN NEURONS AND ASTROCYTES

Four isoforms of the α subunit of the Na<sup>+</sup>/K<sup>+</sup>-ATPase have been cloned (α1-α4) and three isoforms of the β subunit (β1-β3) (Blanco, 2005). Of these cloned isoforms, only α1-α3 and β1-β3 have been detected in the mammalian brain (McGrail et al., 1991; Cameron et al., 1994; Martín-Vasallo et al., 2000) although their quantity and cellular distribution remains unsettled and may depend on the brain region. The general picture that emerges reveals mRNA transcripts of α1 and β1 in both neurons and astrocytes while α2 and β2 are detected in astrocytes, and α3 exclusively in neuronal structures (Watts et al., 1991; Li et al., 2013; Zhang et al., 2014, see **Figure 3**). β3 transcript is prevalent in oligodendrocytes but negligible in astrocytes and neurons (Zhang et al., 2014). The total Na<sup>+</sup>/K<sup>+</sup>-ATPase α subunit mRNA increases about 10-fold from the fetal to adult stage with the α3 transcript reaching maximal and adult levels around post-natal day 7 which is only achieved for α1 and α2 around post-natal day 25 (Orlowski and Lingrel, 1988). The isoform expression pattern obtained at the mRNA level is more or less reflected at the protein level: α1 and β1 are detected in neurons and astrocytes, α2 and β2 predominantly in astrocytes and α3 in neurons (McGrail and Sweadner, 1986; McGrail et al., 1991; Cameron et al., 1994; Cholet et al., 2002; Richards et al., 2007). Alternative expression patterns have been reported which may be due to distinct regional differences in various brain structures and/or developmental stages, subpopulations




of neurons with altered isoform preference, altered expression in cultured neurons/astrocytes, and/or non-specific antibodies. Examples include detection of neuronal expression of the  $\alpha 2$  isoform in late gestation mouse embryos (Moseley et al., 2003) and in cultures and sections of rat hippocampus (McGrail et al., 1991; Cameron et al., 1994) as well as neuronal expression of  $\beta 2$  in cerebellar sections and co-cultures (Peng et al., 1997).

As opposed to the uniform plasma membrane distribution of  $\alpha 1$  in primary cultures of rat hippocampal astrocytes and neurons,  $\alpha 2$  and  $\alpha 3$  appeared to be organized in a reticular pattern paralleling the underlying endoplasmic reticulum (Juhászová and Blaustein, 1997). In addition,  $\alpha 2$  displayed co-localization with the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger in the plasma membrane of rat astrocytes (Juhászová and Blaustein, 1997) and localized to the glial leaflets surrounding dendritic spines (Cholet et al., 2002). The distinct cellular ( $\alpha 1$ ,  $\alpha 3$ ,  $\beta 1$  in neurons and  $\alpha 1$ ,  $\alpha 2$ ,  $\beta 1$ ,  $\beta 2$  in astrocytes) and subcellular localization pattern underscores the notion of the different Na<sup>+</sup>/K<sup>+</sup>-ATPase isoforms serving distinct physiological roles in the CNS.

## Na<sup>+</sup> AND K<sup>+</sup> AFFINITIES OF THE NEURONAL AND ASTROCYTIC Na<sup>+</sup>/K<sup>+</sup>-ATPase ISOFORM COMBINATIONS

The obvious manner in which the Na<sup>+</sup>/K<sup>+</sup>-ATPase may be exerting isoform-specific roles in neurophysiology may be by the distinct apparent ion affinities ( $K_m$ ) of the different  $\alpha$  and  $\beta$  isoform combinations. Of the isoforms expressed in the CNS, all subunit isoform combinations are able to assemble and be catalytically competent in heterologous expression systems (Crambert et al., 2000; Blanco, 2005; Larsen et al., 2014). The apparent ion affinities of the individual isoform combinations

have been investigated in a number of cellular and membranous systems with each their advantages and disadvantages. Most commonly employed are leaky membrane preparations (non-sided preparations) obtained from membrane extracts of diverse tissue, mammalian cell lines (i.e., HeLa, COS-1), Sf-9 insect cells, or yeast cells all expressing select Na<sup>+</sup>/K<sup>+</sup>-ATPase isoforms (Sweadner, 1985; Jewell and Lingrel, 1991; Blanco et al., 1995; Therien et al., 1996; Müller-Ehmsen et al., 2001; Toustrup-Jensen et al., 2001), although intact cellular systems (sided preparations), i.e., Na<sup>+</sup>/K<sup>+</sup>-ATPase-expressing HeLa cells or *Xenopus* oocytes have also been employed (Munzer et al., 1994; Zahler et al., 1997; Hasler et al., 1998; Crambert et al., 2000; Horisberger and Kharoubi-Hess, 2002; Larsen et al., 2014). The advantage of the leaky membrane approach is the complete control of all ion concentrations, the experimental ability to address select steps of the Na<sup>+</sup>/K<sup>+</sup>-ATPase transport cycle, and the high degree of accuracy. An inherent disadvantage is the non-physiological ion concentrations facing the two sides of the Na<sup>+</sup>/K<sup>+</sup>-ATPase, which, due to the reciprocal influence of Na<sup>+</sup> and K<sup>+</sup> on the other ion's ability to bind, affect the apparent ion affinities of the Na<sup>+</sup>/K<sup>+</sup>-ATPase (Toustrup-Jensen et al., 2001). The transmembrane potential (or lack thereof in isolated membranes) also affects Na<sup>+</sup>/K<sup>+</sup>-ATPase activity and ion affinities in an isoform-specific manner (Larsen et al., 2014, also see Figure 4A). While the apparent K<sup>+</sup> affinity is uncomplicated to approach in intact cellular systems with physiological ion concentrations and membrane potential, determination of the apparent Na<sup>+</sup> affinity at the intracellular face of the Na<sup>+</sup>/K<sup>+</sup>-ATPase presents an experimental challenge. For both approaches, expression of the different Na<sup>+</sup>/K<sup>+</sup>-ATPase  $\alpha$  isoforms in mammalian cells endogenously expressing  $\beta 1$  (Orlowski and Lingrel, 1988; Therien et al., 1996; Toustrup-Jensen et al., 2001) has prevented detailed analysis of the kinetic impact of  $\beta 2$  and  $\beta 3$  on the different  $\alpha$  isoforms. Sf9 insect cells and *Xenopus laevis* oocytes express low levels of endogenous Na<sup>+</sup>/K<sup>+</sup>-ATPase and kinetic



**FIGURE 4 | The Na<sup>+</sup>/K<sup>+</sup>-ATPase activity increases in an isoform-specific manner as a function of [K<sup>+</sup>]<sub>o</sub> and V<sub>m</sub>.** The Na<sup>+</sup>/K<sup>+</sup>-ATPase-mediated current was determined with two-electrode voltage clamp in *Xenopus* oocytes heterologously expressing the various astrocytic and neuronal subunit isoform combinations. **(A)** The  $\alpha 2\beta 2$  Na<sup>+</sup>/K<sup>+</sup>-ATPase displayed the most prominent voltage-sensitivity with increased Na<sup>+</sup>/K<sup>+</sup>-ATPase activity as a function of membrane depolarization. For clarity only  $\alpha 2\beta 2$  and  $\alpha 1\beta 1$  are illustrated, the other combinations all had intermediate I/V relationships. **(B)** the Na<sup>+</sup>/K<sup>+</sup>-ATPase-mediated current was determined as a function of [K<sup>+</sup>]<sub>o</sub> at V<sub>m</sub> = -80 mV. The  $\alpha 2\beta 2$  combination displayed a low apparent affinity for K<sup>+</sup>, K<sub>m</sub> ~3.6 mM, whereas all other tested isoform combinations saturated at lower K<sup>+</sup> concentrations, K<sub>m</sub> < 1.0 mM. Figure modified from Larsen et al. (2014) with permission.



analysis of Na<sup>+</sup>/K<sup>+</sup>-ATPase expressed in these cell types have thereby paved the road toward determination of the apparent ion affinities of all isoform combinations (Blanco et al., 1995; Crambert et al., 2000; Larsen et al., 2014). With few studies comparing all, or the majority of, isoform combinations in parallel experimental sessions, a certain diversity exists in the obtained apparent ion affinities of the Na<sup>+</sup>/K<sup>+</sup>-ATPase. The general consensus emerging for the neuronal and astrocytic isoform combinations is as follows: When paired with the ubiquitous β1 isoform, the three α isoforms display similar apparent K<sup>+</sup> affinities in the 1–2 mM range, irrespective of the sidedness of the membrane preparations (Urayama and Nakao, 1979; Sweadner, 1985; Jewell and Lingrel, 1991; Therien et al., 1996; Crambert et al., 2000; Horisberger and Kharoubi-Hess, 2002; Larsen et al., 2014), although a few studies reported a slightly reduced K<sup>+</sup> affinity of the α2 isoform (Blanco et al., 1995; Müller-Ehmsen et al., 2001). The apparent Na<sup>+</sup> affinities of α1 and α2, when in constellation with β1, have been reported at quite a range of values (with K<sub>m</sub>'s centered around 10–16 mM), although roughly similar between the isoforms in each study, with no general trend regarding which of the two displayed a tendency toward higher affinity than the other (Jewell and Lingrel, 1991; Blanco et al., 1995; Zahler et al., 1997; Crambert et al., 2000; Müller-Ehmsen et al., 2001; Horisberger and Kharoubi-Hess, 2002). α3, on the other hand, displayed a much reduced Na<sup>+</sup> affinity with values centered around K<sub>m</sub>' of 30 mM, when determined in sided systems with physiological intracellular K<sup>+</sup> concentrations (Lytton, 1985; Zahler et al., 1997; Crambert et al., 2000; Horisberger and Kharoubi-Hess, 2002; Blanco, 2005). In the leaky membrane preparations, the apparent affinity of α3 toward Na<sup>+</sup> was either similar or only slightly different from that of α1 (Sweadner, 1985; Jewell and Lingrel, 1991; Therien et al., 1996). Pairing with the astrocytic β2 isoform did not significantly alter the apparent K<sup>+</sup> affinity of α1 or α3 whereas the α2β2 combination displayed a reduced K<sup>+</sup> affinity with K<sub>m</sub>'s in the 3–5 mM range (Blanco et al., 1995; Crambert et al., 2000; Larsen et al., 2014, also see **Figure 4B**) while the apparent Na<sup>+</sup> affinity of α2 increased to around 7–8 mM upon pairing with β2 (Blanco et al., 1995). The increased apparent Na<sup>+</sup> affinity may be induced by a β2-dependent shift of the conformational equilibrium toward the E1P state (Hilbers et al., 2016). The isoform-specific relative ion affinities, which are of physiological relevance for clearance of K<sup>+</sup> from the extracellular space of the central nervous system, are summarized as follows:

$$K'_{m(K^+)} : \alpha1\beta1 \approx \alpha1\beta2 \approx \alpha2\beta1 \approx \alpha3\beta1 < \alpha2\beta2$$

$$K'_{m(Na^+)} : \alpha1\beta2 \approx \alpha2\beta2 < \alpha1\beta1 \approx \alpha2\beta1 \ll \alpha3\beta1$$

## THE PHYSIOLOGICAL ROLE OF THE Na<sup>+</sup>/K<sup>+</sup>-ATPase ISOFORM COMBINATIONS IN K<sup>+</sup> CLEARANCE

The diverse roles of the Na<sup>+</sup>/K<sup>+</sup>-ATPase in management of [K<sup>+</sup>]<sub>o</sub> in the central nervous system suggest a scenario in which explicit subunit isoform combinations perform distinct temporal


and spatial roles. During the neuronal stimulus, the astrocytic Na<sup>+</sup>/K<sup>+</sup>-ATPase α2β2 combination appears specifically geared to respond to the resultant increase in extracellular [K<sup>+</sup>] due to its low apparent K<sup>+</sup> affinity: With saturation of all other isoform combinations at basal [K<sup>+</sup>]<sub>o</sub>, an affinity constant of α2β2 in the 3–4 mM range allows this isoform combination to increase its transport activity when faced with a K<sup>+</sup> load and thus increase the rate of K<sup>+</sup> clearance from the extracellular space and into the nearby astrocytes (Larsen et al., 2014, **Figure 4B**). In addition, the membrane depolarization brought about by the increased extracellular [K<sup>+</sup>] promoted enhanced transport activity of all isoform combinations although the voltage-sensitivity was most pronounced for the α2β2 combination (Larsen et al., 2014, **Figure 4A**). The select glial expression of the Na<sup>+</sup>/K<sup>+</sup>-ATPase α2β2 isoform combination (McGrail et al., 1991; Cameron et al., 1994; Zhang et al., 2014) and localization of α2 to the glial processes surrounding neuronal dendrites (Cholet et al., 2002) taken together with the voltage- and K<sup>+</sup>-sensitivity of the α2β2 isoform combination, renders it an efficient responder to stimulus-evoked K<sup>+</sup> transients in the extracellular space and thus an important contributor to management of the extracellular K<sup>+</sup> during neuronal activity.

Given the nature of the Na<sup>+</sup>/K<sup>+</sup>-ATPase, the transport activity can, in addition to its K<sup>+</sup>-dependent activation, also be governed by activity-dependent fluctuations in the intracellular Na<sup>+</sup> concentration. As neurons propagate action potentials with subsequent synaptic activity, neuronal Na<sup>+</sup> accumulation occurs gradually via ligand- and voltage-gated Na<sup>+</sup> channels in parallel to the release of K<sup>+</sup> to the extracellular space (Langer and Rose, 2009). This [Na<sup>+</sup>]<sub>i</sub> increase prompts enhanced activation of the neuronal Na<sup>+</sup>/K<sup>+</sup>-ATPase of the isoform combination α3β1, with its characteristically low apparent Na<sup>+</sup> affinity (Lytton, 1985; Zahler et al., 1997; Crambert et al., 2000; Horisberger and Kharoubi-Hess, 2002; Blanco, 2005). The α3β1-mediated extrusion of Na<sup>+</sup> and uptake of K<sup>+</sup> permits the neurons to reestablish the concentration gradients of the two ions following the neuronal activity and the subsequent gradual release of the astrocytic K<sup>+</sup> stores via Kir4.1 (Chever et al., 2010). In this way the neuron is ready to propagate a succeeding volley of action potentials. Since the α3β1-mediated activity is governed primarily by the intracellular Na<sup>+</sup> concentration, its enhanced activity continues until normalization of [Na<sup>+</sup>]<sub>i</sub> has occurred. As this process occurs toward the termination of the neuronal activity (Grafe and Ballanyi, 1987) and is independent of the [K<sup>+</sup>]<sub>o</sub>, it may lead to an undershoot of [K<sup>+</sup>]<sub>o</sub> in which the concentration drops below baseline in the later phase of the stimulus-evoked K<sup>+</sup> transient prior to its stabilization (Ransom et al., 2000; D'Ambrosio et al., 2002).

Glutamatergic signaling represents the majority of the excitatory stimulus in the central nervous system (Danbolt, 2001). Upon vesicular release of glutamate into the synaptic cleft, glutamate is swiftly removed from the extracellular space by Na<sup>+</sup>-coupled glutamate transporters primarily located in adjacent astrocytes (Bergles and Jahr, 1997; Danbolt, 2001). Of the five cloned isoforms of the glutamate transporter (Excitatory

Amino Acid Transporters, EAAT1-5), astrocytes express EAAT1 and EAAT2, with the latter being the dominantly expressed isoform in hippocampal tissue (Bar-Peled et al., 1997; Furuta et al., 1997; Lehre and Danbolt, 1998) and estimated to account for around 90% of the glutamate uptake in the mammalian brain (Danbolt et al., 1992). During one transport cycle, inward translocation of one glutamate is driven by cotransport of three Na<sup>+</sup> and one H<sup>+</sup> while one K<sup>+</sup> is extruded (Levy et al., 1998). Astrocytic Na<sup>+</sup> accumulation thus occurs upon glutamate transporter activity in cultured astrocytes (Chatton et al., 2000; Illarionava et al., 2014) and acute brain slices (Langer and Rose, 2009; Langer et al., 2012). The astrocytic glutamate transporters display some degree of co-localization with the Na<sup>+</sup>/K<sup>+</sup>-ATPase  $\alpha$ 2 in glial leaflets surrounding glutamatergic synapses (Cholet et al., 2002; Rose et al., 2009; Rose and Karus, 2013), suggesting functional interplay between these two transport proteins, as illustrated in **Figure 5**. Indeed, in cultured astrocytes, inhibition of the Na<sup>+</sup>/K<sup>+</sup>-ATPase instantly compromised glutamate uptake (Rose et al., 2009; Bauer et al., 2012; Illarionava et al., 2014) while Na<sup>+</sup> accumulation via glutamate transporter activity (or mimicking thereof via a Na<sup>+</sup> ionophor) stimulated Na<sup>+</sup>/K<sup>+</sup>-ATPase-mediated K<sup>+</sup> uptake driven by the  $\alpha$ 2 isoform (Pellerin and Magistretti, 1997; Bender et al., 1998; Munhoz et al., 2005; Sheean et al., 2013). However, this glutamate transporter-mediated activation of Na<sup>+</sup>/K<sup>+</sup>-ATPase activity in cultured cells was observed exclusively upon prolonged (up to 20 min) exposure to glutamate (Pellerin and Magistretti, 1997; Bender et al., 1998; Munhoz et al., 2005; Sheean et al., 2013) which could suggest an indirect activation of the Na<sup>+</sup>/K<sup>+</sup>-ATPase (Munhoz et al., 2005) rather than a direct effect of a Na<sup>+</sup> transient. It therefore remains to be resolved if the glial glutamate transporter-mediated Na<sup>+</sup> transients occurring during brief neuronal activity (Langer and Rose, 2009) directly activates the glial Na<sup>+</sup>/K<sup>+</sup>-ATPase from the cytosolic face of the membrane. For the Na<sup>+</sup>/K<sup>+</sup>-ATPase to respond to stimulus-evoked intracellular Na<sup>+</sup> transients, it follows that the intracellular face of the transport protein is not already saturated at the basal glial Na<sup>+</sup> concentration of ~12 mM (Langer et al., 2012). With the above-mentioned challenges of recording the intracellular Na<sup>+</sup> affinity of various isoform combinations of the Na<sup>+</sup>/K<sup>+</sup>-ATPase in sided preparations of intact cells, especially in regard to  $\beta$ 2, it remains to be revealed which glial isoform combination(s) may display Na<sup>+</sup> affinities in a range which would allow for increased activity when faced with stimulus-induced intracellular Na<sup>+</sup> transients.

It follows that the kinetic behavior and therefore physiological specialization of the different  $\alpha$  isoforms may depend on the interacting  $\beta$  subunit and the physiological contribution of the different isoforms hinges on their quantitative expression in the different cellular structures of the CNS. Little is known about the quantitative distribution of the different isoforms at the protein level (due to challenges with comparison between different antibodies with different antigen specificity) as is their favored accessory subunit combinations, although  $\alpha$ 2 has been shown to favor interaction with the  $\beta$ 2 subunit in mouse brain extracts (Tokhtaeva et al., 2012).



**FIGURE 5 | The Na<sup>+</sup>/K<sup>+</sup>-ATPase is suggested to work in concert with the glutamate transporter.** During action potential propagation, glutamate and K<sup>+</sup> are both released from the pre-synaptic neuron. Glutamate-induced depolarization of the post-synaptic neuron instigates further K<sup>+</sup> release. By diffusion glutamate and K<sup>+</sup> reach the peri-synaptic astrocyte processes located in close proximity to the synapse. Glutamate is transported into the astrocyte along with Na<sup>+</sup> and in exchange for K<sup>+</sup> (see text for stoichiometry, H<sup>+</sup> transport has been omitted on figure for clarity). The resultant rise in intracellular Na<sup>+</sup> may trigger the activation of the Na<sup>+</sup>/K<sup>+</sup>-ATPase isoform combinations, which are not already saturated at their Na<sup>+</sup> binding site. Increased activity of the Na<sup>+</sup>/K<sup>+</sup>-ATPase would result in enhanced extrusion of the accumulated Na<sup>+</sup> and, concomitantly, in amplified removal of extracellular K<sup>+</sup> in the synaptic area. Although this proposal (based on co-localization patterns and prolonged glutamate exposure of astrocytic cultures) remains to be verified in brain slices during neuronal activity, these two transporters may work in concert in order to efficiently clear both glutamate and K<sup>+</sup> from the extracellular space.

## INVOLVEMENT OF Na<sup>+</sup>/K<sup>+</sup>-ATPase $\alpha$ 2 AND $\alpha$ 3 IN CENTRAL NERVOUS SYSTEM DISEASE

Alterations in expression of the Na<sup>+</sup>/K<sup>+</sup>-ATPase isoforms and/or dysfunctional mutant forms of the Na<sup>+</sup>/K<sup>+</sup>-ATPase may cause disturbances in the K<sup>+</sup> homeostasis and therefore affect a range of cellular parameters, i.e., membrane potential, synaptic signaling, rate of activity-evoked K<sup>+</sup> clearance, glutamate re-absorption etc. Underscoring the importance of these transport proteins for normal physiology, mice with homozygous deletion of the genes *ATP1A2* or *ATP1A3* (encoding  $\alpha$ 2 and  $\alpha$ 3) die immediately after birth (Ikeda et al., 2004, 2013). Severe neurological diseases thus arise following various mutations in *ATP1A2* and *ATP1A3* (de Vries et al., 2009; Benarroch, 2011; Böttger et al., 2012): Familial hemiplegic migraine type 2 (FHM2) arises upon mutations in *ATP1A2* (De Fusco et al., 2003) and is a severe autosomal form of migraine with aura associated with hemiparesis and sometimes accompanied by manifestations such as epilepsy, seizures, ataxia and developmental disabilities (Pietrobon, 2007; Böttger et al., 2012). A variety of different mutations in  $\alpha$ 2 has been detected in patients suffering from

FHM2 and knock-in mice expressing FHM2-derived mutant forms of  $\alpha 2$  have been established and found to mimic FHM2-relevant disease traits (Leo et al., 2011; Böttger et al., 2016 and reviewed elsewhere in this special issue by Lykke-Hartmann and colleagues): The heterozygous  $\alpha 2(+/W887R)$  knock-in mouse model displayed a lower induction threshold for cortical spreading depression (a phenomenon observed in association with migraine with aura) and an increased rate of propagation (Leo et al., 2011) whereas the  $\alpha 2(+/G301R)$  mouse model displayed mood depression, obsessive compulsive disorder, reduced regeneration after cortical spreading depression in males, as well as reduced glutamate uptake in astrocytic cultures obtained from embryonic homozygotes (Böttger et al., 2016). A selection of the mutations found in human FHM2 patients have been generated *in vitro* to determine the functional impact on the Na<sup>+</sup>/K<sup>+</sup>-ATPase activity. Diverse kinetic implications arise as a consequence of the mutations, e.g., loss-of-function, reduced catalytic turnover, shifted E1 $\leftrightarrow$ E2 steady-state conformational equilibrium, increased/decreased apparent K<sup>+</sup> affinity and increased/decreased apparent Na<sup>+</sup> affinity (the latter associated with altered basal [Na<sup>+</sup>]<sub>i</sub>) (De Fusco et al., 2003; Segall et al., 2004, 2005; Schack et al., 2012; Toustrup-Jensen et al., 2014). The diverse functional outcome of these mutations, all giving rise to the same disease, underscores the complexity of the role of  $\alpha 2$  in Na<sup>+</sup> and K<sup>+</sup> management in the central nervous system. However, it remains puzzling why such diverse alterations, e.g., oppositely directed shifts in K<sup>+</sup> affinity, disrupt neuronal signaling in a manner that gives rise to similar phenotypic outcome. Alterations in Na<sup>+</sup>/K<sup>+</sup>-ATPase  $\alpha 2$  isoform expression have been indicated in several neuropathological situations: (i) migraine (with aura) patients displayed altered Na<sup>+</sup>/K<sup>+</sup>-ATPase isoform expression toward versions with higher ouabain sensitivity (Scarrone et al., 2007), (ii) reduced K<sup>+</sup> affinity of partially purified Na<sup>+</sup>/K<sup>+</sup>-ATPase from neocortices of human epileptic patients (Guillaume et al., 1991), (iii) reduced K<sup>+</sup> clearance rates observed in cortex of epileptic monkeys (Lewis et al., 1977), and (iv) glial fractions obtained from cats following freezing lesion-induced epilepsy lost their characteristic K<sup>+</sup>-activation, suggesting that  $\alpha 2$  and/or  $\beta 2$  are down regulated during the time between the freezing lesion and the occurrence of the epileptogenic states (Grisar et al., 1983). Oppositely, ciliary neurotrophic factor up regulated glial  $\alpha 2$  mRNA with subsequent enhanced K<sup>+</sup> uptake and increased threshold for spreading depolarization (Seidel et al., 2015).

Mutations in the gene *ATPIA3* encoding the  $\alpha 3$  isoform have been discovered in patients with rapid-onset dystonia parkinsonism (RDP), alternating hemiplegia of childhood (AHC), and early life epilepsy (de Carvalho Aguiar et al.,

2004; Böttger et al., 2012; Heinzen et al., 2014; Sweney et al., 2015). Expressional analysis of  $\alpha 3$  demonstrated the presence of this subunit in anatomical regions correlating with RDP symptoms (Böttger et al., 2011). Mice with heterozygous deletion of *ATPIA3* displayed movement abnormalities partly in line with the human RDP phenotype (Ikeda et al., 2013; Sugimoto et al., 2014) and were more susceptible to stress-induced depression-like phenotypes (Kirshenbaum et al., 2011). Mice expressing mutant forms of  $\alpha 3$  exhibit a phenotype comparable to that of patients with AHC and display hyperexcitability, K<sup>+</sup>-induced spreading depolarization of longer duration, and enhanced seizure susceptibility (Clapcote et al., 2009; Hunanyan et al., 2015; Kirshenbaum et al., 2015, reviewed in detail elsewhere in this special issue by Lykke-Hartmann and colleagues). Select human mutations in *ATPIA3* have been established *in vitro* and the kinetic alterations of the mutated  $\alpha 3$  isoforms ranged from loss of function to reduced activity and modified cation binding (Einholm et al., 2010; Toustrup-Jensen et al., 2014; Weigand et al., 2014).

## CONCLUSION

The Na<sup>+</sup>/K<sup>+</sup>-ATPase exhibits an isoform-specific expression pattern in the mammalian CNS. The distinct kinetic properties of the different catalytic  $\alpha$  isoforms hinge, in part, on the accessory  $\beta$  subunit with which they pair and determine their temporal and spatial quantitative contribution to management of the extracellular K<sup>+</sup> transients occurring in the wake of neuronal activity. While the astrocytic  $\alpha 2$  isoform appears specifically geared to respond to the activity-evoked K<sup>+</sup> transients upon pairing with  $\beta 2$ , the excessively low apparent Na<sup>+</sup> affinity of the  $\alpha 3$  ensures neuronal re-uptake of K<sup>+</sup> once the neuronal activity is terminated. Mutations in the genes encoding the neuronal  $\alpha 3$  or the astrocytic  $\alpha 2$  give rise to a range of severe neuronal pathologies, although the exact mechanism coupling each point mutation to a distinct disease pattern remains unresolved.

## AUTHOR CONTRIBUTIONS

All authors listed have made substantial, direct and intellectual contribution to the work, and approved it for publication.

## FUNDING

The laboratory is funded by grants from the Lundbeck Foundation, Danish Medical Research Council and Thorberg's Foundation.

## REFERENCES

- Ballanyi, K., Grafe, P., and ten Bruggencate, G. (1987). Ion activities and potassium uptake mechanisms of glial cells in guinea-pig olfactory cortex slices. *J. Physiol.* 382, 159–174. doi: 10.1113/jphysiol.1987.sp016361
- Bar-Peled, O., Ben-Hur, H., Biegón, A., Groner, Y., Dewhurst, S., Furuta, A., et al. (1997). Distribution of glutamate transporter subtypes during human brain development. *J. Neurochem.* 69, 2571–2580. doi: 10.1046/j.1471-4159.1997.69062571.x
- Bauer, D. E., Jackson, J. G., Genda, E. N., Montoya, M. M., Yudkoff, M., and Robinson, M. B. (2012). The glutamate transporter, GLAST, participates in a macromolecular complex that supports glutamate metabolism. *Neurochem. Int.* 61, 566–574. doi: 10.1016/j.neuint.2012.01.013



- Benarroch, E. E. (2011). Na<sup>+</sup>, K<sup>+</sup>-ATPase: Functions in the nervous system and involvement in neurologic disease. *Neurology* 76, 287–293. doi: 10.1212/WNL.0b013e3182074c2f
- Bender, A. S., Schousboe, A., Reichelt, W., and Norenberg, M. D. (1998). Ionic mechanisms in glutamate-induced astrocyte swelling: role of K<sup>+</sup> influx. *J. Neurosci. Res.* 52, 307–321.
- Bergles, D. E., and Jahr, C. E. (1997). Synaptic activation of glutamate transporters in hippocampal astrocytes. *Neuron* 19, 1297–1308. doi: 10.1016/S0896-6273(00)80420-1
- Blanco, G. (2005). Na,K-ATPase subunit heterogeneity as a mechanism for tissue-specific ion regulation. *Semin. Nephrol.* 25, 292–303. doi: 10.1016/j.semnephrol.2005.03.004
- Blanco, G., Koster, J. C., Sanchez, G., and Mercer, R. W. (1995). Kinetic Properties of the alpha.2.beta.1 and alpha.2.beta.2 Isoenzymes of the Na,K-ATPase. *Biochemistry* 34, 319–325. doi: 10.1021/bi00001a039
- Böttger, P., Doğanli, C., and Lykke-Hartmann, K. (2012). Migraine- and dystonia-related disease-mutations of Na<sup>+</sup>/K<sup>+</sup>-ATPases: relevance of behavioral studies in mice to disease symptoms and neurological manifestations in humans. *Neurosci. Biobehav. Rev.* 36, 855–871. doi: 10.1016/j.neubiorev.2011.10.005
- Böttger, P., Glerup, S., Gesslein, B., Illarionova, N. B., Isaksen, T. J., Heuck, A., et al. (2016). Glutamate-system defects behind psychiatric manifestations in a familial hemiplegic migraine type 2 disease-mutation mouse model. *Sci. Rep.* 6:22047. doi: 10.1038/srep22047
- Böttger, P., Tracz, Z., Heuck, A., Nissen, P., Romero-Ramos, M., and Lykke-Hartmann, K. (2011). Distribution of Na/K-ATPase alpha 3 isoform, a sodium-potassium P-type pump associated with rapid-onset of dystonia parkinsonism (RDP) in the adult mouse brain. *J. Comp. Neurol.* 519, 376–404. doi: 10.1002/cne.22524
- Cameron, R., Klein, L., Shyjan, A., Rakic, P., and Levenson, R. (1994). Neurons and astroglia express distinct subsets of Na,K-ATPase alpha and beta subunits. *Brain Res. Mol. Brain Res.* 21, 333–343. doi: 10.1016/0169-328X(94)90264-X
- Chatton, J.-Y., Marquet, P., and Magistretti, P. J. (2000). A quantitative analysis of l-glutamate-regulated Na<sup>+</sup> dynamics in mouse cortical astrocytes: implications for cellular bioenergetics. *Eur. J. Neurosci.* 12, 3843–3853. doi: 10.1046/j.1460-9568.2000.00269.x
- Chever, O., Djukic, B., McCarthy, K. D., and Amzica, F. (2010). Implication of Kir4.1 channel in excess potassium clearance: an *in vivo* study on anesthetized glial-conditional Kir4.1 knock-out mice. *J. Neurosci.* 30, 15769–15777. doi: 10.1523/JNEUROSCI.2078-10.2010
- Cholet, N., Pellerin, L., Magistretti, P. J., and Hamel, E. (2002). Similar perisynaptic glial localization for the Na<sup>+</sup>,K<sup>+</sup>-ATPase alpha 2 subunit and the glutamate transporters GLAST and GLT-1 in the rat somatosensory cortex. *Cereb. Cortex* 12, 515–525. doi: 10.1093/cercor/12.5.515
- Clapcote, S. J., Duffy, S., Xie, G., Kirshenbaum, G., Bechard, A. R., Rodacker Schack, V., et al. (2009). Mutation I810N in the alpha 3 isoform of Na<sup>+</sup>,K<sup>+</sup>-ATPase causes impairments in the sodium pump and hyperexcitability in the CNS. *Proc. Natl. Acad. Sci. U.S.A.* 106, 14085–14090. doi: 10.1073/pnas.0904817106
- Clayton, G. H., Owens, G. C., Wolff, J. S., and Roderic, L. S. (1998). Ontogeny of cation-Cl<sup>-</sup> cotransporter expression in rat neocortex. *Dev. Brain Res.* 109, 281–292. doi: 10.1016/S0165-3806(98)00078-9
- Crambert, G., Hasler, U., Beggah, A. T., Yu, C., Modyanov, N. N., Horisberger, J.-D., et al. (2000). Transport and pharmacological properties of nine different human Na,K-ATPase isozymes. *J. Biol. Chem.* 275, 1976–1986. doi: 10.1074/jbc.275.3.1976
- D'Ambrosio, R., Gordon, D. S., and Winn, H. R. (2002). Differential role of KIR channel and Na<sup>+</sup>/K<sup>+</sup>-pump in the regulation of extracellular K<sup>+</sup> in rat hippocampus. *J. Neurophysiol.* 87, 87–102. doi: 10.1152/jn.00240.2001
- Danbolt, N. C. (2001). Glutamate uptake. *Prog. Neurobiol.* 65, 1–105. doi: 10.1016/S0301-0082(00)00067-8
- Danbolt, N. C., Storm-Mathisen, J., and Kanner, B. I. (1992). An [Na<sup>+</sup> + K<sup>+</sup>]coupled l-glutamate transporter purified from rat brain is located in glial cell processes. *Neuroscience* 51, 295–310. doi: 10.1016/0306-4522(92)90316-T
- de Carvalho Aguiar, P., Swadner, K. J., Penniston, J. T., Zaremba, J., Liu, L., Caton, M., et al. (2004). Mutations in the Na<sup>+</sup>/K<sup>+</sup>-ATPase alpha 3 gene ATP1A3 are associated with rapid-onset dystonia parkinsonism. *Neuron* 43, 169–175. doi: 10.1016/j.neuron.2004.06.028
- De Fusco, M., Marconi, R., Silvestri, L., Atorino, L., Rampoldi, L., Morgante, L., et al. (2003). Haploinsufficiency of ATP1A2 encoding the Na<sup>+</sup>/K<sup>+</sup> pump [alpha]2 subunit associated with familial hemiplegic migraine type 2. *Nat. Genet.* 33, 192–196. doi: 10.1038/ng1081
- de Vries, B., Frants, R. R., Ferrari, M. D., and van den Maagdenberg, A. M. J. M. (2009). Molecular genetics of migraine. *Hum. Genet.* 126, 115–132. doi: 10.1007/s00439-009-0684-z
- Einholm, A. P., Toustrup-Jensen, M. S., Holm, R., Andersen, J. P., and Vilsen, B. (2010). The rapid-onset dystonia parkinsonism mutation D923N of the Na<sup>+</sup>,K<sup>+</sup>-ATPase alpha 3 isoform disrupts Na<sup>+</sup> interaction at the third Na<sup>+</sup> site. *J. Biol. Chem.* 285, 26245–26254. doi: 10.1074/jbc.M110.123976
- Frankenhaeuser, B., and Hodgkin, A. L. (1956). The after-effects of impulses in the giant nerve fibres of Loligo. *J. Physiol.* 131, 341–376. doi: 10.1113/jphysiol.1956.sp005467
- Furuta, A., Rothstein, J. D., and Martin, L. J. (1997). Glutamate transporter protein subtypes are expressed differentially during rat CNS development. *J. Neurosci.* 17, 8363–8375.
- Futamachi, K. J., Mutani, R., and Prince, D. A. (1974). Potassium activity in rabbit cortex. *Brain Res.* 75, 5–25. doi: 10.1016/0006-8993(74)90767-7
- Gardner-Medwin, A. R. (1983). A study of the mechanisms by which potassium moves through brain tissue in the rat. *J. Physiol.* 335, 353–374. doi: 10.1113/jphysiol.1983.sp014539
- Grafe, P., and Ballanyi, K. (1987). Cellular mechanisms of potassium homeostasis in the mammalian nervous system. *Can. J. Physiol. Pharmacol.* 65, 1038–1042. doi: 10.1139/y87-164
- Grisar, T., Franck, G., and Delgado-Escueta, A. V. (1983). Glial contribution to seizure: K<sup>+</sup> activation of (Na<sup>+</sup>, K<sup>+</sup>)-ATPase in bulk isolated glial cells and synaptosomes of epileptogenic cortex. *Brain Res.* 261, 75–84. doi: 10.1016/0006-8993(83)91285-4
- Grisar, T., Franck, G., and Schoffeniels, E. (1980). Glial control of neuronal excitability in mammals: II. Enzymatic evidence: two molecular forms of the (Na<sup>+</sup>,K<sup>+</sup>)-ATPase in brain. *Neurochem. Int.* 2, 311–320. doi: 10.1016/0197-0186(80)90038-8
- Guillaume, D., Grisar, T., Delgado-Escueta, A. V., Laschet, J., and Bureau-Heeren, M. (1991). Phosphorylation of brain (Na<sup>+</sup>,K<sup>+</sup>)-ATPase alpha catalytic subunits in normal and epileptic cerebral cortex: II. Partial seizures in human epilepsy. *J. Neurosci. Res.* 29, 218–224. doi: 10.1002/jnr.490290212
- Gulledge, A. T., Dasari, S., Onoue, K., Stephens, E. K., Hasse, J. M., and Avesar, D. (2013). A sodium-pump-mediated afterhyperpolarization in pyramidal neurons. *J. Neurosci.* 33, 13025–13041. doi: 10.1523/JNEUROSCI.0220-13.2013
- Haglund, M. M., and Schwartzkroin, P. A. (1990). Role of Na-K pump potassium regulation and IPSPs in seizures and spreading depression in immature rabbit hippocampal slices. *J. Neurophysiol.* 63, 225–239.
- Hajek, I., Subbarao, K. V. S., and Hertz, L. (1996). Acute and chronic effects of potassium and noradrenaline on Na<sup>+</sup>, K<sup>+</sup>-ATPase activity in cultured mouse neurons and astrocytes. *Neurochem. Int.* 28, 335–342. doi: 10.1016/0197-0186(95)00081-X
- Hasler, U., Wang, X., Crambert, G., Béguin, P., Jaisser, F., Horisberger, J.-D., et al. (1998). Role of beta-subunit domains in the assembly, stable expression, intracellular routing, and functional properties of Na,K-ATPase. *J. Biol. Chem.* 273, 30826–30835. doi: 10.1074/jbc.273.46.30826
- Heinemann, U., and Lux, H. D. (1977). Ceiling of stimulus induced rises in extracellular potassium concentration in the cerebral cortex of cat. *Brain Res.* 120, 231–249. doi: 10.1016/0006-8993(77)90903-9
- Heinzen, E. L., Arzimanoglou, A., Brashear, A., Clapcote, S. J., Gurreri, F., Goldstein, D. B., et al. (2014). Distinct neurological disorders with ATP1A3 mutations. *Lancet Neurol.* 13, 503–514. doi: 10.1016/S1474-4422(14)70011-0
- Hertz, L., Song, D., Xu, J., Peng, L., and Gibbs, M. (2015). Role of the astrocytic Na<sup>+</sup>, K<sup>+</sup>-ATPase in K<sup>+</sup> Homeostasis in brain: K<sup>+</sup> uptake, signaling pathways and substrate utilization. *Neurochem. Res.* 40, 2505–2516. doi: 10.1007/s11064-014-1505-x
- Hilbers, F., Kopec, W., Isaksen, T. J., Holm, T. H., Lykke-Hartmann, K., Nissen, P., et al. (2016). Tuning of the Na,K-ATPase by the beta subunit. *Sci. Rep.* 6:20442. doi: 10.1038/srep20442
- Horisberger, J.-D., and Kharoubi-Hess, S. (2002). Functional differences between alpha subunit isoforms of the rat Na,K-ATPase expressed in *Xenopus* oocytes. *J. Physiol.* 539, 669–680. doi: 10.1113/jphysiol.2001.013201
- Humanyan, A. S., Fainberg, N. A., Linabarger, M., Arehart, E., Leonard, A. S., Adil, S. M., et al. (2015). Knock-in mouse model of alternating hemiplegia

- of childhood: behavioral and electrophysiologic characterization. *Epilepsia* 56, 82–93. doi: 10.1111/epi.12878
- Ikeda, K., Onimaru, H., Yamada, J., Inoue, K., Ueno, S., Onaka, T., et al. (2004). Malfunction of respiratory-related neuronal activity in Na<sup>+</sup>, K<sup>+</sup>-ATPase  $\alpha$ 2 subunit-deficient mice is attributable to abnormal Cl<sup>-</sup> homeostasis in brainstem neurons. *J. Neurosci.* 24, 10693–10701. doi: 10.1523/JNEUROSCI.2909-04.2004
- Ikeda, K., Satake, S. I., Onaka, T., Sugimoto, H., Takeda, N., Imoto, K., et al. (2013). Enhanced inhibitory neurotransmission in the cerebellar cortex of Atp1a3-deficient heterozygous mice. *J. Physiol.* 591, 3433–3449. doi: 10.1113/jphysiol.2012.247817
- Illarionova, N. B., Brismar, H., Aperia, A., and Gunnarsson, E. (2014). Role of Na<sup>+</sup>,K<sup>+</sup>-ATPase  $\alpha$ 1 and  $\alpha$ 2 isoforms in the support of astrocyte glutamate uptake. *PLoS ONE* 9:e98469. doi: 10.1371/journal.pone.0098469
- Jewell, E. A., and Lingrel, J. B. (1991). Comparison of the substrate dependence properties of the rat Na<sup>+</sup>,K<sup>+</sup>-ATPase alpha 1, alpha 2, and alpha 3 isoforms expressed in HeLa cells. *J. Biol. Chem.* 266, 16925–16930.
- Juhászová, M., and Blaustein, M. P. (1997). Na<sup>+</sup> pump low and high ouabain affinity  $\alpha$  subunit isoforms are differently distributed in cells. *Proc. Natl. Acad. Sci. U.S.A.* 94, 1800–1805. doi: 10.1073/pnas.94.5.1800
- Karwoski, C., Lu, H.-K., and Newman, E. (1989). Spatial buffering of light-evoked potassium increases by retinal müller (Glial) cells. *Science* 244, 578–580. doi: 10.1126/science.2785716
- Kimelberg, H. K., Biddlecome, S., Narumi, S., and Bourke, R. S. (1978). ATPase and carbonic anhydrase activities of bulk-isolated neuron, glia and synaptosome fractions from rat brain. *Brain Res.* 141, 305–323. doi: 10.1016/0006-8993(78)90200-7
- Kirshenbaum, G. S., Dachtler, J., Roder, J. C., and Clapcote, S. J. (2015). Characterization of cognitive deficits in mice with an alternating hemiplegia-linked mutation. *Behav. Neurosci.* 129, 822–831. doi: 10.1037/bne0000097
- Kirshenbaum, G. S., Saltzman, K., Rose, B., Petersen, J., Vilsen, B., and Roder, J. C. (2011). Decreased neuronal Na<sup>+</sup>,K<sup>+</sup>-ATPase activity in Atp1a3 heterozygous mice increases susceptibility to depression-like endophenotypes by chronic variable stress. *Genes Brain Behav.* 10, 542–550. doi: 10.1111/j.1601-183X.2011.00691.x
- Kofuji, P., and Newman, E. A. (2004). Potassium buffering in the central nervous system. *Neuroscience* 129, 1043–1054. doi: 10.1016/j.neuroscience.2004.06.008
- Krnjević, K., and Morris, M. E. (1972). Extracellular K<sup>+</sup> activity and slow potential changes in spinal cord and medulla. *Can. J. Physiol. Pharmacol.* 50, 1214–1217. doi: 10.1139/y72-177
- Kuffler, S. W., and Nicholls, J. (1964). Glial cells in the central nervous system of the leech; their membrane potential and potassium content. *Naunyn Schmiedeberg's Arch. Exp. Pathol. Pharmacol.* 248, 216–222.
- Kuffler, S. W., and Potter, D. D. (1964). Glia in the leech central nervous system: physiological properties and neuron-glia relationship. *J. Neurophysiol.* 27, 290–320.
- Langer, J., and Rose, C. R. (2009). Synaptically induced sodium signals in hippocampal astrocytes *in situ*. *J. Physiol.* 587, 5859–5877. doi: 10.1113/jphysiol.2009.182279
- Langer, J., Stephan, J., Theis, M., and Rose, C. R. (2012). Gap junctions mediate intercellular spread of sodium between hippocampal astrocytes *in situ*. *Glia* 60, 239–252. doi: 10.1002/glia.21259
- Larsen, B. R., Assentoft, M., Cotrina, M. L., Hua, S. Z., Nedergaard, M., Kaila, K., et al. (2014). Contributions of the Na<sup>+</sup>/K<sup>+</sup>-ATPase, NKCC1, and Kir4.1 to hippocampal K<sup>+</sup> clearance and volume responses. *Glia* 62, 608–622. doi: 10.1002/glia.22629
- Larsen, B. R., and MacAulay, N. (2014). Kir4.1-mediated spatial buffering of K<sup>+</sup>: experimental challenges in determination of its temporal and quantitative contribution to K<sup>+</sup> clearance in the brain. *Channels* 8, 544–550. doi: 10.4161/19336950.2014.970448
- Lehre, K. P., and Danbolt, N. C. (1998). The number of glutamate transporter subtype molecules at glutamatergic synapses: chemical and stereological quantification in young adult rat brain. *J. Neurosci.* 18, 8751–8757.
- Leo, L., Gherardini, L., Barone, V., De Fusco, M., Pietrobon, D., Pizzorusso, T., et al. (2011). Increased susceptibility to cortical spreading depression in the mouse model of familial hemiplegic migraine type 2. *PLoS Genet.* 7:e1002129. doi: 10.1371/journal.pgen.1002129
- Levy, L. M., Warr, O., and Attwell, D. (1998). Stoichiometry of the glial glutamate transporter GLT-1 expressed inducibly in a Chinese hamster ovary cell line selected for low endogenous Na<sup>+</sup>-dependent glutamate uptake. *J. Neurosci.* 18, 9620–9628.
- Lewis, D. V., Mutsaers, N., Schuette, W. H., and Van Buren, J. (1977). Potassium clearance and reactive gliosis in the alumina gel lesion. *Epilepsia* 18, 499–506. doi: 10.1111/j.1528-1157.1977.tb04997.x
- Li, B., Hertz, L., and Peng, L. (2013). Cell-specific mRNA alterations in Na<sup>+</sup>, K<sup>+</sup>-ATPase  $\alpha$  and  $\beta$  isoforms and FXFD in mice treated chronically with carbamazepine, an anti-bipolar drug. *Neurochem. Res.* 38, 834–841. doi: 10.1007/s11064-013-0986-3
- Lytton, J. (1985). Insulin affects the sodium affinity of the rat adipocyte (Na<sup>+</sup>,K<sup>+</sup>)-ATPase. *J. Biol. Chem.* 260, 10075–10080.
- MacAulay, N., and Zeuthen, T. (2012). Glial K<sup>+</sup> clearance and cell swelling: key roles for cotransporters and pumps. *Neurochem. Res.* 37, 2299–2309. doi: 10.1007/s11064-012-0731-3
- Martin-Vasallo, P., Wetzel, R. K., García-Segura, L. M., Molina-Holgado, E., Arystarkhova, E., and Sweadner, K. J. (2000). Oligodendrocytes in brain and optic nerve express the  $\beta$ 3 subunit isoform of Na<sup>+</sup>,K<sup>+</sup>-ATPase. *Glia* 31, 206–218. doi: 10.1002/1098-1136(200009)31:3<206::AID-GLIA20>3.0.CO;2-1
- McGrail, K. M., Phillips, J. M., and Sweadner, K. M. (1991). Immunofluorescent localization of three Na<sup>+</sup>,K<sup>+</sup>-ATPase isozymes in the rat central nervous system: both neurons and glia can express more than one Na<sup>+</sup>,K<sup>+</sup>-ATPase. *J. Neurosci.* 11, 381–391.
- McGrail, K. M., and Sweadner, K. M. (1986). Immunofluorescent localization of two different Na<sup>+</sup>,K<sup>+</sup>-ATPases in the rat retina and in identified dissociated retinal cells. *J. Neurosci.* 6, 1272–1283.
- Moseley, A. E., Lieske, S. P., Wetzel, R. K., James, P. F., He, S., Shelly, D. A., et al. (2003). The Na<sup>+</sup>,K<sup>+</sup>-ATPase  $\alpha$ 2 isoform is expressed in neurons, and its absence disrupts neuronal activity in newborn mice. *J. Biol. Chem.* 278, 5317–5324. doi: 10.1074/jbc.M211315200
- Müller-Ehmsen, J., Juvvadi, P., Thompson, C. B., Tumyan, L., Croyle, M., Lingrel, J. B., et al. (2001). Ouabain and substrate affinities of human Na<sup>+</sup>-K<sup>+</sup>-ATPase  $\alpha$ 1 $\beta$ 1,  $\alpha$ 2 $\beta$ 1, and  $\alpha$ 3 $\beta$ 1 when expressed separately in yeast cells. *Am. J. Physiol. Cell Physiol.* 281, C1355–C1364.
- Munhoz, C. D., Kawamoto, E. M., de Sá Lima, L., Lepsch, L. B., Glezer, I., Marcourakis, T., et al. (2005). Glutamate modulates sodium-potassium-ATPase through cyclic GMP and cyclic GMP-dependent protein kinase in rat striatum. *Cell Biochem. Funct.* 23, 115–123. doi: 10.1002/cbf.1217
- Munzer, J. S., Daly, S. E., Jewell-Motz, E. A., Lingrel, J. B., and Blostein, R. (1994). Tissue- and isoform-specific kinetic behavior of the Na<sup>+</sup>,K<sup>+</sup>-ATPase. *J. Biol. Chem.* 269, 16668–16676.
- Orkand, R. K., Nicholls, J. G., and Kuffler, S. W. (1966). Effect of nerve impulses on the membrane potential of glial cells in the central nervous system of amphibia. *J. Neurophysiol.* 29, 788–806.
- Orlowski, J., and Lingrel, J. B. (1988). Tissue-specific and developmental regulation of rat Na<sup>+</sup>,K<sup>+</sup>-ATPase catalytic alpha isoform and beta subunit mRNAs. *J. Biol. Chem.* 263, 10436–10442.
- Pellerin, L., and Magistretti, P. J. (1997). Glutamate uptake stimulates Na<sup>+</sup>,K<sup>+</sup>-ATPase activity in astrocytes via activation of a distinct subunit highly sensitive to ouabain. *J. Neurochem.* 69, 2132–2137. doi: 10.1046/j.1471-4159.1997.69052132.x
- Peng, L., Martin-Vasallo, P., and Sweadner, K. J. (1997). Isoforms of Na<sup>+</sup>,K<sup>+</sup>-ATPase  $\alpha$  and  $\beta$  subunits in the rat cerebellum and in granule cell cultures. *J. Neurosci.* 17, 3488–3502.
- Pietrobon, D. (2007). Familial hemiplegic migraine. *Neurotherapeutics* 4, 274–284. doi: 10.1016/j.nurt.2007.01.008
- Plotkin, M. D., Kaplan, M. R., Peterson, L. N., Gullans, S. R., Hebert, S. C., and Delpire, E. (1997). Expression of the Na<sup>+</sup>(+)-K<sup>+</sup>(+)-2Cl<sup>-</sup> cotransporter BSC2 in the nervous system. *Am. J. Physiol. Cell Physiol.* 272(Pt 1), C173–C183.
- Prince, D. A., Lux, H. D., and Neher, E. (1973). Measurement of extracellular potassium activity in cat cortex. *Brain Res.* 50, 489–495. doi: 10.1016/0006-8993(73)90758-0
- Raat, N. J., Delpire, E., van Os, C. H., and Bindels, R. J. (1996). Culturing induced expression of basolateral Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransporter BSC2 in proximal tubule, aortic endothelium, and vascular smooth muscle. *Pflugers Arch.* 431, 458–460. doi: 10.1007/bf02207286

- Ransom, C. B., Ransom, B. R., and Sontheimer, H. (2000). Activity-dependent extracellular K<sup>+</sup> accumulation in rat optic nerve: the role of glial and axonal Na<sup>+</sup> pumps. *J. Physiol.* 522, 427–442. doi: 10.1111/j.1469-7793.2000.00427.x
- Richards, K. S., Bommert, K., Szabo, G., and Miles, R. (2007). Differential expression of Na<sup>+</sup>/K<sup>+</sup>-ATPase  $\alpha$ -subunits in mouse hippocampal interneurons and pyramidal cells. *J. Physiol.* 585, 491–505. doi: 10.1113/jphysiol.2007.144733
- Rose, C. R., and Karus, C. (2013). Two sides of the same coin: sodium homeostasis and signaling in astrocytes under physiological and pathophysiological conditions. *Glia* 61, 1191–1205. doi: 10.1002/glia.22492
- Rose, E. M., Koo, J. C. P., Antflick, J. E., Ahmed, S. M., Angers, S., and Hampson, D. R. (2009). Glutamate transporter coupling to Na<sub>2</sub>K-ATPase. *J. Neurosci.* 29, 8143–8155. doi: 10.1523/JNEUROSCI.1081-09.2009
- Scarrone, S., Podestà, M., Cupello, A., Finocchi, C., Frassoni, F., Gandolfo, C., et al. (2007). Abnormalities of Na/K ATPase in migraine with aura. *Cephalalgia* 27, 128–132. doi: 10.1111/j.1468-2982.2007.01267.x
- Schack, V. R., Holm, R., and Vilsen, B. (2012). Inhibition of Phosphorylation of Na<sup>+</sup>,K<sup>+</sup>-ATPase by mutations causing familial hemiplegic migraine. *J. Biol. Chem.* 287, 2191–2202. doi: 10.1074/jbc.M111.323022
- Segall, L., Mezzetti, A., Scanzano, R., Gargus, J. J., Purisima, E., and Blostein, R. (2005). Alterations in the  $\alpha$ 2 isoform of Na<sub>2</sub>K-ATPase associated with familial hemiplegic migraine type 2. *Proc. Natl. Acad. Sci. U.S.A.* 102, 11106–11111. doi: 10.1073/pnas.0504323102
- Segall, L., Scanzano, R., Kaunisto, M. A., Wessman, M., Palotie, A., Gargus, J. J., et al. (2004). Kinetic alterations due to a missense mutation in the Na<sub>2</sub>K-ATPase  $\alpha$ 2 subunit cause familial hemiplegic migraine type 2. *J. Biol. Chem.* 279, 43692–43696. doi: 10.1074/jbc.M407471200
- Seidel, J. L., Faideau, M., Aiba, I., Pannasch, U., Escartin, C., Rouach, N., et al. (2015). Ciliary neurotrophic factor (CNTF) activation of astrocytes decreases spreading depolarization susceptibility and increases potassium clearance. *Glia* 63, 91–103. doi: 10.1002/glia.22735
- Sheean, R. K., Lau, C. L., Shin, Y. S., O'Shea, R. D., and Beart, P. M. (2013). Links between l-glutamate transporters, Na<sup>+</sup>/K<sup>+</sup>-ATPase and cytoskeleton in astrocytes: Evidence following inhibition with rottlerin. *Neuroscience* 254, 335–346. doi: 10.1016/j.neuroscience.2013.09.043
- Singer, W., and Lux, H. D. (1975). Extracellular potassium gradients and visual receptive fields in the cat striate cortex. *Brain Res.* 96, 378–383. doi: 10.1016/0006-8993(75)90751-9
- Su, G., Haworth, R. A., Dempsey, R. J., and Sun, D. (2000). Regulation of Na<sup>+</sup>-K<sup>+</sup>-Cl<sup>-</sup>-cotransporter in primary astrocytes by dibutyryl cAMP and high [K<sup>+</sup>]<sub>o</sub>. *Am. J. Physiol. Cell Physiol.* 279, C1710–C1721.
- Sugimoto, H., Ikeda, K., and Kawakami, K. (2014). Heterozygous mice deficient in Atp1a3 exhibit motor deficits by chronic restraint stress. *Behav. Brain Res.* 272, 100–110. doi: 10.1016/j.bbr.2014.06.048
- Swadner, K. J. (1985). Enzymatic properties of separated isozymes of the Na<sub>2</sub>K-ATPase. Substrate affinities, kinetic cooperativity, and ion transport stoichiometry. *J. Biol. Chem.* 260, 11508–11513.
- Sweney, M. T., Newcomb, T. M., and Swoboda, K. J. (2015). The expanding spectrum of neurological phenotypes in children with ATP1A3 mutations, alternating hemiplegia of childhood, rapid-onset dystonia-parkinsonism, CAPOS and beyond. *Pediatr. Neurol.* 52, 56–64. doi: 10.1016/j.pediatrneurol.2014.09.015
- Syková, E., Rothenberg, S., and Krekule, I. (1974). Changes of extracellular potassium concentration during spontaneous activity in the mesencephalic reticular formation of the rat. *Brain Res.* 79, 333–337. doi: 10.1016/0006-8993(74)90428-4
- Therien, A. G., Nestor, N. B., Ball, W. J., and Blostein, R. (1996). Tissue-specific versus isoform-specific differences in cation activation kinetics of the Na<sub>2</sub>K-ATPase. *J. Biol. Chem.* 271, 7104–7112. doi: 10.1074/jbc.271.12.7104
- Tokhtaeva, E., Clifford, R. J., Kaplan, J. H., Sachs, G., and Vagin, O. (2012). Subunit isoform selectivity in assembly of Na<sub>2</sub>K-ATPase  $\alpha$ - $\beta$  heterodimers. *J. Biol. Chem.* 287, 26115–26125. doi: 10.1074/jbc.M112.370734
- Toustrup-Jensen, M., Hauge, M., and Vilsen, B. (2001). Mutational effects on conformational changes of the dephospho- and phospho-forms of the Na<sup>+</sup>,K<sup>+</sup>-ATPase<sup>f</sup>. *Biochemistry* 40, 5521–5532. doi: 10.1021/bi002367m
- Toustrup-Jensen, M. S., Einholm, A. P., Schack, V. R., Nielsen, H. N., Holm, R., Sobrido, M.-J., et al. (2014). Relationship between intracellular Na<sup>+</sup> concentration and reduced Na<sup>+</sup> affinity in Na<sup>+</sup>,K<sup>+</sup>-ATPase mutants causing neurological disease. *J. Biol. Chem.* 289, 3186–3197. doi: 10.1074/jbc.M113.543272
- Urayama, O., and Nakao, M. (1979). Organ specificity of rat sodium- and potassium-activated adenosine triphosphatase. *J. Biochem.* 86, 1371–1381.
- Walz, W. (2000). Role of astrocytes in the clearance of excess extracellular potassium. *Neurochem. Int.* 36, 291–300. doi: 10.1016/S0197-0186(99)00137-0
- Walz, W., and Hertz, L. (1982). Ouabain-sensitive and ouabain-resistant net uptake of potassium into astrocytes and neurons in primary cultures. *J. Neurochem.* 39, 70–77. doi: 10.1111/j.1471-4159.1982.tb04702.x
- Walz, W., and Hertz, L. (1984). Intense furosemide-sensitive potassium accumulation in astrocytes in the presence of pathologically high extracellular potassium levels. *J. Cereb. Blood Flow Metab.* 4, 301–304. doi: 10.1038/jcbfm.1984.42
- Wang, F., Smith, N. A., Xu, Q., Fujita, T., Baba, A., Matsuda, T., et al. (2012). Astrocytes modulate neural network activity by Ca<sup>2+</sup>-dependent uptake of extracellular K<sup>+</sup>. *Sci. Signal.* 5:ra26. doi: 10.1126/scisignal.2002334
- Watts, A. G., Sanchez-Watts, G., Emanuel, J. R., and Levenson, R. (1991). Cell-specific expression of mRNAs encoding Na<sup>+</sup>,K<sup>(+)</sup>-ATPase alpha- and beta-subunit isoforms within the rat central nervous system. *Proc. Natl. Acad. Sci. U.S.A.* 88, 7425–7429. doi: 10.1073/pnas.88.16.7425
- Weigand, K. M., Messchaert, M., Swarts, H. G. P., Russel, F. G. M., and Koenderink, J. B. (2014). Alternating Hemiplegia of Childhood mutations have a differential effect on Na<sup>+</sup>,K<sup>+</sup>-ATPase activity and ouabain binding. *Biochim. Biophys. Acta* 1842, 1010–1016. doi: 10.1016/j.bbdis.2014.03.002
- White, H. S., Chow, S. Y., Yen-Chow, Y. C., and Woodbury, D. M. (1992). Effect of elevated potassium on the ion content of mouse astrocytes and neurons. *Can. J. Physiol. Pharmacol.* 70, S263–S268. doi: 10.1139/y92-271
- Zahler, R., Zhang, Z.-T., Manor, M., and Boron, W. F. (1997). Sodium kinetics of Na<sub>2</sub>K-ATPase  $\alpha$  isoforms in intact transfected HeLa cells. *J. Gen. Physiol.* 110, 201–213. doi: 10.1085/jgp.110.2.201
- Zhang, Y., Chen, K., Sloan, S. A., Bennett, M. L., Scholze, A. R., O'Keefe, S., et al. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J. Neurosci.* 34, 11929–11947. doi: 10.1523/JNEUROSCI.1860-14.2014

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Larsen, Stoica and MacAulay. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.