



## **Late Holocene changes in carbon accumulation across major ecosystem transitions of a raised bog in southern Sweden**

Kokfelt, Ulla; Rundgren, Mats

*Published in:*  
LUCCI Annual Report 2011-2012

*Publication date:*  
2012

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Kokfelt, U., & Rundgren, M. (2012). Late Holocene changes in carbon accumulation across major ecosystem transitions of a raised bog in southern Sweden. In M. Hellström, M-L. Linderson, & A. Lindroth (Eds.), *LUCCI Annual Report 2011-2012: Lund Centre for Studies of Carbon Cycle and Climate Interactions* (pp. 84). Lund.

## Late Holocene changes in carbon accumulation across major ecosystem transitions of a raised bog in southern Sweden

Ulla Kokfelt (a) and Mats Rundgren (b)

(a) Department of Geography and Geology, University of Copenhagen, Denmark

(b) Quaternary Sciences, Department of Geology, Lund University

Peatlands provide significant stores for carbon (C) derived from the atmosphere, but C-accumulation rates vary over time dependent of factors such as climate, vegetation and hydrology. Recurrence surfaces or RY's are characteristic in peat profiles from raised bogs, where peat is formed by *Sphagnum* mosses. RY's constitute distinct transitions from peat with a high degree of humification to peat with a low degree of humification. Based on the contrasting depositional environments reflected by the surfaces, RY's are traditionally interpreted to reflect rapid climatic shifts towards wetter conditions.

Twenty years after the onset of the comprehensive research on the overall Holocene stratigraphy and biogeochemistry of Store Mosse in Southern Sweden [1,2,3] we returned to the bog in order to conduct an in depth study of one selected core point in the southern part of the bog. High resolution sampling and examination of paleo-ecology and biogeochemistry in combination with an extensive dating programme across two late Holocene RY's, allowed for detailed assessment of changes in C-accumulation during the relatively rapid shifts in the ecosystem as represented by the RY's. Overall, the results support previous investigations, but our high resolution approach provides new insights to short term changes in C-accumulation during major ecosystem transitions of the bog. Hence, very different C-accumulation patterns were found across the two transitions, despite their similar appearance. This observation warrants cautiousness when estimating past changes in C accumulation across major ecosystem transitions identified in peat records.

### References

- [1] Svensson, G. (1988a) Bog development and environmental conditions as shown by the stratigraphy of Store Mosse mire in southern Sweden. *Boreas* 17, 89-111
- [2] Svensson, G. (1988b) Fossil Plant Communities and Regeneration Patterns on a Raised Bog in South Sweden. *Journal of Ecology* 76, 41-59.
- [3] Malmer, N., Svensson, G. and Wallén, B. (1997) Mass balance and nitrogen accumulation in hummocks on a South Swedish bog during the late Holocene. *Ecography* 20, 535-549.