



Trophic interactions among aphidophages in Europe

the case of the invasive ladybird *Harmonia axyridis* and the flower bug *Anthocoris nemoralis*

Howe, Andrew Gordon; Aebi, Alex; Ravn, Hans Peter

Publication date:
2010


Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Howe, A. G., Aebi, A., & Ravn, H. P. (2010). *Trophic interactions among aphidophages in Europe: the case of the invasive ladybird *Harmonia axyridis* and the flower bug *Anthocoris nemoralis**. Poster session presented at International Symposium Ecology of Aphidophaga, Perugia, Italy.

H. P. RAVN

International Symposium Ecology of Aphidophaga 11

Perugia, Italy 19 - 24 September 2010



IOBC Global



Proceedings

**TROPHIC INTERACTIONS AMONG APHIDOPHAGES IN EUROPE:
THE CASE OF THE INVASIVE LADYBIRD *HARMONIA AXYRIDIS*
AND THE FLOWER BUG *ANTHOCORIS NEMORALIS***

Howe, Andy^{1*}; Aebi, Alex² and Ravn, Hans P.¹

¹Forest & Landscape, Faculty of Life Sciences, University of Copenhagen, Frederiksberg, Denmark; ²Agroscope Reckenholz-Tänikon Research Station ART, Biodiversity and Environmental Management, Biosafety, Reckenholz, Switzerland

* andy@life.ku.dk

The establishment of *Harmonia axyridis* (released as a biocontrol agent) in North America has been linked with adverse impacts on non-target species including native ladybirds. A similar effect on native European ladybirds (Coleoptera: Coccinellidae) e.g. the two-spot ladybird *Adalia bipunctata*, could be a result of its successful, ongoing establishment. The ecological mechanisms behind these trends are believed to include resource competition and intraguild predation. Potentially, the entire aphidophagous guild of insect predators might be influenced by the presence of *H. axyridis*, e.g., ladybirds, lacewings (Neuroptera), hoverflies (Diptera: Syrphidae) and flower bugs (Heteroptera: Anthocoridae). In Denmark, when the second generation of *H. axyridis* are actively feeding, most native coccinellids are least active. Anthocorids, on the other hand, are often found feeding on aphids at this time of year and thus there potentially exists a period of several months during late summer/autumn when anthocorids and *H. axyridis* are in competition for prey. As little attention has been given to anthocorids in connection with invasive species (and in particular in connection with *H. axyridis*) this Ph.D. project is in part investigating trophic relationships between *H. axyridis* and flower bugs of the genus *Anthocoris*. Preliminary studies of intraguild predation in Petri dishes have shown third and fourth instar ladybird larvae to be consistent intraguild predators of adult *Anthocoris nemoralis*. Here we present further results from laboratory assays of intraguild predation as well as future directions based on gut detection of flower bug remains using DNA barcoding.

Key words: Intraguild predation

**EFFECTS OF *LYSIPHLE*
APHIS GLYCINES MAT**

Kaiser, Matthew C.^{1*} a

¹University of Minnesota, and Behavior, St. Paul, MN Department of Entomology * kais0101@umn.edu

Soybean aphid (*Aphis gl*) of soybeans in North America several parasitoid species imported into quarantine control agents. One such population of *L. orientalis* making it more likely to control agent. This in turn reared in quarantine, cages contain both greater number of live aphids than cages: aphid parasitoid increase advantageous for the parasitoid control. Here, we present to reveal if *L. orientalis* reproduction. Potential new work on this subject are c

Key words: soybean aphid