



Analysing biodiversity and conservation knowledge products to support regional environmental assessments

Brooks, Thomas M.; Akcakaya, H. Resit; Burgess, Neil David; Butchart, Stuart H. M.; Hilton-Taylor, Craig; Hoffmann, Michael; Juffe-Bignoli, Diego; Kingston, Naomi; MacSharry, Brian; Parr, Mike; Perianin, Laurence; Regan, Eugenie C.; Rodrigues, Ana S. L.; Rondinini, Carlo; Shennan-Farpon, Yara; Young, Bruce E.

Published in:
Scientific Data

DOI:
[10.1038/sdata.2016.7](https://doi.org/10.1038/sdata.2016.7)

Publication date:
2016

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Brooks, T. M., Akcakaya, H. R., Burgess, N. D., Butchart, S. H. M., Hilton-Taylor, C., Hoffmann, M., ... Young, B. E. (2016). Analysing biodiversity and conservation knowledge products to support regional environmental assessments. *Scientific Data*, 3, [160007]. <https://doi.org/10.1038/sdata.2016.7>

SCIENTIFIC DATA

OPEN

SUBJECT CATEGORIES

- » Biogeography
- » Conservation biology
- » Biodiversity
- » Sustainability

Analysing biodiversity and conservation knowledge products to support regional environmental assessments

Thomas M. Brooks^{1,2,3}, H. Resit Akçakaya⁴, Neil D. Burgess^{5,6}, Stuart H.M. Butchart⁷, Craig Hilton-Taylor¹, Michael Hoffmann^{1,5}, Diego Juffe-Bignoli⁵, Naomi Kingston⁵, Brian MacSharry⁵, Mike Parr⁸, Laurence Perianin¹, Eugenie C. Regan^{5,9}, Ana S.L. Rodrigues¹⁰, Carlo Rondinini¹¹, Yara Shennan-Farpon⁵ & Bruce E. Young¹²

Received: 2 October 2015

Accepted: 9 December 2015

Published: 16 February 2016

Two processes for regional environmental assessment are currently underway: the Global Environment Outlook (GEO) and Intergovernmental Platform on Biodiversity and Ecosystem Services (IPBES). Both face constraints of data, time, capacity, and resources. To support these assessments, we disaggregate three global knowledge products according to their regions and subregions. These products are: The IUCN Red List of Threatened Species, Key Biodiversity Areas (specifically Important Bird & Biodiversity Areas [IBAs], and Alliance for Zero Extinction [AZE] sites), and Protected Planet. We present fourteen Data citations: numbers of species occurring and percentages threatened; numbers of endemics and percentages threatened; downscaled Red List Indices for mammals, birds, and amphibians; numbers, mean sizes, and percentage coverages of IBAs and AZE sites; percentage coverage of land and sea by protected areas; and trends in percentages of IBAs and AZE sites wholly covered by protected areas. These data will inform the regional/subregional assessment chapters on the status of biodiversity, drivers of its decline, and institutional responses, and greatly facilitate comparability and consistency between the different regional/subregional assessments.

¹International Union for Conservation of Nature, 28 Rue Mauverney, 1196 Gland, Switzerland. ²World Agroforestry Center (ICRAF), University of the Philippines Los Baños, Laguna 4031, Philippines. ³School of Geography and Environmental Studies, University of Tasmania, Hobart, Tasmania 7001, Australia. ⁴Department of Ecology and Evolution, Stony Brook University, Stony Brook, New York 11794, USA. ⁵United Nations Environment Programme World Conservation Monitoring Centre, 219 Huntingdon Road, Cambridge CB3 0DL, UK. ⁶Center for Macroecology, Evolution and Climate, Natural History Museum, University of Copenhagen, Copenhagen DK-2100, Denmark. ⁷BirdLife International, Wellbrook Court, Cambridge CB2 3QZ, UK. ⁸American Bird Conservancy, The Plains, Virginia 20198, USA. ⁹The Biodiversity Consultancy, 3E King's Parade, Cambridge CB1 2RR, UK. ¹⁰Centre d'Ecologie Fonctionnelle et Evolutive, CNRS UMR5175, 1919 Route de Mende, 34293 Montpellier, France. ¹¹Global Mammal Assessment programme, Department of Biology and Biotechnologies, Sapienza Università di Roma, Viale dell'Università 32, 00185 Roma, Italy. ¹²NatureServe, Apdo. 358-1260, Plaza Colonial, San José, Costa Rica. Correspondence and requests for materials should be addressed to T.M.B. (email: t.brooks@iucn.org).

Design Type(s)	data integration objective • time series design • observation design
Measurement Type(s)	species extinction risk • identification of biodiverse areas • identification of protected areas
Technology Type(s)	biodiversity assessment objective • ecological observations • independent data collection method
Factor Type(s)	
Sample Characteristic(s)	habitat • South Asia • Central Europe • North Africa • South Pacific • Western Europe • South Africa • Caribbean Region • South America • Eastern Europe • Australasia • Arabian Peninsula • Mesoamerica • West Africa • Southeast Asia • East Africa • Central Africa • North America • Northeast Asia • Western Indian Ocean • Mashriq • Central Asia • Central and Western Europe • Oceania • Caribbean region • Western Asia • Antarctica • Arctic Ocean

Background & Summary

Assessment of evidence to underpin responses to challenging societal issues is increasingly recognised as one approach to bridging the science-policy interface. The Intergovernmental Panel on Climate Change is a frequently cited example¹. This is now paralleled by assessment processes on biodiversity and ecosystem services². However, while much climate change science is primarily global, biodiversity and ecosystem services are much more geographically variable³. Their assessment is therefore most useful in a multi-scale framework⁴.

Such multi-scale environmental assessment processes are now underway. The United Nations Environment Programme (UNEP) compiles a periodic GEO. The fifth edition⁵ was published in 2012, and the sixth, now underway, will incorporate regional/subregional assessments. Meanwhile, IPBES has been established, and its regional/subregional assessments are currently beginning⁶.

Global biodiversity and conservation databases hold much relevant information for these assessment processes, given the importance of consistency between different regions. Such databases are typically delivered as spatially explicit global knowledge products, and so can be analysed to inform regional/subregional assessment processes. However, both limited GIS capacity and frequently abbreviated timeframes pose severe challenges to the timely preparation of such analyses. Moreover, such regionalisation is sensitive to assumptions in data preparation and analytical settings, which if not standardised among regions can yield non-comparable results. Given the intention that the GEO and IPBES regional/subregional assessments will feed into respective global assessments, this standardisation is critically important.

Here, as a contribution towards these regional/subregional environmental assessment processes, we provide disaggregations of three scientifically robust and commonly used global knowledge products. These present assessment of the risk of species extinction and associated distributional and other information (The IUCN Red List of Threatened Species), of sites contributing significantly to the global persistence of biodiversity (Key Biodiversity Areas, specifically IBAs and AZE sites), and of protected areas (Protected Planet). We regionalise policy-relevant combinations of each according to the GEO and IPBES regions/subregions. These will inform the regional assessment chapters relating to the status of biodiversity, drivers of its decline, and institutional responses, reflected as, e.g., Chapters 3, 4, and 6 respectively in the IPBES 'Generic scoping report for the regional and subregional assessments of biodiversity and ecosystem services' (Decision IPBES-3/1: Work programme for the period 2014–2018, Annex III), and Section 2 'State of the Environment' of the GEO regional assessments in assessing the status and trends of biota and ecosystems. The generic scoping report for IPBES regional/subregional assessments (IPBES/3/6/Add.1), the rationale for which is to promote coherence across the assessments, specifies that these will draw on relevant datasets.

The variation between these regionalisations is remarkable. It provides great support to the regional assessment approach, given that this variation is masked by global reporting. We leave inference regarding causes of this variation to the assessment processes themselves, but note that contributing factors will surely include variation in underlying biogeography, intensity of anthropogenic drivers, and capacity for conservation response.

Disaggregation of other knowledge products would complement our work. Examples could include regionalisation of forest cover⁷, species populations⁸, forest carbon⁹, and protected area management effectiveness, equitability, ecological representativeness, and connectedness¹⁰. UNEP-WCMC is undertaking such regionalisation for other measures of the state of nature and drivers of change. Other emerging knowledge products, for example to assess risk of ecosystem collapse¹¹, and will also be candidates for such disaggregation once they achieve global coverage.

Finally, given the dynamic nature of all three knowledge products used here, we emphasise the importance that assessment processes build capacity for customised analysis. For future regional/subregional analyses, it will be necessary to repeat such analyses, because the data will be outdated by then. If systematic disaggregations of global datasets such as those presented here become unnecessary in the future, because capacity to undertake such analyses has by that point been developed within each region, this would be an excellent measure of the success of the capacity-building efforts associated with processes to assess biodiversity and ecosystem services. Clear guidelines will still be necessary to standardise data analysis across regions, as a precondition for comparisons in space and time.

Methods

The regionalisations slated for use by GEO¹² (updated to classify the five Central Asian countries into the 'Europe' region) and IPBES¹³ (specifically IPBES-3/1 Annexes IV–VII) are documented in Table 1 (available online only) and shown in Fig. 1. Where data are available, we also include 'Areas Beyond National Jurisdiction' (ABNJ) as a region for both GEO and IPBES, because while the high seas are not the subject of regional assessment in these current processes, they may be incorporated into GEO and IPBES in the future. We also include a region for 'Excluded' for IPBES, for completeness (this is the Antarctic, which might be included in a future regional IPBES marine assessment). Importantly, these regions do not have (and do not claim to have) any biogeographic basis. Rather they are established based on assumed policy relevance, given economic, cultural, and political similarities among their constituent countries. The three knowledge products that we disaggregate regionally and subregionally comprise assessments of species, important sites, and protected areas. All literature references were accessed 1 September 2015.

The IUCN Red List of Threatened Species

The IUCN Red List of Threatened Species¹⁴ (<http://www.iucnredlist.org>) is a knowledge product derived from assessment of species extinction risk against the IUCN Red List Categories and Criteria¹⁵. The IUCN Red List of Threatened Species dates back five decades¹⁶. Stimulated by the 1984 Road to Extinction Conference¹⁷, a process was initiated to establish quantitative categories and criteria for assessment of extinction risk¹⁸, and this standard was eventually approved in 2000 by IUCN Council¹⁹. The nine mutually-exclusive categories of extinction risk are: Not Evaluated (NE); Data Deficient (DD); Least Concern (LC); Near Threatened (NT); Vulnerable (VU); Endangered (EN); Critically Endangered (CR); Extinct in the Wild (EW); and Extinct (EX). In addition, a flag can be applied to denote Critically Endangered species which are 'Possibly Extinct' and 'Possibly Extinct in the Wild'²⁰. It incorporates robust protocols for handling uncertainty²¹ and guidelines for application at national and regional levels²². Required documentation for all assessments includes not only application of the categories and criteria, but also distribution maps, and application of standard classification schemes, e.g., for threats²³. Its application is supported by detailed, regularly updated guidelines²⁴. As a risk assessment protocol, The IUCN Red List of Threatened Species does not drive priorities for any particular type of action²⁵ but rather informs a broad scope of policy and practice ranging from threatened species legislation through to Environmental Impact Assessment^{26,27}.

The IUCN Red List of Threatened Species is a dynamic knowledge product. Version 2015-2 includes assessments of 77,340 species against the IUCN Red List Categories and Criteria¹⁹. These data are maintained in an underlying database, the Species Information Service (SIS), and are freely available for non-commercial use according to published terms²⁸, and under data licence for commercial use through IBAT²⁹.

All bird species have been assessed six times by BirdLife International³⁰, and there have been two assessments of all mammals³¹, amphibians³², and reef-building corals³³. Third reassessments are underway for mammals and amphibians, and comprehensive global assessments of reptiles and fishes are far-advanced, with the former building off existing assessments of all chameleons¹⁴, and all snakes³⁴, and the latter all sharks and rays³⁵, tarpons and ladyfishes³⁶, parrotfishes and surgeonfishes³⁷, groupers³⁸, tunas and billfishes³⁹, and hagfishes⁴⁰, as well as angelfishes, blennies, butterflyfishes, picarels, porgies, pufferfishes, seabreams, sturgeon, and wrasses¹⁴. Other animal groups that are already comprehensively assessed include freshwater caridean shrimps⁴¹, cone snails⁴², freshwater crabs⁴³, freshwater crayfish⁴⁴, and lobsters¹⁴. Among plants, comprehensive assessments are complete for cacti⁴⁵, conifers⁴⁶, cycads⁴⁷, seagrasses⁴⁸, and species occurring in mangrove ecosystems⁴⁹. In addition, a sampled approach to Red Listing⁵⁰ has been implemented for reptiles⁵¹ and dragonflies and damselflies⁵², and is being implemented for various other invertebrate⁵³ and plant^{54,55} taxa. The IUCN Red List has a target of assessing 160,000 species, stratified taxonomically, to serve as a 'barometer of life' representative across species and ecosystems⁵⁶.

The IUCN Red List of Threatened Species is governed by a Red List Committee, the Chair of which is appointed by the Chair of the IUCN Species Survival Commission (SSC; a position elected by the IUCN Membership of 216 governments and state agencies and 1,043 NGOs at the World Conservation Congress, once every four years). The Red List Committee comprises equal representation from SSC, the Global Species Programme of the IUCN Secretariat, and the Red List Partnership. The latter comprises a dozen institutions who contribute \$200,000 or more annually towards the delivery of the Red List Strategic Plan⁵⁷. A Standards & Petitions Sub-Committee, independently appointed by and accountable




Figure 1. GEO and IPBES regions and subregions. Regionalisation documenting each GEO region and subregion (a) and IPBES region and subregion (b).

to the SSC Chair, serves to adjudicate petitions against particular assessments or disputes related to the Red List Categories and Criteria.

For larger taxonomic groups that have been assessed comprehensively (i.e., for which >90% of species have been assessed), listed above, we present total numbers of species occurring in each region/subregion, for GEO (Data citation 1, Fig. 2a,b) and IPBES (Data citation 2, Fig. 3a,b). The numbers of species endemic to each region and subregion are shown separately for GEO (Data citation 3, Fig. 2c,d) and IPBES (Data citation 4, Fig. 3c,d). Extinction risk has been assessed for all species in these taxonomic groups. We therefore present in each of the Data citations 1-4 the numbers of species in each Red List Category and the overall percentage threatened (presented as lower, upper and best estimates).

For the taxonomic groups that have been assessed multiple times, it is possible to derive Red List Indices, which are indicators of the aggregate rate at which all species in a given taxonomic group are moving towards extinction^{58,59}. Critically, the derivation of the Red List Indices requires extracting only those changes in Red List category between assessments that are caused by genuine increases or decreases in extinction risk, while those caused by changing knowledge or revised taxonomy are accounted for so




Figure 2. Red List Categories of species occurring in and endemic to each GEO region and subregion. Proportion of species, by Red List Category, in comprehensively assessed groups on The IUCN Red List of Threatened Species (Version 2015-2) occurring in each GEO region (a) and subregion (b); and proportion of endemic species, by Red List Category, in comprehensively assessed groups on The IUCN Red List of Threatened Species (Version 2015-2) occurring in each GEO region (c) and subregion (d). The vertical red lines show the best estimate for the proportion of extant species considered threatened (CR, EN and VU) if Data Deficient species are Threatened in the same proportion as data-sufficient species. The numbers to the right of each bar represent the total number of species assessed and in parentheses the best estimate of the percentage threatened. CR, critically endangered; DD, data deficient; EN, endangered; EW, extinct in the wild; EX, extinct; LC, least concern; NT, near threatened; VU, vulnerable.

that they do not drive trends in the index⁶⁰. This avoids the fundamental flaw in earlier approaches to developing indicators from Red Lists^{61,62}. The Red List Index has been widely applied as a biodiversity indicator^{63–68}. Red List Indices can also be disaggregated by themes including, among others, biogeographic realm, ecosystem, taxonomy, habitat association, threats, ecosystem services, and life-history traits^{58,69–75}. For example, the 2010 Millennium Development Goals Report⁶⁶ disaggregated the Red List Index between developed and developing countries (page 57).

It is possible to downscale Red List Indices spatially by combining information on species' changes in Red List status with the range maps compiled as required documentation for each species' assessment. For any given taxon and over a given assessment period, the weighted annual change in Red List status is a measure of the relative annual contribution of each region/subregion to the overall change in the global Red List Index for that taxon. Specifically, this measure is calculated for any given region/subregion as a weighted species richness divided by the number of years in the assessment period, where each species is weighted by: i) the number of genuine category changes in The IUCN Red List of Threatened Species during the assessment period (one category change of increasing extinction risk = -1; one category change of decreasing extinction risk = +1), and ii) by the fraction of the species' range occurring within the region/subregion⁷⁶. Rodrigues *et al.*⁷⁶ successfully applied this downscaling technique to grid cells, ecological regions, and countries. We extend this downscaling to the regionalisation slated for GEO (Data citation 5) and IPBES (Data citation 6).

Key Biodiversity Areas (specifically IBAs and AZE sites)

Key Biodiversity Areas (KBAs) are sites contributing significantly to the global persistence of biodiversity⁷⁷. They are identified by assessment of sites against standard criteria for the presence of threshold levels of significance for threatened biodiversity (based on Red Lists), range-restricted biodiversity, ecological integrity, and biological process. In 2004, IUCN's Membership requested




Figure 3. Red List Categories of species occurring in and endemic to each IPBES region and subregion. Proportion of species, by Red List Category, in comprehensively assessed groups on The IUCN Red List of Threatened Species (Version 2015-2) occurring in each IPBES region (**a**) and subregion (**b**); and proportion of endemic species, by Red List Category, in comprehensively assessed groups on The IUCN Red List of Threatened Species (Version 2015-2) occurring in each IPBES region (**c**) and subregion (**d**). The vertical red lines show the best estimate for the proportion of extant species considered threatened (CR, EN and VU) if Data Deficient species are Threatened in the same proportion as data-sufficient species. The numbers to the right of each bar represent the total number of species assessed and in parentheses the best estimate of the percentage threatened. CR, critically endangered; DD, data deficient; EN, endangered; EW, extinct in the wild; EX, extinct; LC, least concern; NT, near threatened; VU, vulnerable.

‘a worldwide consultative process to agree a methodology to enable countries to identify Key Biodiversity Areas, drawing on data from the IUCN Red List of Threatened Species and other datasets, building on existing approaches’ (WCC-2004-Res-013). An initial formulation of the scientific basis for identification of KBAs was published by Eken *et al.*⁷⁸, extended into best practice guidelines by Langhammer *et al.*⁷⁹ In response to published critique^{80,81}, the World Commission on Protected Areas (WCPA) and SSC convened a joint taskforce to consolidate standards for KBA identification. Building from six technical workshops and 12 regional workshops, this taskforce released a consultation document for public comment⁷⁷ in 2014 and final revisions to this are currently underway for presentation to IUCN’s Council.

The mandate that the process for consolidation of the standard for identification of KBAs must ‘building on existing approaches’ is important, because a number of such processes have been in place for four decades. BirdLife International (then the International Council for Bird Preservation) first established such criteria for the identification of Important Bird Areas (now Important Bird & Biodiversity Areas; IBAs) in the late 1970s. National processes, led by BirdLife International partner NGOs, have now undertaken site assessment following these criteria in >200 countries and territories, yielding identification of >12,800 IBAs in total³⁰. The criteria have also been applied in the marine environment to identify >3,000 marine IBAs including 120 in Areas Beyond National Jurisdiction⁸². IBA data are freely available for non-commercial use according to published terms⁸³, and under data licence for commercial use through IBAT²⁹.

Numerous other organisations have utilised similar criteria to identify important sites for, e.g., amphibians⁸⁴, butterflies⁸⁵, plants⁸⁶, freshwater biodiversity⁸⁷, and marine turtles⁸⁸, mammals⁸⁹, and other biodiversity⁹⁰. In North America, the Natural Heritage Programs have since the 1970s utilised similar criteria to identify ‘B-ranked’ sites⁹¹. The Alliance for Zero Extinction (AZE), established in 2004 and comprising 88 biodiversity conservation NGOs, is dedicated to the identification and safeguard of all

KBAs holding effectively the entire global population of at least one Critically Endangered or Endangered species⁹². A total of 587 AZE sites have been identified, with these data freely available for non-commercial use⁹³, and available under data licence for commercial use through IBAT²⁹. Among other contributors to the identification of KBAs⁹⁴, the Critical Ecosystem Partnership Fund, which has based its 'ecosystem profiles' on KBAs for more than a decade, particularly stands out.

While all of these approaches, and thus KBAs as an umbrella standard, identify sites of importance for biodiversity, these are not necessarily important for any particular type of conservation action. Thus, they are intended to inform, but not prescribe site-level actions for practice and policy, including the establishment of protected areas at national level, and the designation of sites according to regional directives (e.g., Natura 2000 in the European Union) and international conventions (e.g., through the Ramsar Convention on Wetlands of International Importance, natural sites under the World Heritage Convention, Ecologically and Biologically Sensitive Areas in the marine realm under the CBD, etc).

While KBAs have not yet been identified globally using the criteria in the standard being developed, the mandate that the KBA standard and its application must be implemented 'building on existing approaches' allows us to have confidence that sites identified by those existing approaches which have been applied globally will be retained as KBAs of international importance under the new standard. The two such existing approaches that have been applied globally are IBAs and AZE sites⁹⁵. Here, we therefore report the total numbers, mean sizes, and percentage coverages of IBAs and AZE sites across the regionalisation proposed for GEO (Data citation 7) and IPBES (Data citation 8).

Protected Planet

Protected Planet is a knowledge product reporting on the location, status, and management of the world's protected areas, underpinned by the World Database on Protected Areas⁹⁶ (<http://www.protectedplanet.net>). It is based on a United Nations Economic and Social Council (ECOSOC) mandate dating back to 1959 for the compilation of the UN List of Protected Areas (ECOSOC Resolution 713 (XXVIII)), and implemented by IUCN-WCPA and the UNEP-World Conservation Monitoring Centre⁹⁷. Protected Planet data are freely available for non-commercial use according to published terms⁹⁸, and under data licence for commercial use through IBAT²⁹. It follows the IUCN definition of a protected area as 'A clearly defined geographical space, recognised, dedicated and managed, through legal or other effective means, to achieve the long-term conservation of nature with associated ecosystem services and cultural values' across six protected area management categories⁹⁹, where the data are available. Since 1981 it has been maintained as a database, and for the last decade it has been made available as an online knowledge product¹⁰⁰. In addition to formal, government-reported data, Protected Planet also compiles protected area data from other sources, and is in the process of strengthening its coverage across protected area governance types¹⁰¹. In the future, the WDPA is likely to be expanded to include data on 'other effective area-based conservation measures', once the definition of such sites has been agreed⁹⁷. This definition is still under discussion¹⁰², and an IUCN-WCPA task force has been created to provide recommendations on such definition.

Global summary statistics for the coverage of land and sea by different protected area categories are presented in the Protected Planet reports (Fig. 2.8 in Juffe-Bignoli *et al.*¹⁰). Here, we disaggregate the latest such statistics according to the regionalisations for GEO (Data citation 9), and IPBES (Data citation 10).

An oft-cited limitation of percentage protected area coverage and growth is that it does not account for the distribution of biodiversity which stands to benefit from site-level conservation¹⁰³. Complementing percentage area coverage, much more relevant statistics can be derived by considering the coverage by protected areas of IBAs and AZE sites (Fig. 2c in Butchart *et al.*¹⁰⁴). These indicators were incorporated into recent assessments of progress against the targets of the Convention on Biological Diversity's Strategic Plan for Biodiversity^{64,68}. Here, we therefore derive the latest regional statistics for: proportions of IBAs fully covered by protected areas, for GEO (Data citation 11, Supplementary Fig. 1) and IPBES (Data citation 12, Supplementary Fig. 2); and proportions of AZE sites fully covered by protected areas, for GEO (Data citation 13, Supplementary Fig. 3) and IPBES (Data citation 14, Supplementary Fig. 4).

Data Records

Regional and subregional species diversity and endemism

For both GEO and IPBES, total numbers of species occurring within each region/subregion, and number of species endemic to each region/subregion, are derived using the drop-down menus for countries of occurrence in the SIS, the database which underlies The IUCN Red List of Threatened Species (which are based on documented occurrence in each country), rather than through GIS analysis of the range maps. The only exception is for ABNJ, within which species occurrences are by definition not coded for national occurrence, and so for which we derive occurrence from GIS analysis of species range maps (none of the species included occur only in ABNJ). We count species as occurring if they occur in at least one of a region's countries; and species as endemic if they are not listed as occurring in any countries outside of a given region (excluding records of vagrants, records of uncertain origin, and introduced populations). We present these data for each of the following taxa: mammals; birds; seasnakes; chameleons; amphibians; sharks and rays; selected bony fish groups (angelfishes; butterflyfishes; tarpons and ladyfishes; parrotfishes and surgeonfishes; groupers; wrasses; tunas and billfishes; hagfishes; sturgeon; blennies; pufferfishes;

seabreams; porgies; and picarels); freshwater caridean shrimps; cone snails; freshwater crabs; freshwater crayfish; lobsters; reef-building corals; cacti; conifers; cycads; seagrasses; and plant species occurring in mangrove ecosystems. Resulting data can be found in Data citation 1 and Data citation 3 for GEO (Fig. 2) and in Data citation 2 and Data citation 4 for IPBES (Fig. 3).

Regional and subregional species prevalence of species extinction risk

For both GEO and IPBES, total numbers of threatened species occurring in and endemic to each region/subregion, and proportions of the total number of species and number of endemic species occurring within each region/subregion that are threatened, are derived from the documented occurrence of species in countries. The only exception is for ABNJ, within which species occurrences are by definition not coded for national occurrence, and so for which we derive occurrence from GIS analysis of species range maps (none of the species included occur only in ABNJ). We count species as occurring if they occur in at least one of a region's countries; and species as endemic if they are not listed as occurring in any countries outside of a given region (excluding records of vagrants, records of uncertain origin, and introduced populations). We present these data for each of the following taxa: mammals; birds; seasnakes; chameleons; amphibians; sharks and rays; selected bony fish groups (angelfishes and butterflyfishes; tarpons and ladyfishes; parrotfishes and surgeonfishes; groupers; wrasses; tunas and billfishes; hagfishes; sturgeon; blennies; pufferfishes; seabreams; porgies; and picarels); freshwater caridean shrimps; cone snails; freshwater crabs; freshwater crayfish; lobsters; reef-building corals; cacti; conifers; cycads; seagrasses; and plant species occurring in mangrove ecosystems.

The proportion of threatened species can be calculated for all groups that have been comprehensively assessed, but the number of threatened species is often uncertain because it is not known whether Data Deficient (DD) species are actually threatened or not. Some taxonomic groups are much better known than others (i.e., they have fewer DD species), and therefore a more accurate figure for proportion of threatened species can be calculated. Other, less well known groups have a large proportion of DD species, which brings uncertainty into the estimate for proportion of threatened species. The reported percentage of threatened species for each region and sub-region is therefore presented as a best estimate within a range of possible values bounded by lower and upper estimates:

- Lower estimate ('lower bound') = % threatened extant species, assuming all DD species are not threatened, i.e., $(CR+EN+VU)/(\text{total assessed}-EX)$
- Best estimate ('mid-point') = % threatened extant species, assuming DD species are equally threatened as data sufficient species, i.e., $(CR+EN+VU)/(\text{total assessed}-EX-DD)$
- Upper estimate ('upper bound') = % threatened extant species, assuming all DD species are threatened, i.e., $(CR+EN+VU+DD)/(\text{total assessed}-EX)$

Resulting data can be found in Data citation 1 and Data citation 3 for GEO (Fig. 2) and in Data citation 2 and Data citation 4 for IPBES (Fig. 3).

Regional and Subregional Red List Indices

For both GEO and IPBES, downscaled Red List Indices for mammals, birds, and amphibians are derived as the sum, for all species occurring in the region/subregion, of each species' number of genuine category changes on The IUCN Red List of Threatened Species (one category change of increasing extinction risk = -1; one category change of decreasing extinction risk = +1), multiplied by the proportion of the species range occurring within the region/subregion, divided by the number of years of the total assessment period⁷⁶. Category changes encompass the following categories, in order of increasing extinction risk: LC; NT; VU; EN; CR; EW and EX (including CR (Possibly Extinct) and CR (Possibly Extinct in the Wild)). Resulting data can be found in Data citation 5 for GEO and in Data citation 6 for IPBES. This approach requires an assumption that changes in extinction risk are evenly spread across all species ranges. While this will never be wholly valid, for regions as broad as those proposed for the three assessment processes in question, it will be a close approximation, because such a high proportion of species are endemic to single regions (86% for GEO, 90% for IPBES) and even to single subregions (66% for both GEO and IPBES).

Regional and subregional coverage of Key Biodiversity Areas (specifically of IBAs and AZE sites)

For both GEO and IPBES regions and subregions, we present the total numbers and mean sizes of IBAs and AZE sites, and merge site coverage into two layers (one each for IBAs and AZE sites) to calculate percentage areas of each region and subregion covered by IBAs and AZE sites respectively. Resulting data can be found in Data citation 7 for GEO and in Data citation 8 for IPBES.

Regional and subregional coverage of protected areas

For both GEO and IPBES, we present regional and subregional summary statistics for the coverage of land and sea by different protected area categories. Resulting data can be found in Data citation 9 for GEO and in Data citation 10 for IPBES.

Regional and subregional trends in coverage of Key Biodiversity Areas (specifically of IBAs and AZE sites) by protected areas

For both GEO and IPBES, we present regional and subregional summary statistics for trends in the proportions of IBAs and of AZE sites fully covered by protected areas, using data on the year of protected area establishment recorded in the January 2013 version of the World Database on Protected Areas. We overlaid digital boundaries of protected areas onto IBAs and AZEs to quantify the degree of overlap. Uncertainty in tracking changes in protected area coverage is generated by the fact that dates of establishment are not documented for 14.3% of terrestrial and 8.6% of marine protected areas. We reflected this by assigning dates of establishment 1,000 times selected at random from those for dated protected areas in the same country to these un-dated sites (or, for countries with less than five protected areas with known year of establishment, from all terrestrial or marine protected areas), and plotting the 95% confidence intervals around median protected area coverage accordingly^{64,95,104}. Resulting data can be found in Data citation 11 for GEO for IBAs (Supplementary Fig. 1), in Data citation 13 for GEO for AZE (Supplementary Fig. 3), in Data citation 12 for IPBES for IBAs (Supplementary Fig. 2), and in Data citation 14 for IPBES for AZE (Supplementary Fig. 4). No AZE sites have yet been identified in the GEO Polar region or subregions, the GEO and IPBES West Asia region and subregions, or in the GEO and IPBES Central Asia subregions, and so these are excluded from Data citation 13 and Data citation 14, and from Supplementary Figs 3 and 4, accordingly. No IBAs or AZE sites are covered by protected areas in ABNJ, and so ABNJ are also excluded from Data citation 11, Data citation 12, Data citation 13, Data citation 14 and Supplementary Fig. 1.

Technical Validation

The IUCN Red List of Threatened Species

The primary technical validation of the IUCN Red List Categories and Criteria undertaken to date compared actual movement of threatened species through the Red List Categories with that predicted assuming equivalence of the E Criterion (formal quantitative analysis of extinction risk, e.g., using Population Viability Analysis¹⁰⁵) to the other four criteria¹⁰⁶. They examined all bird species globally for 1988–2004 and all Australian species for the much longer period, 1750–2000. They found that the expected rates associated with the thresholds for extinction risk used under the E Criterion were broadly consistent with the observed rates, with the exception of category change from Critically Endangered to Extinct, for which the data revealed many fewer extinctions than predicted from the E Criterion thresholds. They speculated that this mismatch is explained by the disproportionate focus of conservation action on Critically Endangered species, and indeed there is good evidence that conservation efforts have prevented the extinction of a substantial number of Critically Endangered species^{107,108}. Other approaches to validating the IUCN Red List Categories and Criteria involved retrospective testing based on a reconstructed past extinction¹⁰⁹, and prospective testing based on projected future extinctions due to climate change^{110,111}. Both types of studies concluded that the IUCN criteria can not only identify species that would be extinct without conservation effort, but do so with substantial warning time.

In addition, a number of studies have conducted inter-model comparisons between the IUCN Red List Categories and Criteria and other protocols for assessment of extinction risk^{112–114}, comparisons of the consistency of assessments within each of these¹¹⁵, and comparisons of the results of application of the IUCN Red List Categories and Criteria with other methods for predicting extinctions such as species-area curves¹¹⁶ and random forest decision trees of traits¹¹⁷. These found relatively high levels of agreement between the protocols.

Finally, estimates of uncertainty around measurement of threat prevalence (Data citation 1, Data citation 2, Data citation 3, Data citation 4, Figs 2 and 3) can be derived from consideration of the numbers of species classified in the Data Deficient category, wherein a species is assessed as having insufficient data available to apply the IUCN Red List Criteria⁶⁴.

Key Biodiversity Areas (specifically IBAs and AZE sites)

Di Marco *et al.*¹¹⁸ undertook a major validation exercise for KBAs, specifically for IBAs in Australia, Europe, and Southern Africa, the three regions from which major independent datasets are available in the form of comprehensive bird atlases. The study set a quantitative conservation targets for all birds in the regions (in terms of number of atlas grid cells to be protected, and compared the selection frequency of atlas grid cells covering IBAs, selected through application of a simulated annealing software (Marxan¹¹⁹), with the selection frequency of cells outside IBAs. It found the former to be much higher than the latter, as predicted if such threshold approaches to identification of important sites genuinely identify sites of high irreplaceability for the global persistence of biodiversity. Montesino Pouzols *et al.*¹²⁰ undertook a similar validation, comparing KBAs identified in Madagascar, Myanmar, and the Philippines to results of their global analysis of important areas for terrestrial vertebrates, finding similarly consistent results.

Several tests have also been undertaken to examine the degree to which IBAs represent important sites for biodiversity more generally^{121–123}, finding that representation of threatened species in other taxonomic groups is high. Equivalent tests have not yet been undertaken for marine or freshwater biodiversity. Finally, provision also exists for documentation of KBAs as ‘candidate sites’ suspected by experts as holding threshold levels of threatened species, but for taxa that have not yet been formally

File name	Description	Authors	Dryad DOI
Total_Species_GEO.csv	Total numbers of species and of threatened species occurring in each GEO region and subregion	IUCN	http://dx.doi.org/10.5061/dryad.6gb90.2/1.2
Total_Species_IPBES.csv	Total numbers of species and of threatened species occurring in each IPBES region and subregion		http://dx.doi.org/10.5061/dryad.6gb90.2/2.2
Endemic_Species_GEO.csv	Total numbers of species and of threatened species endemic to each GEO region and subregion		http://dx.doi.org/10.5061/dryad.6gb90.2/3.2
Endemic_Species_IPBES.csv	Total numbers of species and of threatened species endemic to each IPBES region and subregion		http://dx.doi.org/10.5061/dryad.6gb90.2/4.2
Red_List_Index_GEO.csv	Relative annual contribution to the global Red List Index for mammals, birds, and amphibians in each GEO region and subregion	IUCN & BirdLife International	http://dx.doi.org/10.5061/dryad.6gb90.2/5.2
Red_List_Index_IPBES.csv	Relative annual contribution to the global Red List Index for mammals, birds, and amphibians in each IPBES region and subregion		http://dx.doi.org/10.5061/dryad.6gb90.2/6.2
IBAs_AZEs_GEO.csv	Total numbers, mean sizes, and percentage coverages of IBAs and AZEs in each GEO region and subregion	BirdLife International & AZE	http://dx.doi.org/10.5061/dryad.6gb90.2/7.2
IBAs_AZEs_IPBES.csv	Total numbers, mean sizes, and percentage coverages of IBAs and AZEs in each IPBES region and subregion		http://dx.doi.org/10.5061/dryad.6gb90.2/8.2
PAs_GEO.csv	Percentage protected area coverage of land and sea, for each GEO region and subregion	UNEP-WCMC & IUCN	http://dx.doi.org/10.5061/dryad.6gb90.2/9.2
PAs_IPBES.csv	Percentage protected area coverage of land and sea, for each IPBES region and subregion		http://dx.doi.org/10.5061/dryad.6gb90.2/10.2
Protected_IBAs_GEO.csv	Percentage of IBAs wholly covered by protected areas, over time, for each GEO region and subregion	BirdLife International, IUCN & UNEP-WCMC	http://dx.doi.org/10.5061/dryad.6gb90.2/11.2
Protected_IBAs_IPBES.csv	Percentage of IBAs wholly covered by protected areas, over time, for each IPBES region and subregion		http://dx.doi.org/10.5061/dryad.6gb90.2/12.2
Protected_AZEs_GEO.csv	Percentage of AZE sites wholly covered by protected areas, over time, for each GEO region and subregion	AZE, BirdLife International, IUCN & UNEP-WCMC	http://dx.doi.org/10.5061/dryad.6gb90.2/13.2
Protected_AZEs_IPBES.csv	Percentage of AZE sites wholly covered by protected areas, over time, for each IPBES region and subregion		http://dx.doi.org/10.5061/dryad.6gb90.2/14.2

Table 2. Individual datasets published in the data citations.

assessed for the global IUCN Red List or which have not yet been formally documented to occur at the site. No aggregated analyses have yet been undertaken of these candidate sites.

Protected Planet

The compilation of the World Database on Protected Areas is underpinned by a formal ECOSOC mandate for national submission of protected area datasets to compile the UN List of Protected Areas (ECOSOC Resolution 713 (XXVIII)), technical validation of the data focuses on tracking changes in the completeness of the dataset and working closely with data providers to ensure the veracity and quality of the information provided through a clearly predefined working protocol⁹⁷. The data are collected directly from governmental agencies while data gaps are provided by other authoritative sources such as NGOs or secretariats of international conventions (e.g., Ramsar Convention) or regional entities (e.g., European Environment Agency). All data providers are requested to sign a Data Contributor Agreement which ensures the data published in the WDPA have been approved by the data provider and comply with the terms and conditions of use of the original data.

The quality of the WDPA is ensured by the WDPA data standards⁹⁷ implemented in 2010. As a result, the WDPA has experienced a great improvement in quality in the past years. For example, the proportion of protected areas represented in the dataset as mapped polygons, as opposed to latitude-longitude centre points alone. Between 2003 and 2015, this proportion increased from 39.5 to 91.1%.

Usage Notes

For the numbers of species and of endemic species in each of the GEO and IPBES regions and subregions (Data citations 1–4), we recommend presentation following the horizontal proportional bar chart used in Figs 2 and 3. For relative annual contribution to the global Red List Index (Data citation 5, Data citation 6), coverage of KBAs, specifically IBAs and AZEs (Data citation 7, Data citation 8), and protected area coverage (Data citation 9, Data citation 10), the most appropriate presentation would be in tables, maybe supplemented with absolute bar chart. Changing protected area coverage of KBAs, specifically IBAs and AZEs, over time (Data citations 11–14) is best presented as line graphs, following those presented in Supplementary Fig. 1. In each case, it would be suitable to structure presentation for the regional assessments to show the component subregions. By extension, for the global assessments, it would be suitable to structure them to show the component regions. Table 2 documents the individual datasets published in the data citations.

References

- Bolin, B. *A History of the Science and Politics of Climate Change: the Role of the Intergovernmental Panel on Climate Change* (Cambridge University Press: Cambridge, UK, 2007).
- Díaz, S., Demissew, S., Joly, C., Lonsdale, W. M. & Larigauderie, A. A rosetta stone for nature's benefits to people. *PLoS Biol.* **13**, e1002040 (2015).
- Brooks, T. M., Lamoreux, J. F. & Soberón, J. IPBES≠IPCC. *Trends Ecol. Evol.* **29**, 543–545 (2014).

4. Soberón, J. M. & Sarukhan, J. K. A new mechanism for science-policy transfer and biodiversity governance? *Environ. Conserv.* **36**, 265–267 (2009).
5. UNEP. *Global Environmental Outlook 5: Environment for the Future we Want*. Available at <http://www.unep.org/geo/geo5.asp> (UNEP: Nairobi, Kenya, 2012).
6. Opgenoorth, L. & Faith, D. P. The Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES), up and walking. *Front. Biogeogr.* **5**, 207–211 (2014).
7. Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change. *Science* **342**, 850–853 (2013).
8. Loh, J. *et al.* The Living Planet Index: using species population time series to track trends in biodiversity. *Philos. T. Roy. Soc. B* **360**, 289–295 (2005).
9. Ruesch, A. S. & Gibbs, H. K. *New Global Biomass Carbon Map for the Year 2000 Based On IPCC Tier-1 Methodology*. Available at http://cdiac.ornl.gov/epubs/ndp/global_carbon/carbon_documentation.html (Oak Ridge National Laboratory: Oak Ridge, USA, 2008).
10. Juffe-Bignoli, D. *et al.* *Protected Planet Report 2014*. Available at <https://portals.iucn.org/library/node/44896> (UNEP-WCMC: Cambridge, UK, 2014).
11. Keith, D. A. *et al.* Scientific foundations for an IUCN Red List of ecosystems. *PLoS ONE* **8**, e62111 (2013).
12. UNEP. *GEO-4 Subregional Breakdown*. Available at http://geodata.grid.unep.ch/extras/geo_breakdown.doc (UNEP: Nairobi, Kenya, 2007).
13. IPBES. *Report of the Third Session of the Plenary of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES-3/1*. Available at <http://www.ipbes.net/index.php/plenary/ipbes-3#one> (IPBES: Bonn, Germany, 2015).
14. IUCN. *IUCN Red List of Threatened Species. Version 2015-2*. Available at <http://www.iucnredlist.org> (IUCN: Gland, Switzerland, 2015).
15. Mace, G. M. *et al.* Quantification of extinction risk: IUCN's system for classifying threatened species. *Conserv. Biol.* **22**, 1424–1442 (2008).
16. Smart, J., Hilton-Taylor, C. & Mittermeier, R. A. *The IUCN Red List: 50 Years of Conservation*. Available at <https://portals.iucn.org/library/node/44968> (CEMEX: Mexico City, Mexico, 2014).
17. Fitter, R. & Fitter, M. *The Road to Extinction: Problems with Categorizing the Status of Taxa Threatened with Extinction*. Available at <https://portals.iucn.org/library/node/5867> (IUCN: Gland, Switzerland, 1987).
18. Mace, G. & Lande, R. Assessing extinction threats: toward a reevaluation of IUCN threatened species categories. *Conserv. Biol.* **5**, 148–157 (1991).
19. IUCN. *IUCN Red List Categories and Criteria: Version 3.1*. Available at <https://portals.iucn.org/library/node/7977> (IUCN Species Survival Commission, IUCN: Gland, Switzerland, 2001).
20. Butchart, S. H. M., Stattersfield, A. J. & Brooks, T. M. Going or gone: defining 'Possibly Extinct' species to give a truer picture of recent extinctions. *Bull. Brit. Ornithol. Club* suppl. **126**, 7–24 (2006).
21. Akçakaya, H. R. *et al.* Making consistent IUCN classifications under uncertainty. *Conserv. Biol.* **14**, 1001–1013 (2000).
22. Gärdenfors, U., Hilton-Taylor, C., Mace, G. M. & Rodriguez, J. P. The application of IUCN Red List criteria at regional levels. *Conserv. Biol.* **15**, 1206–1212 (2001).
23. Salafsky, N. *et al.* A standard lexicon for biodiversity conservation: unified classifications of threats and actions. *Conserv. Biol.* **22**, 897–911 (2008).
24. IUCN Standards and Petitions Subcommittee. *Guidelines for Using the IUCN Red List Categories and Criteria. Version 11*. Available at <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> (Standards and Petitions Subcommittee, Species Survival Commission, IUCN: Gland, Switzerland, 2014).
25. Lamoreux, J. *et al.* Value of the IUCN Red List. *Trends Ecol. Evol.* **18**, 214–215 (2003).
26. Rodrigues, A. S. L., Pilgrim, J. D., Lamoreux, J. F., Hoffmann, M. & Brooks, T. M. The value of the IUCN Red List for conservation. *Trends Ecol. Evol.* **21**, 71–76 (2006).
27. Lacher, T. E. Jr, Boitani, L. & da Fonseca, G. A. B. The IUCN global assessments: partnerships, collaboration and data sharing for biodiversity science and policy. *Conserv. Lett.* **5**, 327–333 (2012).
28. IUCN. *The IUCN Red List Terms and Conditions of Use (version 2.1)*. Available at <http://www.iucnredlist.org/info/terms-of-use> (IUCN: Gland, Switzerland, 2015).
29. IBAT. *Integrated Biodiversity Assessment Tool*. Available at <https://www.ibatforbusiness.org/> (IBAT: Washington DC, USA, 2015).
30. BirdLife International. *DataZone*. Available at <http://www.birdlife.org/datazone/site/search> (BirdLife International: Cambridge, UK, 2015).
31. Schipper, J. *et al.* The status of the world's land and marine mammals: diversity, threat, and knowledge. *Science* **322**, 225–230 (2008).
32. Stuart, S. N. *et al.* Status and trends of amphibian declines and extinctions worldwide. *Science* **306**, 1783–1786 (2004).
33. Carpenter, K. E. *et al.* One-third of reef-building corals face elevated extinction risk from climate change and local impacts. *Science* **321**, 560–563 (2008).
34. Elfes, C. T. *et al.* Fascinating and forgotten: the conservation status of the world's sea snakes. *Herpetol. Conserv. Biol.* **8**, 37–52 (2013).
35. Dulvy, N. K. *et al.* Extinction risk and conservation of the world's sharks and rays. *eLife* **3**, e00590 (2014).
36. Adams, A. J. *et al.* Global conservation status and research needs for tarpons (Megalopidae), ladyfishes (Elopidae) and bonefishes (Albulidae). *Fish Fish.* **8**, 280–311 (2013).
37. Comeros-Raynal, M. T. *et al.* The likelihood of extinction of iconic and dominant herbivores and detritivores of coral reefs: the parrotfishes and surgeonfishes. *PLoS ONE* **7**, e39825 (2012).
38. Sadovy de Mitcheson, Y. S. *et al.* Fishing groupers towards extinction: a global assessment of threats and extinction risks in a billion dollar fishery. *Fish Fish.* **14**, 119–136 (2012).
39. Collette, B. B. *et al.* High value and long-lived: a double jeopardy for threatened tunas and billfishes. *Science* **333**, 291–292 (2011).
40. Knapp, L. *et al.* Conservation status of the world's hagfish species and the loss of phylogenetic diversity and ecosystem function. *Aquat. Conserv.* **21**, 401–411 (2011).
41. De Grave, S. *et al.* Dead shrimp blues: a global assessment of extinction risk in freshwater shrimps (Crustacea: Decapoda: Caridea). *PLoS ONE* **10**, e0120198 (2015).
42. Peters, H., O'Leary, B. C., Hawkins, J. P., Carpenter, K. E. & Roberts, C. M. *Conus*: first comprehensive conservation Red List assessment of a marine gastropod mollusc genus. *PLoS ONE* **8**, e83353 (2013).
43. Cumberlidge, N. *et al.* Freshwater crabs and the biodiversity crisis: importance, threats, status, and conservation challenges. *Biol. Conserv.* **142**, 1665–1673 (2009).
44. Richman, N. I. *et al.* Multiple drivers of decline in the global status of freshwater crayfish (Decapoda: Astacidea). *Philos. T. Roy. Soc. B* **370**, 20140060 (2015).
45. Goettsch, B. *et al.* High proportion of cactus species threatened with extinction. *Nature Plants* **370**, doi:10.1038/NPLANTS.2015.142 (2015).

46. Farjon, A. & Page, C.N. *Conifers. Status Survey and Conservation Action Plan*. Available at <https://portals.iucn.org/library/node/7565> (IUCN/SSC Conifer Specialist Group, IUCN: Gland, Switzerland, 1999).
47. Donaldson, J. *Cycads. Status Survey and Conservation Action Plan*. Available at <https://portals.iucn.org/library/node/8203> (IUCN/SSC Cycad Specialist Group, IUCN: Gland, Switzerland, 2003).
48. Short, F. T. *et al.* Extinction risk assessment of the world's seagrass species. *Biol. Conserv.* **144**, 1961–1971 (2011).
49. Polidoro, B. A. *et al.* The loss of species: mangrove extinction risk and geographic areas of global concern. *PLoS ONE* **5**, e10095 (2010).
50. Baillie, J. E. M. *et al.* Toward monitoring global biodiversity. *Conservation Lett.* **1**, 18–26 (2008).
51. Böhm, M. *et al.* The conservation status of the world's reptiles. *Biol. Conserv.* **157**, 372–385 (2013).
52. Clausnitzer, V. *et al.* Odonata enter the biodiversity crisis debate: the first global assessment of an insect group. *Biol. Conserv.* **142**, 1864–1869 (2009).
53. Lewis, O. T. & Senior, M. J. M. Assessing conservation status and trends for the world's butterflies: the Sampled Red List Index approach. *J. Insect Conserv.* **15**, 121–128 (2011).
54. Brummitt, N. *et al.* The Sampled Red List Index for Plants, phase II: ground-truthing specimen-based conservation assessments. *Philos. T. Roy. Soc. B* **370**, 20140015 (2015).
55. Brummitt, N. *et al.* Green Plants in the Red: A Baseline Global Assessment for the IUCN Sampled Red List Index for Plants. *PLoS ONE* **10**, e0135152 (2015).
56. Stuart, S. N., Wilson, E. O., McNeely, J. A., Mittermeier, R. A. & Rodríguez, J. P. The barometer of life. *Science* **328**, 177–177 (2010).
57. IUCN. *The IUCN Red List of Threatened Species Strategic Plan 2013–2020. Version 1.0*. Available at http://cmsdata.iucn.org/downloads/red_list_strategic_plan_2013_2020.pdf (IUCN: Gland, Switzerland, 2013).
58. Butchart, S. H. M. *et al.* Measuring global trends in the status of biodiversity: Red List indices for birds. *PLoS Biology* **2**, e383 (2004).
59. Butchart, S. H. M. *et al.* Improvements to the Red List Index. *PLoS ONE* **2**, e140 (2007).
60. Brooks, T. & Kennedy, E. Biodiversity barometers. *Nature* **431**, 1046–1047 (2004).
61. Cuarón, A. D. Extinction rate estimates. *Nature* **366**, 118 (1993).
62. Burgman, M. A. Turner review no 5: are listed threatened plant species actually at risk? *Aust. J. Bot.* **50**, 1–13 (2002).
63. Butchart, S. H. M. *et al.* Using Red List indices to measure progress towards the 2010 target and beyond. *Philos. T. Roy. Soc. B* **360**, 359–372 (2005).
64. Butchart, S. H. M. *et al.* Global biodiversity: indicators of recent declines. *Science* **328**, 1164–1168 (2010).
65. Hoffmann, M. *et al.* The impact of conservation on the status of the world's vertebrates. *Science* **330**, 1503–1509 (2010).
66. UN. *The Millennium Development Goals Report 2010*. Available at <http://www.un.org/millenniumgoals/pdf/MDG%20Report%202010%20En%20r15%20-low%20res%2020100615%20-.pdf> (United Nations: New York, USA, 2010).
67. SCBD. *Global Biodiversity Outlook 4: A Mid-Term Assessment of Progress towards the Implementation of the Strategic Plan for Biodiversity 2011–2020*. Available at <https://www.cbd.int/gbo/gbo4/publication/gbo4-en.pdf> (SCBD: Montréal, Canada, 2014).
68. Tittensor, D. P. *et al.* A mid-term analysis of progress toward international biodiversity targets. *Science* **346**, 241–244 (2014).
69. Butchart, S. H. M. Red List Indices to measure the sustainability of species use and impacts of invasive alien species. *Bird Conserv. Int.* **18**(suppl.): 245–262 (2008).
70. Stuart, S. N. *et al.* *Threatened Amphibians of the World* (Lynx Edicions: Barcelona, Spain, 2008).
71. Kirby, J. S. *et al.* Key conservation issues for migratory land- and waterbird species on the world's major flyways. *Bird Conserv. Int.* **18**(suppl.): 49–73 (2008).
72. McGeoch, M. A. *et al.* Global indicators of biological invasion: species numbers, biodiversity impact and policy responses. *Divers. Distrib.* **16**, 95–108 (2010).
73. Hoffmann, M. *et al.* The changing fates of the world's mammals. *Philos. T. Roy. Soc. B* **366**, 2598–2610 (2011).
74. Croxall, J. P. *et al.* Seabird conservation status, threats and priority actions: a global assessment. *Bird Conserv. Int.* **22**, 1–34 (2012).
75. Regan, E. C. *et al.* Global trends in the status of bird and mammal pollinators. *Conserv. Lett.* **8**, 397–403 (2016).
76. Rodrigues, A. S. L. *et al.* Spatially explicit trends in the global conservation status of vertebrates. *PLoS ONE* **9**, e113934 (2014).
77. IUCN. *Consultation Document on an IUCN Standard for the Identification of Key Biodiversity Areas*. Available at https://portals.iucn.org/union/sites/union/files/doc/consultation_document_iucn_kba_standard_01oct2014.pdf (IUCN, Gland: Switzerland, 2014).
78. Eken, G. *et al.* Key biodiversity areas as site conservation targets. *BioScience* **54**, 1110–1118 (2004).
79. Langhammer, P. F. *et al.* Identification and Gap Analysis of Key Biodiversity Areas: Targets for Comprehensive Protected Area Systems. *IUCN World Commission on Protected Areas Best Practice Protected Area Guidelines Series No. 15*. Available at <https://portals.iucn.org/library/node/9055> (IUCN: Gland, Switzerland, 2007).
80. Knight, A. T. *et al.* Improving the Key Biodiversity Areas approach for effective conservation planning. *BioScience* **57**, 256–261 (2007).
81. Bennun, L., Bakarr, M., Eken, G. & da Fonseca, G. A. B. Clarifying the Key Biodiversity Areas approach. *BioScience* **207**, 645 (2007).
82. BirdLife International. *Marine IBA e-Atlas: Delivering Site Networks for Seabird Conservation*. Available at <http://maps.birdlife.org/marineIBAs/default.html> (BirdLife International: Cambridge, UK, 2012).
83. BirdLife International. *Terms of Use*. Available at <http://www.birdlife.org/datazone/info/termsofuse> (BirdLife International: Cambridge, UK, 2015b).
84. Silvano, D., Angulo, A., Carnaval, A. C. O. Q. & Pethiyagoda, R. in *Amphibian Conservation Action Plan* (eds Gascon C. *et al.*). Available at <https://portals.iucn.org/library/node/9071>. 12–15 (IUCN/SSC Amphibian Specialist Group: Gland, Switzerland, 2007).
85. van Swaay, C. A. M. & Warren, M. S. Prime butterfly areas in Europe: an initial selection of priority sites for conservation. *J. Insect Conserv.* **10**, 5–11 (2006).
86. Plantlife International. *Identifying and Protecting the World's Most Important Plant Areas*. Available at http://www.plantlife.org.uk/uploads/documents/Guide_to_Implementing_IPAs_2004.pdf (Plantlife International: Salisbury, UK, 2004).
87. Holland, R. A., Darwall, W. R. T. & Smith, K. G. Conservation priorities for freshwater biodiversity: the key biodiversity area approach refined and tested for continental Africa. *Biol. Conserv.* **148**, 167–179 (2012).
88. Bass, D., Anderson, P. & De Silva, N. Applying thresholds to identify key biodiversity areas for marine turtles in Melanesia. *Anim. Conserv.* **14**, 1–11 (2011).
89. Corrigan, C. M. *et al.* Developing important marine mammal area criteria: learning from ecologically or biologically significant areas and key biodiversity areas. *Aquat. Conserv.* **24**(suppl.): 166–183 (2014).
90. Edgar, G. J. *et al.* Key Biodiversity Areas as globally significant target sites for the conservation of marine biological diversity. *Aquat. Conserv.* **18**, 969–983 (2008).
91. Jenkins, R. E. in *Biodiversity* (ed. Wilson E. O.) 231–239 (National Academy Press: Washington DC, USA, 1988).

92. Ricketts, T. H. *et al.* Pinpointing and preventing imminent extinctions. *Proc. Natl. Acad. Sci. USA* **102**, 18497–18501 (2005).
93. AZE. *2010 AZE Update*. Available at <http://www.zeroextinction.org> (AZE: Washington DC, USA, 2010).
94. Foster, M. N. *et al.* The identification of sites of biodiversity conservation significance: progress with the application of a global standard. *J. Threatened Taxa* **4**, 2733–2744 (2012).
95. Butchart, S. H. M. *et al.* Protecting important sites for biodiversity contributes to meeting global conservation targets. *PLoS ONE* **7**, e32529 (2012).
96. IUCN & UNEP-WCMC. *The World Database on Protected Areas (WDPA). February 2015*. Available at <http://www.protectedplanet.net> (UNEP-WCMC: Cambridge, UK, 2015).
97. UNEP-WCMC. *World Database on Protected Areas User Manual 1.1*. Available at http://wcmc.io/WDPA_Manual (UNEP-WCMC: Cambridge, UK, 2015).
98. UNEP-WCMC. *Terms of Use*. Available at <http://www.protectedplanet.net/terms> (UNEP-WCMC: Cambridge, UK, 2015).
99. Dudley, N. *Guidelines for Applying Protected Area Management Categories*. Available at <https://portals.iucn.org/library/node/9243> (IUCN: Gland, Switzerland, 2008).
100. Bertzky, B. *et al.* *Protected Planet Report 2012: Tracking Progress towards Global Targets for Protected Areas*. Available at <https://portals.iucn.org/library/node/10233> (UNEP-WCMC: Cambridge, UK, 2012).
101. Borriini-Feyerabend, G. *et al.* *Governance of Protected Areas: From Understanding to Action. IUCN Best Practice Protected Area Guidelines Series No. 20*. Available at <https://portals.iucn.org/library/node/29138> (IUCN: Gland, Switzerland, 2013).
102. Jonas, H. D., Barbuto, V., Jonas, H., Kothari, A. & Nelson, F. New steps of change: looking beyond protected areas to consider other effective area-based conservation measures. *Parks* **20**, 111–128 (2014).
103. Rodrigues, A. S. L. *et al.* Effectiveness of the global protected area network in representing species diversity. *Nature* **428**, 640–643 (2004).
104. Butchart, S. H. M. *et al.* Shortfalls and solutions for meeting national and global conservation area targets. *Conserv. Lett.* **8**, 329–337 (2015).
105. Brook, B. W. *et al.* Predictive accuracy of population viability analysis in conservation biology. *Nature* **404**, 385–387 (2000).
106. Brooke, M. L. *et al.* Rates of movement of threatened bird species between IUCN Red List categories and toward extinction. *Conserv. Biol.* **22**, 417–427 (2008).
107. Butchart, S. H. M., Stattersfield, A. J. & Collar, N. J. How many bird extinctions have we prevented? *Oryx* **40**, 266–278 (2006).
108. Rodrigues, A. S. L. Are global conservation efforts successful? *Science* **313**, 1051–1052 (2006).
109. Stanton, J. C. Present-day risk assessment would have predicted the extinction of the passenger pigeon (*Ectopistes migratorius*). *Biol. Conserv.* **180**, 11–20 (2014).
110. Keith, D. A. *et al.* Detecting extinction risk from climate change by IUCN Red List criteria. *Conserv. Biol.* **28**, 810–819 (2014).
111. Stanton, J. C., Shoemaker, K. T., Pearson, R. G. & Akçakaya, H. R. Warning times for species extinctions due to climate change. *Glob. Change Biol.* **21**, 1066–1077 (2015).
112. Keith, D. A. *et al.* Protocols for listing threatened species can forecast extinction. *Ecol. Lett.* **7**, 1101–1108 (2004).
113. O’Grady, J. J. *et al.* Correlations among extinction risks assessed by different threatened species categorization systems. *Conserv. Biol.* **18**, 1624–1635 (2004).
114. De Grammont, P. C. & Cuarón, A. D. An evaluation of threatened species categorization systems used on the American continent. *Conserv. Biol.* **20**, 14–27 (2006).
115. Regan, T. J. *et al.* The consistency of extinction risk classification protocols. *Conserv. Biol.* **19**, 1969–1977 (2005).
116. Brooks, T. M. *et al.* Habitat loss and extinction in the hotspots of biodiversity. *Conserv. Biol.* **16**, 909–923 (2002).
117. Davidson, A. D., Hamilton, M. J., Boyer, A. G., Brown, J. H. & Ceballos, G. Multiple ecological pathways to extinction in mammals. *Proc. Natl. Acad. Sci. USA* **106**, 10702–10705 (2009).
118. Di Marco, M. *et al.* Quantifying the relative irreplaceability of Important Bird and Biodiversity Areas. *Conserv. Biol.* <http://onlinelibrary.wiley.com/doi/10.1111/cobi.12609/abstract> (2016).
119. Ball, I. R. & Possingham, H. P. *Marxan (V1.8.2): Marine Reserve Design Using Spatially Explicit Annealing, a Manual prepared for The Great Barrier Reef Marine Park Authority*. Available at http://www.uq.edu.au/marxan/docs/marxan_manual_1_8_2.pdf (GBRMPA: Brisbane, Australia, 2000).
120. Montesino Pouzols, F. *et al.* Global protected area expansion is compromised by projected land-use and parochialism. *Nature* **516**, 383–386 (2014).
121. Brooks, T. *et al.* Conservation priorities for birds and biodiversity: do East African Important Bird Areas represent species diversity in other terrestrial vertebrate groups? *Ostrich suppl.* **15**, 3–12 (2001).
122. Pain, D. J., Fishpool, L., Byaruhanga, A., Arinaitwe, J. & Balmford, A. Biodiversity representation in Uganda’s forest IBAs. *Biol. Conserv.* **125**, 133–138 (2005).
123. BirdLife International. *Important Bird and Biodiversity Areas: a Global Network for Conserving Nature and Benefiting People*. Available at <http://www.birdlife.org/datazone/sowb/sowbpubs#IBA> (BirdLife International, Cambridge: UK, 2014).

Data Citations

1. IUCN. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/1.2> (2015).
2. IUCN. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/2.2> (2015).
3. IUCN. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/3.2> (2015).
4. IUCN. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/4.2> (2015).
5. IUCN & BirdLife International. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/5.2> (2015).
6. IUCN & BirdLife International. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/6.2> (2015).
7. BirdLife International & AZE. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/7.2> (2015).
8. BirdLife International & AZE. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/8.2> (2015).
9. UNEP-WCMC & IUCN. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/9.2> (2015).
10. UNEP-WCMC & IUCN. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/10.2> (2015).
11. BirdLife International, IUCN & UNEP-WCMC. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/11.2> (2015).
12. BirdLife International, IUCN & UNEP-WCMC. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/12.2> (2015).
13. AZE, BirdLife International, IUCN & UNEP-WCMC. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/13.2> (2015).
14. AZE, BirdLife International, IUCN & UNEP-WCMC. *Dryad* <http://dx.doi.org/10.5061/dryad.6gb90.2/14.2> (2015).

Acknowledgements

We thank the many thousands of individuals and organisations who contribute towards the compilation and maintenance of the three knowledge products upon which our analyses are based, as well as J. Harrison, A. Hufton, V. Khodiyar, C. Lortie, J. Watson, and an anonymous reviewer. IUCN thanks the French Ministry of Ecology and the Gordon and Betty Moore Foundation for supporting IUCN’s

engagement with IPBES. NatureServe acknowledges support from the National Science Foundation's Dimensions of Biodiversity program (award 1136586). UNEP DEWA contributed funding to undertake the regionalisation of the protected area data. T.M.B. had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Author Contributions

T.M.B. conceived, drafted, and revised the manuscript, and led the development of Table 1 (available online only). R.A. commented on the draft manuscript. N.D.B. contributed towards the development of Table 1 (available online only), of Data citations 9–14, and of Supplementary Figs 1–4, and commented on the draft manuscript. S.H.M.B. led the development of Data citations 7–8 and Data citations 11–14 and of Supplementary Figs 1–4, and commented on the draft manuscript. C.H.-T. led the development of Data citations 1–4 and of Figs 2–3, and commented on the draft manuscript. M.H. contributed towards the development of Data citations 1–4 and of Figs 2–3, and commented on the draft manuscript. D.J.-B. led the development of Data citations 9–10, contributed towards the development of Fig. 1, of Data citations 11–14, and of Supplementary Figs 1–4, and commented on the draft manuscript. N.K. contributed towards the development of Data citations 9–14 and of Supplementary Figs 1–4, and commented on the draft manuscript. B.M.S. contributed towards the development of Table 1 (available online only), of Data citations 9–10, Data citations 11–14, and of Supplementary Figs 1–4, and commented on the draft manuscript. M.P. commented on the draft manuscript. L.P. commented on the draft manuscript. E.R. commented on the draft manuscript. A.S.L.R. led the development of Data citations 5–6, and commented on the draft manuscript. C.R. commented on the draft manuscript. Y.S.-F. led the development of Fig. 1, contributed towards the development of Data citations 9–14 and of Supplementary Figs 1–4, and commented on the draft manuscript. B.Y. commented on the draft manuscript.

Additional Information

Table 1 is only available in the online version of this paper.

Supplementary information accompanies this paper at <http://www.nature.com/sdata>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Brooks, T. M. *et al.* Analysing biodiversity and conservation knowledge products to support regional environmental assessments. *Sci. Data* 3:160007 doi: 10.1038/sdata.2016.7 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0>

Metadata associated with this Data Descriptor is available at <http://www.nature.com/sdata/> and is released under the CC0 waiver to maximize reuse.