



Morbidity and mortality after bilateral simultaneous total knee arthroplasty in a fast-track setting

Gromov, Kirill; Troelsen, Anders; Otte, Kristian Stahl; Ørsnes, Thue; Husted, Henrik

Published in:
Acta Orthopaedica


DOI:
[10.3109/17453674.2016.1141631](https://doi.org/10.3109/17453674.2016.1141631)

Publication date:
2016

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY-NC](#)

Citation for published version (APA):
Gromov, K., Troelsen, A., Otte, K. S., Ørsnes, T., & Husted, H. (2016). Morbidity and mortality after bilateral simultaneous total knee arthroplasty in a fast-track setting. *Acta Orthopaedica*, 87(3), 286-290.
<https://doi.org/10.3109/17453674.2016.1141631>



Morbidity and mortality after bilateral simultaneous total knee arthroplasty in a fast-track setting

Kirill Gromov, Anders Troelsen, Kristian Stahl Otte, Thue Ørsnes & Henrik Husted

To cite this article: Kirill Gromov, Anders Troelsen, Kristian Stahl Otte, Thue Ørsnes & Henrik Husted (2016) Morbidity and mortality after bilateral simultaneous total knee arthroplasty in a fast-track setting, *Acta Orthopaedica*, 87:3, 286-290, DOI: [10.3109/17453674.2016.1141631](https://doi.org/10.3109/17453674.2016.1141631)

To link to this article: <http://dx.doi.org/10.3109/17453674.2016.1141631>



© 2016 The Author(s). Published by Taylor & Francis on behalf of the Nordic Orthopedic Federation.



Published online: 28 Jan 2016.



Submit your article to this journal [↗](#)



Article views: 711



View related articles [↗](#)



View Crossmark data [↗](#)

Morbidity and mortality after bilateral simultaneous total knee arthroplasty in a fast-track setting

Kirill GROMOV¹, Anders TROELSEN¹, Kristian Stahl OTTE¹, Thue ØRSNES¹, and Henrik HUSTED^{1,2}

¹ Department of Orthopedic Surgery, Copenhagen University Hospital Hvidovre, Copenhagen; ² The Lundbeck Foundation Center for Fast-Track Hip and Knee Replacement, Copenhagen, Denmark.

Correspondence: kirgromov@gmail.com

Submitted 2015-07-15. Accepted 2015-12-07.

Background and purpose — The safety aspects of bilateral simultaneous total knee arthroplasty (BSTKA) are still debated. In this retrospective single-center study, we investigated early morbidity and mortality following BSTKA in a modern fast-track setting. We also identified risk factors for re-admission within 90 days and for a length of stay (LOS) of more than 5 days.

Patients and methods — 284 patients were selected to receive BSTKA at our institution from 2008 through 2014 in a well-described, standardized fast-track setup (Husted 2012 a, b). All re-admissions within 90 days were identified and mortality rates and time until death were recorded. Transfusion rates and numbers of transfusions were also recorded. Logistic regression analysis was used to identify risk factors for re-admission within 90 days, and also for a LOS of more than 5 days.

Results — 90-day mortality was 0%. 10% of the patients were re-admitted within 90 days. Median time to re-admission was 18 (3–75) days. 153 patients (54%) received postoperative blood transfusions. An ASA score of 3 was identified as an independent risk factor for re-admission within 90 days (OR = 5, 95% CI: 1.3–19) and for LOS of > 5 days (OR = 6, 95% CI: 1.6–21). Higher BMI was a weak risk factor for re-admission within 90 days.

Interpretation — BSTKA in selected patients without cardiopulmonary disease in a fast-track setting appears to be safe with respect to early postoperative morbidity and mortality. Surgeons should be aware that patients with an ASA score of 3 have an increased risk of re-admission and a prolonged length of stay, while patients with higher BMI have an increased risk of re-admission following BSTKA.

Fast-track knee arthroplasty is a cost-efficient and safe treatment method for osteoarthritis of the knee, with excellent patient satisfaction, early functional recovery, and low post-

operative morbidity and mortality (Kehlet 2013). Close to one-third of patients scheduled to undergo primary TKA present with bilateral symptoms (Metcalf et al. 2012). In these cases, the surgeon can choose to either perform a staged procedure or a simultaneous bilateral procedure, depending on comorbidities, the preference of the patient, and so on. The proposed benefits of bilateral simultaneous TKA (BSTKA), including reduced total financial costs and shorter total convalescence period (Odum et al. 2013, Lin et al. 2014), have led to an increased number of simultaneous procedures in recent decades (Hooper et al. 2009, Memtsoudis et al. 2013, Poultsides et al. 2014, Lindberg-Larsen et al. 2015). However, the safety aspects of BSTKA are still debated, as several studies have shown a higher risk of postoperative complications and even higher postoperative mortality after simultaneous procedures than after staged procedures (Restrepo et al. 2007, Stefánsdóttir et al. 2008, Hu et al. 2011, Bolognesi et al. 2013). On the other hand, other authors have reported that BSTKA is safe without any increase in postoperative morbidity and mortality (Hooper et al. 2009, Bini et al. 2014, Lindberg-Larsen et al. 2015).

Comparison of studies on BSTKA is complicated by the use of different inclusion criteria, different patient populations, and different perioperative protocols. Few studies have investigated mortality and morbidity following BSTKA in a fast-track setting (Husted et al. 2011, Hart et al. 2015). Further studies are needed to evaluate the safety aspects of simultaneous procedures performed in a modern fast-track setup, with surgery performed without drains or tourniquet, same-day mobilization, and a short length of stay (LOS).

In this retrospective single-center study, we investigated early morbidity and mortality following BSTKA in a modern fast-track setting. We also identified risk factors for re-admission within 90 days and for a LOS of more than 5 days.

Patients and methods

284 patients were selected to receive BSTKA at our institution from 2008 through 2014 in a well-described, standardized fast-track setup (Husted 2012). All patients presented with painful bilateral knee osteoarthritis, had no known cardiopulmonary comorbidity, and were operated by a consultant specialized in joint replacement surgery. Minimum follow-up was 1 year (range: 1–7.5). Median age was 65 (33–86) years, 53% of the patients (150/284) were female, and mean BMI was 30 (19–50). 26% of the patients had an ASA score of 1, 68% had an ASA score of 2, and 6% had an ASA score of 3.

All patients received a total knee prostheses—AGC (Biomet), Vanguard (Biomet), NexGen (Zimmer), or PFC Sigma (DePuy Synthes)—using a standard medial parapatellar approach without the use of a tourniquet. The left knee was operated first and surgery was performed sequentially. Cutting guides were applied externally on the tibia and intramedullary on the femur. The standard surgical protocol included spinal analgesia, standardized fluid management, use of preoperative intravenous tranexamic acid (TXA), plugging of the femoral medullary canal, absence of drains, application of local infiltration analgesia (LIA), and use of postoperative compression bandaging. The multimodal opioid-sparing analgesic regime included use of non-steroidal anti-inflammatory drugs, paracetamol, local infiltration anesthesia (LIA applied at the end of surgery only, no catheters), and opioids only upon request.

The patients were transferred from the postoperative recovery unit to the patient ward after a few hours, where immediate mobilization was attempted, allowing full-weight bearing. Physiotherapy was started on the day of surgery and continued until discharge. Rivaroxaban (Bayer, Denmark) was used as oral thromboprophylaxis, starting 6–8 hours postoperatively and continuing on a daily basis until discharge. Mechanical thromboprophylaxis and extended oral thromboprophylaxis were not used. A transfusion protocol was applied with blood transfusions triggered by a 25% decrease in postoperative hemoglobin compared to preoperative values, combined with clinical signs of anemia such as dizziness, shortness of breath, and/or pallor. Patients were discharged when the functional discharge criteria were met, which included being able to undertake independent personal care, being able to walk 70 m with or without crutches, being able to get in and out of bed, and being able to get up independently from a chair or toilet.

LOS was recorded and counted in whole days as the number of postoperative nights in the hospital. All re-admissions within 90 days were identified and the reason for re-admission recorded. Mortality rates and time until death were recorded. Transfusion rates and number of transfusions were also recorded.

Statistics

Comparisons were performed using 2-sample t-test for normally distributed data and a two-sample Wilcoxon rank-sum

Table 1. Reasons for 29 re-admissions within 90 days of BSTKA

Reason for re-admission	n
Suspicion of DVT ^a	2
Suspicion of infection ^b	6
Cardiac ^c	4
Thromboembolic ^d	2
Infection ^e	8
Other ^f	7
Total	29

^a Suspicion of deep-vein thrombosis (DVT), refuted by ultrasound.

^b Suspicion of infection, refuted by clinical examination and blood sampling.

^c Patients with chest pains. Cardiac ischemia refuted by blood sampling. 1 patient with atrial fibrillation (medically treated).

^d 2 patients with DVT confirmed by ultrasound (medically treated).

^e 6 patients with clinically confirmed infection (treated with surgical revision). 2 patients with superficial site infection (treated with antibiotics).

^f 2 patients admitted for further mobilization, 1 patient admitted due to wound oozing without suspicion of infection, 1 patient with urinary tract infection, 2 patients with gastric symptoms, and 1 patient with dehydration.

test for data that were not normally distributed. Proportions were compared using the Pearson chi-squared test. Logistic regression analysis was used to identify risk factors for re-admission within 90 days and for LOS of more than 5 days, while adjusting for age, sex, ASA score, and BMI.

Any p-value less than 0.05 was considered statistically significant. All analyses were performed using SPSS statistics software version 21.

Results

90-day mortality was 0%. 10% of patients were re-admitted within 90 days, with 8% being re-admitted within 30 days. Median time to re-admission was 18 (3–75) days. Surgical site infection treated either with antibiotics alone (n = 2) or with revision (n = 6) was the most common reason for re-admission (Table 1). The proportion of re-admitted patients was not significantly different in between years during the study period (p = 0.9). Median LOS for all patients was 3 (2–30) days. 14% of all patients had a LOS of more than 5 days (data not shown).

54% of patients received blood transfusions postoperatively and 21% received more than 2 units of blood.

Patients with an ASA score of 3 had an OR of 4.9 (95% CI: 1.3–19) for re-admission within 90 days (Table 2) and an OR of 5.8 (95% CI: 1.6–21) for a LOS of more than 5 days (Table 3). Higher BMI was associated with an increased risk of re-admission within 90 days (OR = 1.1, 95% CI: 1.01–1.2) (Table 2).

Table 2. Odds ratio for re-admission within 90 days of BSTKA

Parameter	OR (95% CI)	p-value
Age, years ^a	1.1 (0.99–1.1)	0.09
Female sex	0.93 (0.40–2.2)	0.9
ASA score 2	0.70 (0.25–2.0)	0.5
ASA score 3	4.9 (1.3–19)	0.02
BMI ^b	1.1 (1.01–1.2)	0.04

^a Age included as a continuous variable.

^b BMI included as a continuous variable.

Table 3. Odds ratio for a length of stay (LOS) of more than 5 days following primary BSTKA

Parameter	OR (95% CI)	p-value
Age, years ^a	1.0 (0.99–1.1)	0.1
Female sex	1.2 (0.59–2.5)	0.6
ASA score 2	1.2 (0.48–3.1)	0.7
ASA score 3	5.8 (1.6–21)	0.008
BMI ^b	1.0 (0.95–1.1)	0.7

^a Age included as a continuous variable.

^b BMI included as a continuous variable.

Discussion

The safety of BSTKA regarding mortality and morbidity is debatable, as several studies have shown higher mortality following simultaneous bilateral procedures than after staged bilateral procedures (Parvizi et al. 2001, Stefánsdóttir et al. 2008) while others have failed to show any differences (Bullcock et al. 2003, Lindberg-Larsen et al. 2015). The 3 most recent meta-analyses comparing morbidity and mortality following BSTKA (Restrepo et al. 2007, Fu et al. 2013, Hussain et al. 2013) all concluded that BSTKA carries a higher risk of postoperative mortality. Conclusions on postoperative complications are less sure, as meta-analyses by both Hussein et al. (2007) and Fu et al. (2001) found similar rates of thromboembolic and cardiac complications and also infections, while Restrepo et al. (2007) found a higher risk of cardiac and pulmonary complications following BSTKA than after staged procedures.

Our finding of a 90-day mortality of 0% following BSTKA in a standardized fast-track setting is encouraging, and it is similar to previous reports on early postoperative mortality for bilateral procedures in a fast-track setting, as both Lindberg et al. (2015) and Husted et al. (2011) also reported 0% 90-day mortality following BSTKA, while Poultside et al. (2014) found a 30-day mortality rate of 0.04% in a cohort of 4,825 BSTKA patients. It is important to point out that the study by Husted et al. (2011) included patients who were operated in 2008 (n = 32), thus having a small overlap with the present study. The low mortality rates stand in contrast to the 0.5%

30-day mortality following BSTKA reported by Parvizi et al. (2001), the 0.15% in-hospital mortality reported by Odum et al. (2014), and the 0.3% in-hospital mortality reported by Memtsoudis et al. (2009). One possible explanation for the zero mortality found in our study was the fast-track setup itself, which may have reduced postoperative morbidity and mortality for both total knee arthroplasty patients and total hip arthroplasty patients (Husted et al. 2010a,b, Kehlet 2013, Lindberg-Larsen et al. 2013, 2015).

The 10% re-admission rate within 90 days and the 8% re-admission rate within 30 days found in the present study is comparable to the 7% re-admission rate within 30 days following simultaneous bilateral procedures reported by Lindberg et al. (2015) in a nationwide study, and to the 7% re-admission rate within 30 days following staged bilateral procedures reported by Kheir et al. (2014). Hart et al. (2015) have reported lower 30-day re-admission rates following BSTKA—4% as compared to the 8% value found in the present study, but this might be explained by varying definitions of re-admission between studies, and the possibility of complete follow-up (Husted 2012). A 90-day re-admission rate of 9% following unilateral TKA in a modern fast-track setting has been reported in a recent nationwide study (Jørgensen and Kehlet 2013). This supports the safety aspect of BSTKA, which in our hands has similar re-admission rates to those of unilateral TKA, as the patients receiving 2 unilateral TKAs would have a higher cumulative re-admission rate over time.

Our finding that postoperative infection is the most common cause of re-admission is also in agreement with the findings of both Kheir et al. (2014) and Lindberg et al. (2015). It is interesting that there were no re-admissions due to pulmonary embolism (PE) and only 2 re-admissions (0.7%) due to deep-vein thrombosis (DVT) in our patient cohort, which can be compared with the DVT rate of 0.9% and the PE rate of 1% within 60 days reported by Meehan et al. (2011) and the DVT and PE rates of 1% and 0.8%, respectively, following BSTKA reported by Memtsoudis et al. (2009) in nationwide studies. The 0.7% (2/284) risk of re-admission within 90 days due to thromboembolic events found in our study is lower than the 2% 90-day risk of re-admission due to thromboembolic events reported in fast-track departments in a nationwide study (Glassou et al. 2014). Less comorbidity in patients undergoing BSTKA is the most likely explanation for this difference.

The median LOS of 3 days found in the present study is similar to LOS figures in previous studies on fast-track total knee arthroplasty (Husted et al. 2011, Jørgensen and Kehlet 2013). This shows that almost all the patients selected can fulfill the functional discharge criteria within just a few days after this major surgical procedure.

We found a high transfusion rate, as more than 50% of patients required postoperative blood transfusions. This finding is not unexpected, as several previous studies have found high transfusion rates following simultaneous bilateral procedures: 95% was found by Stubbs et al. (2005), 55% was

found by Jenny et al. (2013), and 45% was found by Husted et al. (2011). A recent study performed by Kim et al. (2014) did, however, show a much lower transfusion rate of 7% for BSTKA when TXA was administered in multiple doses, indicating the possibility of reducing transfusions more than the blood-saving protocol currently being used at our institution. However, we do not know what role the high transfusion rate reported in our study plays regarding the complication rate and postoperative outcomes, as blood transfusion has been linked to infections (Friedman et al. 2014). Further investigation is therefore required, focusing on intraoperative and total blood loss following BSTKA in order to optimize conservation of blood.

Finally, we found patients with an ASA score of 3 to have an increased risk of re-admission within 90 days and an increased risk of having a LOS of more than 5 days. This is hardly surprising, as similar results have been reported for unilateral procedures (Husted et al. 2010a, Husted 2012). It is, however, surprising that 17 patients (6%) with an ASA score of 3 were offered—and underwent—BSTKA in our study. This contrasts with a recent consensus paper, which stated that only patients with few comorbidities should be offered BSTKA (Memtsoudis et al. 2013). This is supported by our finding that 7 of the 17 patients with an ASA score of 3 were re-admitted within 90 days. However, it is not possible to speculate whether these re-admissions could have been avoided if patients with an ASA score of 3 had undergone a staged bilateral procedure instead.

Obesity is a well-known risk factor for complications following TKA, including increased risk of infections, of prolonged wound drainage, and of thromboembolic complications (Belmont et al. 2014, Thornqvist et al. 2014, Alvi et al. 2015, Si et al. 2015). Thus, our finding that higher BMI was associated with an increased risk, albeit small, of re-admission within 90 days is in agreement with the previous literature (Fabi et al. 2011).

The main limitation of our study was the retrospective design without any control group. The retrospective data collection allowed possible inaccuracies regarding the parameters collected, and the lack of a control group did not allow comparisons and therefore limited our conclusions. The optimal design for evaluating the safety and efficiency of BSTKA would be a randomized controlled study comparing staged and simultaneous procedures. However, such a setup would have a challenging patient inclusion, which is one of the reasons why such trials have not been performed to date (Memtsoudis et al. 2013). Addition of a control group of staged bilateral procedures, as previously done by Husted et al. (2011), would allow comparison of simultaneous with staged procedures. However, such case-control studies with matching of 1 or multiple parameters would still permit selection bias, as the surgeon's choice for performing a staged rather than a simultaneous procedure and vice versa would be unaccounted for. We have therefore presented strictly descriptive data showing safety

aspects with respect to mortality and re-admission following BSTKA in selected patients. Secondly, we only investigated re-admissions and mortality without investigating patient-reported outcomes. However, we have already reported patient satisfaction in another cohort of BSTKA patients from our study population (Husted et al. 2011). Thirdly, ASA score was the only parameter that we included to account for patient comorbidities, and it is known to be associated with great interobserver variability (Mak et al. 2002). Finally, the follow-up for re-admission and mortality was limited to 90 days, which does not allow us to make any conclusions on late surgical complications such as aseptic loosening. However, evaluation of late surgical complications is outside the scope of the current study, and we believe that death more than 90 days after the procedure is unlikely to have been associated with the procedure itself.

In conclusion, BSTKA in selected patients without cardiopulmonary disease in a fast-track setting appears to be safe regarding early postoperative morbidity and mortality. Surgeons should be aware that patients with an ASA score of 3 have an increased risk of re-admission and prolonged length of stay, while patients with higher BMI have an increased risk of re-admission after BSTKA.

KG, AT, and HH wrote the protocol and all the authors revised it. KG, KSO, and TØ undertook all data gathering. KG, AT, and HH performed and evaluated all the statistical analyses. KG wrote the first draft of the manuscript and HH revised it; all the authors revised the draft and approved the final version. All the authors had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analyses.

No competing interests declared.

- Alvi H M, Mednick R E, Krishnan V, Kwasny M J, Beal M D, Manning D W. The Effect of BMI on 30 Day Outcomes Following Total Joint Arthroplasty. *J Arthroplasty* 2015; 30(7): 1113-7.
- Belmont P J, Goodman G P, Waterman B R, Bader J O, Schoenfeld A J. Thirty-day postoperative complications and mortality following total knee arthroplasty: incidence and risk factors among a national sample of 15,321 patients. *J Bone Joint Surg Am* 2014; 96(1): 20-6.
- Bini S A, Khatod M, Inacio MCS, Paxton EW. Same-day versus staged bilateral total knee arthroplasty poses no increase in complications in 6672 primary procedures. *J Arthroplasty* 2014; 29(4): 694-7.
- Bolognesi M P, Watters T S, Attarian D E, Wellman S S, Setoguchi S. Simultaneous vs staged bilateral total knee arthroplasty among Medicare beneficiaries, 2000-2009. *J Arthroplasty* 2013; 28(8 Suppl): 87-91.
- Bullock D P, Sporer S M, Shirreffs T G. Comparison of simultaneous bilateral with unilateral total knee arthroplasty in terms of perioperative complications. *J Bone Joint Surg Am* 2003; 85-A(10): 1981-6.
- Fabi D W, Mohan V, Goldstein W M, Dunn J H, Murphy B P. Unilateral vs bilateral total knee arthroplasty risk factors increasing morbidity. *J Arthroplasty* 2011; 26(5): 668-73.
- Friedman R, Homering M, Holberg G, Berkowitz S D. Allogeneic blood transfusions and postoperative infections after total hip or knee arthroplasty. *J Bone Joint Surg Am* 2014 19; 96(4): 272-8.

- Fu D, Li G, Chen K, Zeng H, Zhang X, Cai Z. Comparison of clinical outcome between simultaneous-bilateral and staged-bilateral total knee arthroplasty: a systematic review of retrospective studies. *J Arthroplasty* 2013; 28(7): 1141-7.
- Glassou E N, Pedersen A B, Hansen T B. Risk of re-admission, reoperation, and mortality within 90 days of total hip and knee arthroplasty in fast-track departments in Denmark from 2005 to 2011. *Acta Orthop* 2014; 85(5): 493-500.
- Hart A, Antoniou J, Brin Y S, Huk O L, Zukor D J, Bergeron S G. Simultaneous bilateral versus unilateral total knee arthroplasty: a comparison of 30-day readmission rates and major complications. *J Arthroplasty* 2015; Jul 21. doi: 10.1016/j.arth.2015.07.031. [Epub ahead of print]
- Hooper G J, Hooper N M, Rothwell A G, Hobbs T. Bilateral total joint arthroplasty: the early results from the New Zealand National Joint Registry. *J Arthroplasty* 2009; 24(8): 1174-7.
- Hu J, Liu Y, Lv Z, Li X, Qin X, Fan W. Mortality and morbidity associated with simultaneous bilateral or staged bilateral total knee arthroplasty: a meta-analysis. *Arch Orthop Trauma Surg* 2011; 131(9): 1291-8.
- Hussain N, Chien T, Hussain F, Bookwala A, Simunovic N, Shetty V, et al. Simultaneous versus staged bilateral total knee arthroplasty: a meta-analysis evaluating mortality, peri-operative complications and infection rates. *HSS J* 2013; 9(1): 50-9.
- Husted H. Fast-track hip and knee arthroplasty: clinical and organizational aspects. *Acta Orthop (Suppl 346)* 2012; 83: 1-39.
- Husted H, Otte KS, Kristensen BB, Orsnes T, Kehlet H. Readmissions after fast-track hip and knee arthroplasty. *Arch Orthop Trauma Surg* 2010a; 130(9): 1185-91.
- Husted H, Otte K S, Kristensen B B, Ørsnes T, Wong C, Kehlet H. Low risk of thromboembolic complications after fast-track hip and knee arthroplasty. *Acta Orthop* 2010b; 81(5): 599-605.
- Husted H, Troelsen A, Otte K S, Kristensen B B, Holm G, Kehlet H. Fast-track surgery for bilateral total knee replacement. *J Bone Joint Surg Br* 2011; 93(3): 351-6.
- Jenny J-Y, Trojani C, Prudhon J-L, Vielpeau C, Saragaglia D, Houillon C, et al. Simultaneous bilateral total knee arthroplasty. A multicenter feasibility study. *Orthop Traumatol Surg Res* 2013; 99(2): 191-5.
- Jørgensen C C, Kehlet H. Role of patient characteristics for fast-track hip and knee arthroplasty. *Br J Anaesth* 2013; 110(6): 972-80.
- Kehlet H. Fast-track hip and knee arthroplasty. *Lancet* 2013; 381(9878): 1600-2.
- Kheir M M, Clement R C, Derman P B, Flynn D N, Speck R M, Levin L S, et al. Are there identifiable risk factors and causes associated with unplanned readmissions following total knee arthroplasty? *J Arthroplasty* 2014; 29(11): 2192-6.
- Kim T K, Chang C B, Kang Y G, Seo E S, Lee J H, Yun J H, et al. Clinical value of tranexamic acid in unilateral and simultaneous bilateral TKAs under a contemporary blood-saving protocol: a randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc* 2014; 22(8): 1870-8.
- Lin AC-C, Chao E, Yang C-M, Wen H-C, Ma H-L, Lu T-C. Costs of staged versus simultaneous bilateral total knee arthroplasty: a population-based study of the Taiwanese National Health Insurance Database. *J Orthop Surg Res* 2014; 9: 59.
- Lindberg-Larsen M, Joergensen C C, Husted H, Kehlet H. Simultaneous and staged bilateral total hip arthroplasty: a Danish nationwide study. *Arch Orthop Trauma Surg* 2013; 133(11): 1601-5.
- Lindberg-Larsen M, Jørgensen C C, Husted H, Kehlet H. Early morbidity after simultaneous and staged bilateral total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2015; 23(3): 831-7.
- Mak P H K, Campbell R C H, Irwin M G. The ASA Physical Status Classification: inter-observer consistency. *American Society of Anesthesiologists. Anaesth Intensive Care* 2002; 30(5): 633-40.
- Meehan J P, Danielsen B, Tancredi D J, Kim S, Jamali A A, White R H. A population-based comparison of the incidence of adverse outcomes after simultaneous-bilateral and staged-bilateral total knee arthroplasty. *J Bone Joint Surg Am* 2011; 93(23): 2203-13.
- Memtsoudis S G, Hargett M, Russell L A, Parvizi J, Cats-Baril W L, Stundner O, et al. Consensus statement from the consensus conference on bilateral total knee arthroplasty group. *Clin Orthop Relat Res* 2013; 471(8): 2649-57.
- Memtsoudis S G, Ma Y, González Della Valle A, Mazumdar M, Gaber-Baylis L K, MacKenzie C R, et al. Perioperative outcomes after unilateral and bilateral total knee arthroplasty. *Anesthesiology* 2009; 111(6): 1206-16.
- Metcalfe A J, Andersson M L E, Goodfellow R, Thorstensson C A. Is knee osteoarthritis a symmetrical disease? Analysis of a 12 year prospective cohort study. *BMC Musculoskelet Disord* 2012; 13: 153.
- Odum S M, Troyer J L, Kelly M P, Dedini R D, Bozic K J. A cost-utility analysis comparing the cost-effectiveness of simultaneous and staged bilateral total knee arthroplasty. *J Bone Joint Surg Am* 2013; 95(16): 1441-9.
- Parvizi J, Sullivan T A, Trousdale R T, Lewallen D G. Thirty-day mortality after total knee arthroplasty. *J Bone Joint Surg Am* 2001; 83-A(8): 1157-61.
- Poultides L A, Rasouli M R, Maltenfort M G, Parvizi J, Memtsoudis S G, Sculco T P. Trends in Same-Day Bilateral Total Knee Arthroplasty. *J Arthroplasty* 2014; 29(9): 1713-6.
- Restrepo C, Parvizi J, Dietrich T, Einhorn T A. Safety of simultaneous bilateral total knee arthroplasty. A meta-analysis. *J Bone Joint Surg Am* 2007; 89(6): 1220-6.
- Si H-B, Zeng Y, Shen B, Yang J, Zhou Z-K, Kang P, et al. The influence of body mass index on the outcomes of primary total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2015; 23(6): 1824-32.
- Stefánsdóttir A, Lidgren L, Robertsson O. Higher early mortality with simultaneous rather than staged bilateral TKAs: results from the Swedish Knee Arthroplasty Register. *Clin Orthop Relat Res* 2008; 466(12): 3066-70.
- Stubbs G, Pryke S E R, Tewari S, Rogers J, Crowe B, Bridgfoot L, et al. Safety and cost benefits of bilateral total knee replacement in an acute hospital. *ANZ J Surg* 2005; 75(9): 739-46.
- Thornqvist C, Gislason G H, Køber L, Jensen P F, Torp-Pedersen C, Anderson C. Body mass index and risk of perioperative cardiovascular adverse events and mortality in 34,744 Danish patients undergoing hip or knee replacement. *Acta Orthop* 2014; 85(5): 456-62.