

UNIVERSITY OF COPENHAGEN



## How does a plant orchestrate defense in time and space? Using glucosinolates in *Arabidopsis* as case study

Burow, Meike; Halkier, Barbara Ann

*Published in:*  
Current Opinion in Plant Biology

*DOI:*  
[10.1016/j.pbi.2017.04.009](https://doi.org/10.1016/j.pbi.2017.04.009)

*Publication date:*  
2017

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[CC BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)

*Citation for published version (APA):*  
Burow, M., & Halkier, B. A. (2017). How does a plant orchestrate defense in time and space? Using glucosinolates in *Arabidopsis* as case study. *Current Opinion in Plant Biology*, 38, 142-147.  
<https://doi.org/10.1016/j.pbi.2017.04.009>



ELSEVIER

# How does a plant orchestrate defense in time and space? Using glucosinolates in *Arabidopsis* as case study

Meike Burow and Barbara Ann Halkier



The sessile nature of plants has caused plants to develop means to defend themselves against attacking organisms. Multiple strategies range from physical barriers to chemical warfare including pre-formed antipins as well as phytoalexins produced only upon attack. While phytoalexins require rapid induction, constitutive defenses can impose ecological costs if they deter pollinators or attract specialized herbivores. In the model *Arabidopsis thaliana*, the well-characterized glucosinolate antipins are categorized into different classes, aliphatic and indole glucosinolates, depending on their amino acid precursor. Using glucosinolates in *Arabidopsis* as case study, we will discuss how plants orchestrate synthesis, storage and activation of pre-formed defense compounds spatially and temporally.

## Address

DynaMo Center, Department of Plant and Environmental Sciences, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark

Corresponding author: Halkier, Barbara Ann ([bah@plen.ku.dk](mailto:bah@plen.ku.dk))

**Current Opinion in Plant Biology** 2017, **38**:142–147

This review comes from a themed issue on **Biotic interactions**

Edited by **Silke Robatzek** and **Sarah Lebeis**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 30th May 2017

<http://dx.doi.org/10.1016/j.pbi.2017.04.009>

1369-5266/© 2017 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Specialized plant metabolites are well-known for their role in interactions between plants and their distinct biotic environments. Consequently, and as opposed to general metabolites, specialized metabolites are restricted to specific taxa. Their enormous diversity in chemical structures, biosynthetic pathways, and modes of action [1,2] has evolved under selection pressure imposed by specific combinations of harmful and beneficial interacting organisms. Chemical defense strategies in plants range from constitutive accumulation over induced defenses to activated defense systems. The latter depend on at least two components—the preformed, inactive precursor of a bioactive compounds and the activating enzyme machinery. For the plant to ensure that

appropriate chemical defenses are present in the right tissue and at the right time, their biosynthesis, transport and storage needs to be tightly regulated in space and time. As different defense strategies underlie different defense pathways, a general discussion of their spatio-temporal regulation is troublesome. Thus, this review focuses on glucosinolates as model for activated chemical defenses.


As primary chemical defense compounds, *Arabidopsis* synthesizes up to 40 different glucosinolates [3]. The glucosinolates themselves show only very limited biological activities, and depend on activation initiated by co-occurring thioglycosidases called myrosinases. Chemical and enzymatical rearrangements after myrosinase-catalyzed hydrolysis give rise to a range of products toxic to a wide range of organisms [1,4]. Depending on the amino acid-derived glucosinolate side chain and the presence of specifier proteins, each glucosinolate can be activated to one or more products with different biological activities [5]. Variation in the type of glucosinolate hydrolysis products is observed among different species within the Brassicaceae, among *Arabidopsis* accessions and even among tissues of the same plant [6–8]. Upon herbivory, glucosinolate levels can be further induced and induction of the nitrile-specifier protein NSP1 can change the outcome of glucosinolate hydrolysis and thereby shift a direct to an indirect defense strategy [9,10]. This underlines the versatile and dynamic nature of glucosinolate-based defense.

## Storage of glucosinolates in S-cells and seeds

In the unattacked *Arabidopsis* plant, glucosinolates are stored in laticifer-like S-cells within the phloem cap region outside the vasculature and along the leaf margin [11–15] (Figure 1). These cells contain >130 mM glucosinolates stored under high turgor pressure. Although laticifer-like cells are known from, for example, rubber trees it remains an open question whether the localization of laticifer-like cells (such as the S-cells) in the phloem cap region is a Brassicales-specific evolution or whether universally found for other defense compounds.

Other storage sites include the seeds that accumulate high levels of glucosinolates that are imported into the seeds that lack *de novo* biosynthesis [16,17]. Knowledge about the seed loading process is limited. The so-far only identified glucosinolate transporters, the plasma

Figure 1



Spatio-temporal control of glucosinolate-based defense. Like numerous other specialized metabolites, glucosinolates play a key role in plant biotic interactions, both above- and below-ground. Their presence and activation in the right tissue at the right time is critical for their biological functions. Glucosinolate profiles characteristic for a given tissue, developmental stage and combination of environmental factors are dynamically shaped by biosynthesis and transport. Along vascular bundles, rapid activation upon tissue damage relies on close proximity of glucosinolate storage cells (S-cells shown in yellow in the schematic stem cross section) and myrosin cells (shown in red).

membrane-localized glucosinolate transporter 1 (GTR1) and glucosinolate transporter 2 (GTR2) importers, are essential for loading glucosinolates into the seeds as demonstrated by glucosinolate-free seeds of the *gtr1 gtr2* mutant [18]. The level of glucosinolates in a given tissue is subject to a complex feedback regulation. Plants overexpressing the MYB28 – a major positive regulator of aliphatic glucosinolates – accumulated threefold higher levels of aliphatic glucosinolates in the foliar tissue, while their seeds showed only a moderate increase [19]. Thus, independent of the levels of glucosinolates in vegetative

parts, the levels in the seeds appear to reach a certain maximum. It is currently not known how the plant senses that the seeds have reached this threshold.

### Sinks within the leaves

Glucosinolate concentration in leaves decreases with age until virtually gone upon senescence [8]. Whether the glucosinolates are (re)mobilized to sink tissue or turned over is unknown. It was suggested that leaf glucosinolates were destined to the seeds based on increased levels in leaves of *gtr1 gtr2* mutant [18]. However, it turned out that

the overaccumulation in leaves was due to glucosinolates derived from roots via the xylem [20,21]. This suggests that upon senescence the leaf-synthesized glucosinolates are not transported long distance to the seeds, but rather turned over. How leaf-synthesized glucosinolates are turned over in the intact leaves upon senescence is a puzzle, although glucosinolate breakdown independent of the classical myrosinases thioglucoside glucohydrolase 1 (TGG1) and thioglucoside glucohydrolase 2 (TGG2) has been shown [22]. By contrast, the root-derived glucosinolates accumulating in leaves in the *gr1 gr2* mutant [21] are apparently turned over at much lower rate. Possibly, glucosinolates that arrive via xylem accumulate in different sites than the leaf-synthesized ones.

Recently, the cuticular leaf layer was identified as glucosinolate sink [23<sup>\*</sup>]. The surface localization of glucosinolates has been a controversial topic for many years as various methods have yielded ambiguous results. Recently, Shroff *et al.* [23<sup>\*</sup>] used three independent mass spectrometry methods to elegantly detect and quantify intact glucosinolates at the surface. In addition to previous findings, where glucosinolates were shown to accumulate along the major midrib and the leaf margin [24], the surface-localized glucosinolates represent 1–5% of total glucosinolates and are evenly distributed across the epidermis, except for the midrib [23<sup>\*</sup>]. The distribution between adaxial and abaxial surfaces was found to be uneven with 15–30-fold more on the abaxial surface. How glucosinolates get exported out of epidermis to the cuticula is not known. Noticeably, particularly 4-methylthiobutyl glucosinolate (4MTB) (eightfold) and indole-3-methyl glucosinolate (I3M) (threefold) are overrepresented in both epidermis [21] and cuticula [23<sup>\*</sup>]. The significance of this is not known, but the levels are sufficient to function as oviposition cues [23<sup>\*</sup>].

### The rhizosphere is a sink

Recently, intact glucosinolates [25] as well as hydrolysis products [25,26] were detected in the exudate of Arabidopsis roots, as they have previously been detected in *Brassica* root exudates [27,28]. Intuitively, one would have anticipated that rhizosecreted phytochemicals were produced in the outer cell layers of the root. However, in Arabidopsis the route from site of synthesis to the rhizosphere required GTR-mediated import of stele-synthesized glucosinolates into the symplasm [25]. This indicated that glucosinolates must be exported from the cell in which they are synthesized and that GTR1 and GTR2 might be essential in balancing above- and below-ground defense in response to environmental challenge. The effects of *Brassica* root exudates in suppression of soil-born pests have primed an interest in use of glucosinolate-containing plants in pest management, a process termed biofumigation [29]. The endophyte *Piriformospora indica* requires a certain level of glucosinolate hydrolysis to maintain its status as commensalistic/mutualistic

endophyte [30], which indicates that the impact of intact glucosinolates and their hydrolysis products on the microbial community in the rhizosphere represents an interesting future research area.

### Are glucosinolates mobilized upon attack?

Until now we have discussed glucosinolates as phytoanticipines that accumulate in various sinks in the unchallenged plant. However, upon attack by biotrophic pathogens, for example, *Blumeria* and *Phytophthora* species, cell-autonomous defense linked to indole glucosinolates has been shown to play role a plant innate immunity [31,32]. Upon such attacks, unmodified indole glucosinolates I3M produced along the vasculature [33] and stored in the epidermal cell [21] are modified by the induced enzyme CYP81F2 to produce 4-methoxy-indole glucosinolate [34]. It is currently not known whether the epidermal pool of I3M is being replenished by *de novo* synthesis in the epidermal cell or whether the vasculatory synthesis sites deliver upon attack? Also, where in the unchallenged epidermal cell is the pre-formed I3M stored? Is it in the vacuole where glucosinolates are normally stored? If so, how does it get remobilized to the cytosol to encounter the endoplasmic reticulum-associated CYP81F2? Future studies are required to address these questions related to the orchestration of synthesis and storage as well as remobilization and replenishment when under attack.

### How do plants coordinate development and defense?

During development, fluxes through metabolic pathways constantly need to be adjusted to account for changes in source-sink relations [35,36]. In the winter-annual Arabidopsis, the emergence of the florescence coincides with the onset of senescence in rosette leaves which marks an essential development transition. Decreasing levels of glucosinolates in senescing leaves [8] have been attributed to the emergence of the inflorescence as a new source tissue [37], but may as well be a consequence of changes in general metabolism including reduced availability of precursors and increased remobilization of nitrogen. At the same time, higher levels in young and reproductive tissues and lower levels in senescent tissues represent a typical developmental pattern for chemical defenses reflecting re-allocation of resources within the plant and its impact on the behavior of herbivores as well as that of pollinators and other beneficial organisms [38].

Thus, the networks controlling development and defense must be intimately linked to balance metabolic versus ecological costs and thereby maximize plant fitness. And indeed, the same phytohormones provide input to development, growth and regulate chemical defenses [39] supporting a view in which plant specialized metabolism is just another metabolic output of the complex signal transduction networks driving highly conserved biological processes. Despite the metabolic costs associated with

specialized metabolites, the relationship between growth and defense is not necessarily a tradeoff [40]. Instead, molecular decisions made to coordinate growth and defense are determined by the abiotic environment including the availability of nutrients, in the case of glucosinolates most importantly nitrogen and sulfur [41\*\*].

### What mediates feedback regulatory effects of specialized metabolites?

As levels of chemical defenses are strongly intertwined with developmental processes and seasonal changes in the biotic and abiotic environment, they can be expected to provide feedback regulatory input to these developmental processes to maximize the plant's ability to efficiently utilize the resources available. In line with such a model, genetic variation in the glucosinolate biosynthetic *alkenyl- and hydroxyalkyl-producing* locus (*GS-AOP*) was associated with variation in the onset of flowering [42,43] and one of the genes in the *GS-AOP* locus, *AOP2*, is involved in regulatory loop linking jasmonate signaling and glucosinolate biosynthesis [44\*]. The identification of *gulliver1/superroot2-7* – a weak allele of the cytochrome P450 CYP83B1 involved in the biosynthesis of indolic glucosinolates from tryptophan – further highlighted the established link between indole glucosinolate and indole acetic acid synthesis and hinted at a checkpoint for the coordination of the two pathways [45]. Even specific glucosinolate structures can have feedback regulatory effects as illustrated by naturally variable biomass responses in *Arabidopsis* seedlings specifically to the methionine-derived allyl glucosinolate, a product of the enzymatic activity of *AOP2* [46\*,47]. The molecular mechanisms underlying structure-specific fine-tuning of growth and development by glucosinolates and potentially other specialized metabolites remain, however, to be uncovered.

### What are the mechanisms linking circadian clocks and defense?

A multitude of developmental and physiological processes are controlled through cross-talk with molecular clocks as central regulators. Like any other multicellular organism, plants have circadian clocks to coordinate their metabolism with environmental factors by integrating biotic and abiotic factors [48,49]. In plants, this coordination is critical for local adaptation because it allows plants to anticipate predictable fluctuations in the environment and adjust development and physiology accordingly through phytohormone signaling pathways [50–53]. Also pollinators, herbivores, pathogens rely on endogenous clocks and therefore represent – at least to some extent – predictable biotic threats, as indicated by the timing of plant immune responses and increased resistance to herbivores and pathogens that coincides with the highest insect and pathogen activity [54,55].

Depending on temperature, basal levels of root glucosinolates have been reported to show a circadian pattern [56], however, quantitative changes in glucosinolate levels are relatively low and it remains to be tested how much these oscillations contribute to day time-dependent differences in plant resistance. On the level of transcriptional control, the circadian clock and glucosinolates show clear interconnectivity, again at least partially mediated by known regulators of glucosinolate biosynthesis from methionine [43]. *Arabidopsis* mutants with altered levels of *AOP2* or the transcriptional regulators *MYB28* and *MYB29* show a significantly altered circadian period providing yet another example of an output of the clock functioning as feedback regulatory input.

Future research should aim at identifying the parameters that determine whether or not oscillations in transcript levels translate into changes in metabolite levels. Whereas circadian control of glucosinolate transcripts reflects the plant's ability to anticipate herbivore and pathogen attack, diurnal changes in signaling pathways and general metabolism might be decisive factors for basal and inducible levels of defense compounds and thereby plant resistance. The central role of clock components in integrating biotic and abiotic external signal with information from internal feedback loops leaves no doubt about its impact on the dynamics of chemical defenses, but neither about future challenges in identifying the molecular mechanisms behind.

### Biotechnological use of circadian clock to promote postharvest longevity

The modular design of plants enables individual plant organs to manifest autonomous functions and continue aspects of metabolism, for example, respiration, even after separation from the parent plant. Accordingly, harvested vegetables and fruits continue to sense and respond to diverse stimuli, similarly to intact plants. In a recent study, circadian clock entrainment with light/dark cycles during postharvest storage improved plant tissue performance with respect to tissue integrity, green coloration, and chlorophyll content, compared to constant light or darkness [57\*]. In the cruciferous vegetables, kale and cabbage, the levels of the glucosinolates remained at higher levels when stored under light/dark cycles. This suggests that sustained circadian clock entrainment may be a powerful approach to promote postharvest quality and longevity and thereby reduce yield loss. By applying this finding to postharvest storage, more food could be kept for longer without refrigeration.

### Conclusion

The complexity of the orchestration of chemical defenses in plants reflects the complexity of the environments that have shaped the underlying regulatory networks. To understand how plants balance metabolic and ecological



costs to support both development and defense under fluctuating conditions requires detailed knowledge on the spatial and temporal dynamics of synthesis, storage, and mobilization of chemical defenses. We can expect the same knowledge to inspire future biotechnological strategies for improved food quality.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Chew FS: **Biological effects of glucosinolates**. In *Biologically Active Natural Products*. Edited by Cutler HG. American Chemical Society; 1988:155-181.
2. Wink M: **Evolution of secondary metabolites from an ecological and molecular phylogenetic perspective**. *Phytochemistry* 2003, **64**:3-19.
3. Halkier BA, Gershenzon J: **Biology and biochemistry of glucosinolates**. *Annu. Rev. Plant Biol.* 2006, **57**:303-333.
4. Wittstock U, Kliebenstein DJ, Lambrix V, Reichelt M, Gershenzon J: **Chapter five glucosinolate hydrolysis and its impact on generalist and specialist insect herbivores**. *Integrative Phytochemistry: From Ethnobotany to Molecular Ecology*. Elsevier; 2003:101-125.
5. Wittstock U, Burow M: **Glucosinolate breakdown in Arabidopsis: mechanism, regulation and biological significance**. *Arabidopsis Book* 2010, **8**:e0134.
6. Agerbirk N, Olsen CE: **Glucosinolate structures in evolution**. *Phytochemistry* 2012, **77**:16-45.
7. Kliebenstein DJ: **A quantitative genetics and ecological model system: understanding the aliphatic glucosinolate biosynthetic network via QTLs**. *Phytochem. Rev.* 2009, **8**:243-254.
8. Brown PD, Tokuhisa JG, Reichelt M, Gershenzon J: **Variation of glucosinolate accumulation among different organs and developmental stages of Arabidopsis thaliana**. *Phytochemistry* 2003, **62**:471-481.
9. Burow M, Losansky A, Müller R, Plock A, Kliebenstein DJ, Wittstock U: **The genetic basis of constitutive and herbivore-induced ESP-independent nitrile formation in Arabidopsis**. *Plant Physiol.* 2009, **149**:561-574.
10. Mumm R, Burow M, Bukovinszkiné-kiss G, Kazantzidou E, Wittstock U, Dicke M, Gershenzon J: **Formation of simple nitriles upon glucosinolate hydrolysis affects direct and indirect defense against the specialist herbivore, Pieris rapae**. *J. Chem. Ecol.* 2008, **34**:1311-1321.
11. Koroleva OA, Gibson TM, Cramer R, Stain C: **Glucosinolate-accumulating S-cells in Arabidopsis leaves and flower stalks undergo programmed cell death at early stages of differentiation**. *Plant J.* 2010, **64**:456-469.
12. Koroleva OA, Davies A, Deeken R, Thorpe MR, Tomos AD, Hedrich R: **Identification of a new glucosinolate-rich cell type in Arabidopsis flower stalk**. *Plant Physiol.* 2000, **124**:599-608.
13. Koroleva OA, Cramer R: **Single-cell proteomic analysis of glucosinolate-rich S-cells in Arabidopsis thaliana**. *Methods* 2011, **54**:413-423.
14. Sarsby J, Towers MW, Stain C, Cramer R, Koroleva OA: **Mass spectrometry imaging of glucosinolates in Arabidopsis flowers and siliques**. *Phytochemistry* 2012, **77**:110-118.
15. Kissen R, Rossiter JT, Bones AM: **The "mustard oil bomb": not so easy to assemble?! Localization, expression and distribution of the components of the myrosinase enzyme system**. *Phytochem. Rev.* 2009, **8**:69-86.
16. Petersen BL, Chen S, Hansen CH, Olsen CE, Halkier BA: **Composition and content of glucosinolates in developing Arabidopsis thaliana**. *Planta* 2002, **214**:562-571.
17. Magrath R, Mithen R: **Maternal effects on the expression of individual aliphatic glucosinolates in seeds and seedlings of Brassica napus**. *Plant Breed.* 1993, **111**:249-252.
18. Nour-Eldin HH, Andersen TG, Burow M, Madsen SR, Jørgensen ME, Olsen CE, Dreyer I, Hedrich R, Geiger D, Halkier BA: **NRT/PTR transporters essential for allocation of glucosinolate defense compounds to seeds**. 2012.
19. Sønderby IE, Hansen BG, Bjarnholt N, Ticconi C, Halkier BA, Kliebenstein DJ: **A systems biology approach identifies a R2R3 MYB gene subfamily with distinct and overlapping functions in regulation of aliphatic glucosinolates**. *PLoS One* 2007, **2**:e1322.
20. Andersen TG, Nour-Eldin HH, Fuller VL, Olsen CE, Burow M, Halkier BA: **Integration of biosynthesis and long-distance transport establish organ-specific glucosinolate profiles in vegetative Arabidopsis**. *Plant Cell* 2013, **25**:3133-3145.
21. Madsen SR, Olsen CE, Nour-Eldin HH, Halkier BA: **Elucidating the role of transport processes in leaf glucosinolate distribution**. *Plant Physiol.* 2014, **166**:1450-1462.
22. Barth C, Jander G: **Arabidopsis myrosinases TGG1 and TGG2 have redundant function in glucosinolate breakdown and insect defense**. *Plant J.* 2006, **46**:549-562.
23. Shroff R, Schramm K, Jeschke V, Nemes P, Vertes A, Gershenzon J, Svato A: **Quantification of plant surface metabolites by matrix-assisted laser desorption-ionization mass spectrometry imaging: glucosinolates on Arabidopsis thaliana leaves**. *Plant J.* 2015, **81**:961-972.
- A mass spectrometry protocol was developed to detect surface glucosinolates on Arabidopsis leaves by applying the MALDI matrix through sublimation. The results were confirmed by two other mass spectrometry-based techniques, laser ablation electrospray ionization and liquid extraction surface analysis.
24. Sønderby IE, Burow M, Rowe HC, Kliebenstein DJ, Halkier BA: **A complex interplay of three R2R3 MYB transcription factors determines the profile of aliphatic glucosinolates in Arabidopsis**. *Plant Physiol.* 2010, **153**:348-363.
25. Xu D, Hanschen FS, Witzel K, Nintemann SJ, Nour-Eldin HH, Schreiner M, Halkier BA: **Rhizosecretion of stele-synthesized glucosinolates and their catabolites requires GTR-mediated import in Arabidopsis**. *J. Exp. Bot.* 2016 <http://dx.doi.org/10.1093/jxb/erw355>.
26. Strehmel N, Böttcher C, Schmidt S, Scheel D: **Profiling of secondary metabolites in root exudates of Arabidopsis thaliana**. *Phytochemistry* 2014, **108**:35-46.
27. Wadhwa K, Narula N: **A novel technique to collect root exudates from mustard (Brassica juncea)**. *J. Basic Microbiol.* 2012, **52**:613-615.
28. Schreiner M, Krumbein A, Knorr D, Smetanska I: **Enhanced glucosinolates in root exudates of Brassica rapa ssp. rapa mediated by salicylic acid and methyl jasmonate**. *J. Agric. Food Chem.* 2011, **59**:1400-1405.
29. Matthiessen J, Kirkegaard J: **Biofumigation and enhanced biodegradation: opportunity and challenge in soilborne pest and disease management**. *CRC Crit. Rev. Plant Sci.* 2006, **25**:235-265.
30. Sherameti I, Venus Y, Drzewiecki C, Tripathi S, Dan VM, Nitz I, Varma A, Grundler FM, Oelmüller R: **PYK10, a beta-glucosidase located in the endoplasmic reticulum, is crucial for the beneficial interaction between Arabidopsis thaliana and the endophytic fungus Piriformospora indica**. *Plant J.* 2008, **54**:428-439.
31. Clay NK, Adio AM, Denoux C, Jander G, Ausubel FM: **Glucosinolate metabolites required for an Arabidopsis innate immune response**. *Science* 2009, **323**:95-101.
32. Bednarek P, Pislewska-Bednarek M, Svatos A, Schneider B, Doubsky J, Mansurova M, Humphry M, Consonni C, Panstruga R, Sanchez-Vallet A et al.: **A glucosinolate metabolism pathway in**

- living plant cells mediates broad-spectrum antifungal defense. *Science* 2009, **323**:101-106.**
33. Gigolashvili T, Berger B, Mock H-P, Müller C, Weisshaar B, Flügge U-I: **The transcription factor HIG1/MYB51 regulates indolic glucosinolate biosynthesis in *Arabidopsis thaliana*.** *Plant J.* 2007, **50**:886-901.
  34. Fuchs R, Kopischke M, Klapprodt C, Hause G, Meyer AJ, Schwarzländer M, Fricker MD, Lipka V: **Immobilized subpopulations of leaf epidermal mitochondria mediate PENETRATION2-dependent pathogen entry control in *Arabidopsis*.** *Plant Cell* 2016, **28**:130-145.
  35. Li L, Sheen J: **Dynamic and diverse sugar signaling.** *Curr. Opin. Plant Biol.* 2016, **33**:116-125.
  36. Griffiths CA, Paul MJ, Foyer CH: **Metabolite transport and associated sugar signalling systems underpinning source/sink interactions.** *Biochim. Biophys. Acta* 2016, **1857**:1715-1725.
  37. Andersen TG, Halkier BA: **Upon bolting the GTR1 and GTR2 transporters mediate transport of glucosinolates to the inflorescence rather than roots.** *Plant Signal. Behav.* 2014, **9**: e27740.
  38. Boege K, Marquis RJ: **Facing herbivory as you grow up: the ontogeny of resistance in plants.** *Trends Ecol. Evol. (Amst.)* 2005, **20**:441-448.
  39. Robert-Seilaniantz A, Grant M, Jones JDG: **Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism.** *Annu. Rev. Phytopathol.* 2011, **49**:317-343.
  40. Kliebenstein DJ: **False idolatry of the mythical growth versus immunity tradeoff in molecular systems plant pathology.** *Physiol. Mol. Plant Pathol.* 2016, **95**:55-59.
  41. Aarabi F, Kusajima M, Tohge T, Konishi T, Gigolashvili T, Takamune M, Sasazaki Y, Watanabe M, Nakashita H, Fernie AR *et al.*: **Sulfur deficiency-induced repressor proteins optimize glucosinolate biosynthesis in plants.** *Sci. Adv.* 2016, **2**: e1601087.
- The identification of the sulfur deficiency responsive genes SDI1 and SDI2 as negative regulators of glucosinolate biosynthesis provided a molecular mechanism balancing general and specialized sulfur metabolism under sulfur-deprived conditions.
42. Jensen LM, Jepsen HSK, Halkier BA, Kliebenstein DJ, Burow M: **Natural variation in cross-talk between glucosinolates and onset of flowering in *Arabidopsis*.** *Front. Plant Sci.* 2015, **6**:697.
  43. Kerwin RE, Jimenez-Gomez JM, Fulop D, Harmer SL, Maloof JN, Kliebenstein DJ: **Network quantitative trait loci mapping of circadian clock outputs identifies metabolic pathway-to-clock linkages in *Arabidopsis*.** *Plant Cell* 2011, **23**:471-485.
  44. Burow M, Atwell S, Francisco M, Kerwin RE, Halkier BA, Kliebenstein DJ: **The Glucosinolate biosynthetic gene AOP2 mediates feed-back regulation of jasmonic acid signaling in *Arabidopsis*.** *Mol. Plant* 2015, **8**:1201-1212.
- Analysis of mutants altered in the glucosinolate-modifying enzyme AOP2 showed an extensive array of effects upon the plant in the absence of an attacker illustrating that the glucosinolate defenses are not dead end products but functionally influence general processes in the plant.
45. Maharjan PM, Dilkes BP, Fujioka S, Pěnčík A, Ljung K, Burow M, Halkier BA, Choe S: ***Arabidopsis gulliver1/SUPERROOT2-7 identifies a metabolic basis for auxin and brassinosteroid synergy.*** *Plant J.* 2014, **80**:797-808.
  46. Francisco M, Joseph B, Caligagan H, Li B, Corwin JA, Lin C, Kerwin RE, Burow M, Kliebenstein DJ: **The defense metabolite, allyl glucosinolate, modulates *Arabidopsis thaliana* biomass dependent upon the endogenous glucosinolate pathway.** *Front. Plant Sci.* 2016, **7**:774.
- This study identified one specific glucosinolate, allyl glucosinolate, as a feedback regulatory compound that can modulate upstream regulatory networks and thereby alter plant biomass.
47. Francisco M, Joseph B, Caligagan H, Li B, Corwin JA, Lin C, Kerwin RE, Burow M, Kliebenstein DJ: **Genome wide association mapping in *Arabidopsis thaliana* identifies novel genes involved in linking allyl glucosinolate to altered biomass and defense.** *Front. Plant Sci.* 2016, **7**:1010.
  48. Bass J, Lazar MA: **Circadian time signatures of fitness and disease.** *Science* 2016, **354**:994-999.
  49. Harmer SL, Hogenesch JB, Straume M, Chang HS, Han B, Zhu T, Wang X, Kreps JA, Kay SA: **Orchestrated transcription of key pathways in *Arabidopsis* by the circadian clock.** *Science* 2000, **290**:2110-2113.
  50. Sanchez SE, Kay SA: **The plant circadian clock: from a simple timekeeper to a complex developmental manager.** *Cold Spring Harb. Perspect. Biol.* 2016, **8**.
  51. Seo PJ, Mas P: **STRESSing the role of the plant circadian clock.** *Trends Plant Sci.* 2015, **20**:230-237.
  52. Goodspeed D, Chehab EW, Covington MF, Braam J: **Circadian control of jasmonates and salicylates: the clock role in plant defense.** *Plant Signal. Behav.* 2013, **8**:e23123.
  53. Sharma M, Bhatt D: **The circadian clock and defence signalling in plants.** *Mol. Plant Pathol.* 2015, **16**:210-218.
  54. Goodspeed D, Chehab EW, Min-Venditti A, Braam J, Covington MF: ***Arabidopsis* synchronizes jasmonate-mediated defense with insect circadian behavior.** *Proc. Natl. Acad. Sci. U. S. A.* 2012, **109**:4674-4677.
  55. Wang W, Barnaby JY, Tada Y, Li H, Tör M, Caldelari D, Lee D, Fu X-D, Dong X: **Timing of plant immune responses by a central circadian regulator.** *Nature* 2011, **470**:110-114.
  56. Rosa EAS, Rodrigues PMF: **The effect of light and temperature on glucosinolate concentration in the leaves and roots of cabbage seedlings.** *J. Sci. Food Agric.* 1998, **78**:208-212.
  57. Liu JD, Goodspeed D, Sheng Z, Li B, Yang Y, Kliebenstein DJ, Braam J: **Keeping the rhythm: light/dark cycles during postharvest storage preserve the tissue integrity and nutritional content of leafy plants.** *BMC Plant Biol.* 2015, **15**:92.
- This study shows that postharvest storage under day-mimicking conditions maintains and sometimes even improves nutritional quality.